

PILLE MÄND

Light use capacity and carbon and nitrogen budget of plants: remote assessment and physiological determinants



DISSERTATIONES BIOLOGICAE UNIVERSITATIS TARTUENSIS

245

PILLE MÄND

Light use capacity and carbon and nitrogen budget of plants: remote assessment and physiological determinants



Department of Botany, Institute of Ecology and Earth Sciences, Faculty of Science and Technology, University of Tartu, Estonia

Dissertation was accepted for the commencement of the degree of *Doctor philosophiae* in plant ecology and ecophysiology at the University of Tartu on June 17, 2013 by the Scientific Council of the Institute of Ecology and Earth Sciences, University of Tartu.

Supervisors: Professor Olevi Kull (22 June 1955 – 31 January 2007), University of Tartu, Estonia
Professor Josep Peñuelas, CSIC, Global Ecology Unit CREAL-CEAB-CSIC-UAB, Catalonia, Spain
Dr. Anu Sõber, University of Tartu, Estonia

Opponent: Dr. Albert Porcar-Castell, University of Helsinki, Finland

Commencement: Room 218, 40 Lai Street, Tartu, on 11 October 2013 at 12:15 a.m.

Publication of this thesis is granted by the Institute of Ecology and Earth Sciences, University of Tartu and by the Doctoral School of Earth Sciences and Ecology created under the auspices of European Social Fund.



European Union
European Social Fund



Investing in your future

ISSN 1024–6479

ISBN 978–9949–32–379–1 (Print)

ISBN 978–9949–32–380–7 (PDF)

Copyright: Pille Mänd, 2013

University of Tartu Press
www.tyk.ee

CONTENTS

LIST OF ORIGINAL PUBLICATIONS	6
LIST OF TERMS AND ABBREVIATIONS	7
1. INTRODUCTION.....	10
1.1. Carbon and nitrogen budget of ecosystems	10
1.2. Remote sensing of plant productivity	11
1.3. The effect of canopy structure on estimations of light use efficiency	12
1.4. Within-leaf mechanisms influencing light-use efficiency	13
1.5. Aims of the thesis	15
2. MATERIALS AND METHODS.....	17
2.1. Study sites and experimental set-up	17
2.2. Data processing	19
3. RESULTS AND DISCUSSION.....	22
3.1. Primary production of shrublands	22
3.2. Green LAI of shrubland communities, estimated by NDVI.....	23
3.3. LUE of shrubland communities, estimated by PRI	24
3.4. Effect of canopy structure on light use efficiency of canopies and single leaves.....	26
3.5. Chlorophyll distribution within leaves and canopy	27
3.6. Nitrogen distribution within leaves and canopy	31
3.7. Nitrogen budget in relation with carbon production of vegetation	33
4. CONCLUSIONS	35
5. REFERENCES	37
SUMMARY IN ESTONIAN	46
ACKNOWLEDGEMENTS	50
PUBLICATIONS.....	53
CURRICULUM VITAE	113

LIST OF ORIGINAL PUBLICATIONS

This thesis is based on the following papers which are referred to in the text by Roman numerals:

- I** Beier C, Emmett BA, Tietema A, Schmidt IK, Peñuelas J, Kovács Láng E, Duce P, De Angelis P, Gorissen A, Estiarte M, de Dato GD, Sowerby A, Kröel-Dulay G, Lellei-Kovács E, Kull O, **Mänd P**, Petersen H, Gjelstrup P, Spano D. 2009. Carbon and nitrogen balances for 6 shrublands across Europe. *Global Biogeochemical Cycles* 23: 1–13.
- II** **Mänd P**, Hallik L, Peñuelas J, Nilson T, Duce P, Emmett BA, Beier C, Estiarte M, Garadnai J, Kalapos T, Schmidt IK, Kovács-Láng E, Prieto P, Tietema A, Westerveld JW, Kull O. 2010. Responses of the reflectance indices PRI and NDVI to experimental warming and drought in European shrublands along a north–south climatic gradient. *Remote Sensing of Environment* 114: 626–636.
- III** **Mänd P**, Hallik L, Peñuelas J, Kull O. 2013. Electron transport efficiency at opposite leaf sides: effect of vertical distribution of leaf angle, structure, chlorophyll content and species in a forest canopy. *Tree Physiology* 33: 202–210.
- IV** Eichelmann H, Oja V, Rasulov B, Padu E, Bichele I, Pettai H, **Mänd P**, Kull O, Laisk A. 2005. Adjustment of leaf photosynthesis to shade in a natural canopy: reallocation of nitrogen. *Plant, Cell and Environment* 28: 389–401.

Published papers are reproduced with due permission from the publishers.

The participation of the author in preparing the listed publications is as follows:

- Paper I – collected and analysed part of the data and participated in writing;
- Paper II – collected and analysed the data and was leading author in writing the paper;
- Paper III – collected and analysed the data and was leading author in writing the paper;
- Paper IV – collected and analysed part of the data and participated in writing.

LIST OF TERMS AND ABBREVIATIONS

abaxial side of leaf	the lower side (usually) of a bilateral leaf, which is often exposed to lower levels of light than leaf adaxial side
adaxial side of leaf	the upper side (usually) of a bilateral leaf, which is often exposed to higher levels of light than leaf abaxial side
APAR	annually absorbed photosynthetically active radiation ($\text{MJ m}^{-2} \text{yr}^{-1}$)
carboxylation capacity	the maximum rate of CO_2 fixation of photosynthesis ($\mu\text{mol m}^{-2} \text{s}^{-1}$)
Chl a/b	the ratio of concentrations of chlorophyll a to chlorophyll b
Chl b	chlorophyll b
chlorophyll fluorescence	fluorescent light that has been re-emitted after being absorbed by the chlorophyll molecules of a plant
Chl_s	chlorophyll content per unit leaf area ($\mu\text{mol m}^{-2}$)
fAPAR	the fraction of photosynthetically active radiation absorbed by vegetation
far-red irradiation	the red end of the visible light spectrum, wavelength region of 700–800 nm
fluorescence	the emission of light by a substance that has absorbed light, here used in the meaning of chlorophyll fluorescence
F_v/F_m	quantum yield of PSII of dark adapted (relaxed) leaves, calculated as the proportion of absorbed quanta, used in photochemical reactions
GPP	gross primary production, calculated as annual uptake of carbon by vegetation per unit ground area ($\text{g C m}^{-2} \text{yr}^{-1}$)
green LAI	leaf area index of green plant parts, calculated as leaf area per unit ground area ($\text{m}^2 \text{m}^{-2}$)
green LAI_e	effective projected green leaf area, calculated as the average number of contacts of inserted pin (pinpoint method) with green plant parts (leaves and green branches)
LAI	leaf area index, calculated as leaf area per unit ground area ($\text{m}^2 \text{m}^{-2}$)
LAI_e	effective projected leaf area, calculated as average number of contacts of inserted pin (pinpoint method) with plant parts
LHC	light harvesting complex, a complex of proteins and photosynthetic pigments focusing light energy to the reaction centre of the photosystem

LHCI	LHC focusing light energy to the reaction centre of PS I
LHCII	LHC focusing light energy to the reaction centre of PS II
LHCII trimer	large trimeric subunit of LHCII
LMA	leaf dry mass per unit leaf area (g m^{-2})
LUE	light use efficiency, defined as the amount of carbon in produced biomass per unit of absorbed light (g C MJ^{-1})
MODIS	Moderate Resolution Imaging Spectroradiometer acquiring daily reflectance data of Earth's surface
NDVI	Normalized Difference Vegetation Index, acquired from vegetation reflectance at the visible and far-red spectral regions
NDVI₅₇₀	NDVI, calculated from reflectance at 570 and 780 nm
NDVI₆₈₀	NDVI, calculated from reflectance at 680 and 780 nm
N_m	leaf nitrogen content per unit dry mass (g g^{-1})
Non-photochemical quenching	mechanism in plants for dissipating excess light energy as heat
NPP	net primary production, calculated as annual production of carbon per unit ground area when carbon cost of respiration is subtracted ($\text{g C m}^{-2} \text{yr}^{-1}$)
NPQ	non-photochemical quenching, calculated from chlorophyll fluorescence
N_s	leaf nitrogen content per unit leaf area (g m^{-2})
PAR	photosynthetically active radiation ($\text{MJ m}^{-2} \text{yr}^{-1}$)
photosystem	protein complex that carry out the primary absorption of light and the transfer of energy and electrons during photosynthesis
PPFD	photosynthetic photon flux density, the number of photons in the 400–700 nm waveband incident per unit time on a unit surface ($\mu\text{mol photons m}^{-2} \text{s}^{-1}$)
PRI	Photochemical Reflectance Index, calculated using reflectance at wavelengths of 531 and 570 nm
PSI	a protein complex that captures and uses light energy to mediate electron transfer from plastocyanin to ferredoxin
PSI antenna	complex of proteins and photosynthetic pigments focusing light energy toward the reaction centre of PS I
PSII	a protein complex that carries out the primary absorption of light in photosynthesis and uses light energy to energize electrons that are then transferred through a variety of coenzymes and cofactors to reduce plastoquinone to plastoquinol

qN	non-photochemical quenching, calculated on a scale of 0–1 from chlorophyll fluorescence
qP	photochemical quenching, an approximation of the proportion of PSII reaction centres that are open, calculated from chlorophyll fluorescence
quantum use efficiency	the proportion of absorbed light that is used in photochemical reactions
red-irradiation	the wavelength region of 600–700 nm of the visible light spectrum
Rubisco	ribulose-1,5-bisphosphate carboxylase-oxygenase, an enzyme catalysing the fixation of CO ₂ during photosynthesis
Φ_{II}	quantum yield of PSII of light adapted leaves, calculated as the proportion of light absorbed by PSII that is used in photochemical reactions

I. INTRODUCTION

I.1. Carbon and nitrogen budget of ecosystems

Understanding the determinants of the global carbon cycle is becoming a more and more important task of plant ecophysiology due to predictions of large-scale changes in global climate (IPCC, 2007). Roughly 50 Gt of carbon is annually acquired through photosynthesis just by terrestrial plants (Potter, 2012) and even small changes in climate may cause substantial changes in local scale plant production (Parmesan and Yohe 2003, Peñuelas et al. 2007, Bokhorst et al. 2011). The relationships between vegetation production and irradiance, moisture, temperature and nitrogen, as major limiters of vegetation carbon assimilation, have been studied over different biomes and functional groups and extrapolated to the global scale (e.g. Field et al. 1992, Potter et al. 1993, White et al. 2000, Caylor et al. 2003, Wright et al. 2004, Yuan et al. 2007, Street et al. 2012, Peltoniemi et al. 2012). However, compared to abundant studies on forests and grasslands, very few papers on the carbon balances of shrublands are available, although this data is also needed for the determination of global carbon budgets.

In parallel with studies on carbon budget of different ecosystems, more and more attention is paid also to nitrogen content of plants. Already in 1987 Hirose and Weger demonstrated that nitrogen distribution has a strong effect on daily canopy carbon gain, and plants from a denser stand should benefit more from the nonuniform distribution of nitrogen in the canopy (Hirose 2005). Therefore it was assumed that the modelling of canopy photosynthesis with leaf nitrogen distribution enables us to scale up from chloroplast biochemistry to canopy carbon gain (Kull and Jarvis, 1995, De Pury and Farquhar, 1997). Recently canopy-scale measurements have also proven the significant effect of the nitrogen budget of ecosystems on canopy carboxylation capacity, respiration and soil turnover rates, all of which control the canopy carbon cycle and influence the sensitivity of the global carbon cycle to changes in climate (Lambers et al. 2008, Ollinger et al. 2008). Thus C/N ratios of different functional types of plants are often incorporated into plant production models (White et al. 2000, Wania et al. 2012). At the same time the whole concept of nitrogen assimilation by plants has changed (Schimel and Bennett 2004) more efficiently explaining the link between carbon and nitrogen budgets. Nevertheless, there are still large disagreements in calculations of global scale nitrogen balance (Zaehle and Dalmonch 2011, Wania et al. 2012) partly due to changes in C/N ratio in plants as a result of changing climate (Sardans et al. 2012) and partly because of a lack of knowledge on the nitrogen cycles of different ecosystems and plant functional types.

Today the main obstacle for modelling global vegetation processes lies in the identification of a minimal number of plant functional types, in order to reduce the complexity of traits that are needed for productivity predictions (Lavorel et al. 2008). On the other hand, to be able to select those relevant traits and plant groups we need enough knowledge on the whole range of natural variability in

different light-capture and light-use parameters and strategies, which determine the productivity. Thus, two different approaches are used for modelling global productivity: 'top-down' models using remotely sensed data integrate both physiological and structural vegetation parameters, and 'bottom-up' models scale productivity to the canopy level from the known chloroplast- and leaf-level physiology and its relations to resource availability (Houborg et al. 2009).

1.2. Remote sensing of plant productivity

Often the 'top-down' approaches are based on interpretations of remotely sensed canopy reflectance signals (Hilker et al. 2008). To detect changes in global plant cover, remote sensing technologies are constantly developing and there is a search for the best remotely estimated fit for key parameters of productivity (Monteith and Moss 1977): leaf area index (LAI), the fraction of photosynthetically active radiation absorbed by vegetation ($fAPAR$) and the light use efficiency (LUE). Reflectance index NDVI (Rouse et al. 1974) is one remote sensing parameter that is often used as an estimate of the fraction of photosynthetically active radiation absorbed by vegetation ($fAPAR$) (Huemmrich et al. 2010, Peng et al. 2012, Lausch 2013) and a closely related parameter, green leaf area index (green LAI) (Fensholt et al. 2004, Haboudane et al. 2004). The estimations of $fAPAR$ and LAI have already been improved over a number of years and the best outcome is that based on MODIS algorithm global LAI and $fAPAR$ data at a 1×1 km spatial resolution is collected on regular basis (Myneni et al. 2002). However, MODIS sorts the vegetation within each pixel to one of six given biomes leaving room for misclassifications and similarly to small-scale NDVI measurements, at dense canopies MODIS LAI tends to saturate (Myneni et al. 2002). Differently from $fAPAR$ and LAI, estimations of LUE have been improved most extensively over the last decade as spectrometers with better resolution became available.

In general two different approaches for determining LUE from a distance are used: the first option is the prediction of LUE indirectly from environmental stresses, the other possibility is to predict LUE more-or-less directly from changes in spectral reflectance (Hilker et al. 2008). The environmental-stress based LUE estimations using narrow waveband reflectance are highly sensitive to atmospheric scattering and direction of measurements (Liu et al. 2006), yet recently hyperspectral imagery has quite successfully been used for prediction of foliage nitrogen content from narrow waveband reflectance (Martin et al. 2008, Ollinger et al. 2008). The MODIS project also gives estimates of GPP, which use biome-specific information about stress-induced changes in maximum LUE (Heinsch et al. 2003), even though several studies have reported LUE to vary not only between biomes but also between different functional groups of species (Nichol et al. 2000, Ahl et al. 2004, Kergoat et al. 2008).

The alternative approach is estimating LUE more directly from spectral reflectance, for instance by measuring solar induced chlorophyll fluorescence emissions of canopies, or more precisely the impact of fluorescence emissions on apparent spectral reflectance (Zarco-Tejada et al. 2000, Moya et al. 2004). However, this approach has been technically tricky since right now, remote detection of chlorophyll fluorescence is possible only by using sub-nanometre reflectance bands in the red and near infrared regions where solar radiation is not abundant (Meroni and Colombo 2006). Concurrently remotely sensed changes in the photoprotective mechanism are also used for estimations of canopy LUE, since a reflectance index PRI has been proposed as an optical proxy of light use efficiency (Garbulsky et al. 2011). Using a reflectance band at 531 nm, reflectance index PRI involves combined information on changes in the xanthophyll cycle and the aggregation state of PSII antennas (Gamon et al. 1992, Peñuelas et al. 1995, Gamon et al. 1997). For measurements from space the relationship between LUE and PRI is better if an atmospherically corrected band of 678 nm is used for PRI calculations, and PRI measurements from space seem to improve MODIS LUE derivations immensely when compared to the previously-used biome-specific look-up table of maximum LUEs (Drolet et al. 2008). Yet, the relationship between PRI and LUE seems to be species-specific, much less variance in LUE is described by PRI when different functional types of plants are investigated (Garbulsky et al. 2011). Also the reliance of PRI on canopy structure and light conditions has been reported (Barton and North 2001, Grace et al. 2007, Hilker et al. 2008). Thus the applicability of PRI for estimation of LUE of different ecosystem types with varying canopy structure still needs to be tested.

I.3. The effect of canopy structure on estimations of light use efficiency

Currently one of the most challenging aspects of remote sensing is tracking the photosynthetic activity of forests, since the remote estimation of carbon uptake by multilayered forest ecosystems involves several difficulties (Grace et al. 2007). The structure dependence of PRI values applies also to the vertical structure of vegetation (Nichol et al. 2000, Barton and North 2001, Damm et al. 2010). Forests, as very complex systems of coexisting species with different spatial and temporal strategies for light capture may as a whole system in fact be much more efficient in light use than simpler canopy-systems (Ishii and Asano 2010). For instance the productivity of the whole ecosystem appears to be higher in mixed forests, which consist of both broadleaved and coniferous species (Aiba et al. 2007). Mostly the 'top-down' models using reflectance data for estimations of light use efficiency, are very simple. Nevertheless simple LUE models may perform even better than more detailed 'bottom-up' scaled-leaf models containing several species-specific parameters (Houborg et al.

2009). It is difficult to obtain enough input data for multilayer models in order to derive total canopy-scale fluxes.

However, to understand the mechanistic basis of changes in leaf and canopy-level light use, more complex models are necessary. Multilayer models that consider the vertical variability of photosynthesis or at least divide canopy into shaded and sunlit fractions have proven to better estimate the diurnal changes in canopy photosynthesis (Chen et al. 1999, Baldocchi and Wilson, 2001, Damm et al. 2010). Among other factors that influence canopy photosynthesis, the efficiency with which leaf canopy is exposed to light is shown to be a function of inclination angles and spatial aggregation of leaves (Cescatti and Niinemets 2004, Niinemets 2010). However, simpler models assume the inclination angle distribution of plant canopies to be spherical, even if such a generalization may cause significant bias in the estimations of vertical distribution of light and photosynthesis within the canopy (Stadt and Lieffers 2000, Sarlikoti et al. 2011). This is because real vegetation canopies consist of a mixture of species with several alternative leaf inclination distributions (Niinemets 2010).

Another topic that needs to be studied more explicitly is the physiological relationship between leaf level chlorophyll fluorescence measurements and CO₂ assimilation, as there are indications that this relationship may change among leaves with different inclination angles and resultant light conditions (Myers et al. 1997, Tsuyama et al. 2003, Damm et al. 2010). This suggests that if estimations of leaf level quantum use efficiency derived from chlorophyll fluorescence are used for modelling canopy photosynthesis, the inclusion of leaf angle distribution into the model may be necessary. Most probably, leaf-level differences in light acclimation strategies also influence the upscaling from leaf-level quantum use efficiency to canopy light use efficiency (Damm et al. 2010).

I.4. Within-leaf mechanisms influencing light-use efficiency

In order to understand the factors influencing the up-scaling of photosynthetic properties from leaf to canopy the variations in leaf structure and the differences in light-acclimation of chloroplasts within the leaf must be considered. The leaf and chloroplast properties that define light use efficiency of the whole canopy have been studied profoundly (Kull 2002), but the magnitude of these light induced changes in leaf morphology and physiology is not known for different environmental conditions and for different species or functional groups (Ishii and Asano 2010). Valladares and Niinemets (2008) concluded that the magnitude of light acclimation of leaf morphology is larger in shade tolerant species, whereas high-light demanding species change their physiology more plastically. However, the morphology of shade tolerant species can not be the major means of optimizing the quantum use of sunfleck energy, which is one factor strongly influencing the carbon gain of lower canopy regions (Porcar-Castell and Palmroth 2012, Way and Pearcy 2012). Furthermore, the regulation of one trait, such

as leaf chlorophyll content, which has been shown to be a very sensitive indicator of different environmental stresses, can be achieved either through changes in the overall percentage of chlorophyll, through physiological redistribution of chlorophyll between different photosynthetic units, but also through changes in leaf morphology as changes in LMA with constant chlorophyll content may affect light capture efficiency significantly (Niinemets 2010).

The distribution of leaf photosynthetic traits through the canopy seems to be controlled not only on the leaf level but is evidently a whole-canopy phenomenon (Givnish 1988, Kull 2002). Indeed, Moreau et al. (2012) found that the relationship between canopy nitrogen gradient and the corresponding light gradient is a function of canopy size. Yet, changes in leaf nitrogen have proven to be of key importance in the acclimation of leaf photosynthesis to different light conditions (Hirose 2005). In fact, the photosynthetic performance of leaf and canopy is defined not only by within leaf and canopy nitrogen distribution, but also the magnitude of changes in different leaf traits depends on the availability of resources, such as nitrogen (Portsmuth and Niinemets 2007). The reason behind the nitrogen dependence of chlorophyll content and distribution is that chlorophylls need to be bound to proteins that are costly since they contain nitrogen (Evans 1989). The magnitude of light adjustment of the photosynthetic apparatus is not only restricted by the availability of resources (such as nitrogen), but there probably also exists a lower limit of acclimation, as certain constructional features define the minimum amount of nutrients for photosynthetic units to remain intact and active (Kull 2002, Meir et al. 2002).

The consideration of chloroplast-level light acclimation differences within the leaf have not been very common in ecological studies as many gaps are still present in the knowledge of the molecular-level adjustment of photosynthetic compounds to irradiance. Due to technical limitations the differentiation between photosynthetic subunits and the detection of the allometric redistribution of different components of the photosynthetic apparatus *in situ* in leaves from natural canopies has not been possible until recently. That is why the acclimation of the photosystem has remained relatively unnoticed until advances in state transition studies have stressed the variability in light harvesting abilities of PSI (Ruban et al. 2006, Ruban and Johnson 2009, Minagawa 2011). Traditionally a lot of attention has been paid to the light adjustment of photosystem II since in ecological research, leaf chlorophyll a/b ratio has often been observed to decrease with decreasing light availability and has been interpreted as a proportional increase in the chlorophyll b binding light harvesting complex of LHCII complexes (Evans 1989). However, studies on the molecular-scale light acclimation of photosystem II implicate multiple locations for chlorophyll b (Ruban et al. 2006, Kouřil et al. 2013).

In addition to light acclimation of photosystems, other nitrogen-rich units of the photosynthetic apparatus, such as ribulose-bisphosphate carboxylase-oxygenase (Rubisco), are also subjected to changes in light environment. It has been shown that in high-light species the investment of nitrogen into carboxylation enzymes, rather than into chlorophyll, increases photosynthetic

capacity more efficiently (Niinemets and Tenhunen 1997, Terashima et al. 2006). The inclusion of parameters considering light-acclimation-induced changes in the allocation of protein between different components of photosynthesis could improve the predictions of maximum photosynthetic capacity of photosynthesis models (Mott and Woodrow 2000, Porcar-Castell and Palmroth 2012). Nevertheless, too little is known about the natural variability in the amount of Rubisco and other units of photosynthetic apparatus across several species and plant functional groups in order to construct mechanistic models linking nitrogen allocation to electron transport, carboxylation, respiration, storage and light use efficiency (Xu et al. 2012).

I.5. Aims of the thesis

The general aim of this study was to investigate the factors that influence primary production of vegetation and remote estimations of primary production, focusing on the most variable component of primary production – light use efficiency. In particular, we studied which factors are most responsive to changes in leaf and canopy light environment and how do these differences in canopy, plant and leaf architecture and physiology influence optical estimations of leaf- and canopy-level light use capacity and the green leaf area index (green LAI). First, based on previous knowledge of the dependence of remote estimations of vegetation physiology on the wavelength region from where optical signals are obtained, we assumed that measurements from the region of maximum absorption of chlorophyll might be worse estimators of plant properties than measurements from wavelength region where chlorophyll absorption is weaker. Secondly, we expected that canopy reflectance index PRI is well suited for estimating canopy light-use efficiency in European shrublands. We also assumed that the fraction of whole-leaf light capture efficiency that is described by optical measurements of chlorophyll fluorescence, changes within the vegetation canopy as a function of leaf angles, since differences in leaf orientation cause larger or smaller irradiance differences between alternate sides of a bilateral leaf. Additionally we expected that canopy and leaf nitrogen content adjusts more extensively to differences in light conditions than light-absorbing chlorophyll content.

The specific objectives of this thesis were:

1. To produce an integrated scheme of carbon and nitrogen pools and fluxes of European shrublands.
2. To estimate the applicability of reflectance indices NDVI (calculated from two different wavelength regions) and PRI in estimating green LAI and light use efficiency of different shrubland communities that are subjected to varying conditions of irradiance, precipitation and temperature.

3. To investigate the effect of canopy and leaf structure on remote estimations of canopy-level light-use efficiency and leaf-level quantum capture efficiency.
4. To search for differences in leaf chlorophyll distribution within leaves from different canopy positions and different species.
5. To study the natural variation in the fraction of photosynthetic and non-photosynthetic nitrogen in leaves from different irradiance conditions and different species.
6. To find out if canopy nitrogen content can be used for estimating differences in carbon production of aboveground vegetation of European shrublands.

2. MATERIALS AND METHODS

2.1. Study sites and experimental set-up

Field studies were carried out at two different spatial scales. A large-scale manipulation experiment of European shrublands differing in annual irradiation, moisture and temperature has been described in **I** and **II**. The steep vertical canopy light gradient of a multi-layered deciduous forest at Järvselja, Estonia, was investigated in **III** and **IV**.

Papers I and II

The experimental sites of six European shrublands were located in Wales-UK, Denmark, Netherlands, Hungary, Sardinia-Italy, and Catalonia-Spain where next to three control plots, three repetitions of warming and drought systems were installed in order to imitate changed climate. In warming experiment the vegetation was covered with reflective curtains at night: this decreased the loss of heat and increased soil temperature by 0–3 °C, depending on the site. For drought treatments, during precipitation the vegetation was covered with transparent waterproof covers for two-month periods within the growing season, reducing the precipitation of year 2003 by 39–349 mm, depending on the site. This experiment was part of EU projects CLIMOOR and VULCAN (Beier et al. 2004). The meteorological survey and measurements for canopy belowground and aboveground carbon and nitrogen pools and fluxes were performed during the years 1998–2004 for **I**. Optical measurements of **II** were carried out in summer 2003, in a period when drought treatment was being applied at each site. During both studies, air and soil temperature, precipitation and irradiance were constantly recorded by permanent sensors.

In **I** the aboveground vegetation structure and frequency (pinpoint measurements), carbon and nitrogen content of plant tissue and litter, litter decomposition rate, belowground root length, soil carbon and nitrogen content at main rooting depth and in mineral soil, the carbon content of microbial biomass, soil respiration, concentrations of NO_3 and NH_4 in soil, NO_3 and dissolved organic carbon (DOC) concentration in soil water were measured in order to calculate plant biomass, root litterfall, the decomposition of soil organic matter, the fraction of autotrophic and heterotrophic soil respiration, nitrogen mineralization rate, soil water balance, leaching losses of NO_3 and DOC and eventually the carbon and nitrogen pools and fluxes of different shrubland ecosystems.

In **II** canopy reflectance, chlorophyll fluorescence, concentration of leaf chlorophyll, leaf mass per area (LMA) and leaf nitrogen content were measured. Canopy reflectance measurements and part of the leaf-level fluorescence measurements of **II** were performed on the same subplots as measurements of canopy structure (pinpoint measurements) of **I**. Canopy reflectance was measured using a ground-based S2000-FL spectrometer (Ocean Optics Inc., Dunedin, FL, USA) from the spectral region between 400 nm and 950 nm at 1 m height

above the canopy, and vegetation indices PRI and NDVI₆₈₀ and NDVI₅₇₀ were calculated. See more details in **I** and **II**.

Papers III and IV

The experiments were performed in Estonia at Järvelja Experimental forest (58°22'N, 27°20'E) in the summers of 1999–2002. Permanent scaffolding towers were used to access the topmost and lowest canopy layers of the studied tree species: shade intolerant *Betula pendula* Roth. and *Populus tremula* L. and shade tolerant subcanopy *Tilia cordata* Mill. In **IV**, a herbaceous species *Solidago virgaurea* was also taken under examination. Above each measured leaf light conditions were estimated by using hemispherical photography.

In **III**, chlorophyll fluorescence from the adaxial and abaxial side of the leaf, leaf concentrations of chlorophyll a and b, LMA, and leaf angles were measured. Leaf adaxial/abaxial fluorescence ratios were used as estimations of varying intraleaf light acclimation profiles of chloroplasts. In the following year, leaf reflectance from the wavelength range of 655–665 nm (red spectral region) and 550–560 nm (green spectral region), LMA and leaf chlorophyll content and nitrogen content were measured at the same canopy heights of the same trees where the fluorescence measurements of previous year had been carried out. We measured leaf transmittance and reflectance with an integrating sphere (ISP-80-8-R, Ocean Optics) and fibre optic spectrophotometer (S2000, Ocean Optics).

In **IV**, shoots of *B. pendula* and *T. cordata* were cut and immediately placed into water, while the herbaceous species *S. virgaurea* was dug up and planted into pots with moist soil. For non-destructive measurements of partial reactions of photosynthesis a combined gas exchange/optical system was used in order to simultaneously measure quantum flux densities and chamber CO₂ and O₂ concentrations and to record CO₂ uptake, O₂ evolution, chlorophyll fluorescence, and 820 nm absorptance (Laisk et al. 2002, Eichelmann et al. 2004). In addition we measured chlorophyll a and b content, concentration of proteins (including Rubisco), leaf dry weight and area and leaf nitrogen concentration. Data was used for calculations of PSII quantum capture efficiency and PSII e⁻ transport rate, relative optical cross-section of PSII, relative absorption cross-section of PSI, densities of PSII and PSI and antenna sizes of PSII and PSI and the mass of nitrogen in different photosynthetic compartments. See more details in **III** and **IV**.

Leaf chlorophyll fluorescence of was measured using modulated chlorophyll fluorescence equipment PAM-2000 (WalzGmbH, Effeltrich, Germany) at both study-sites. See more details in **II–IV**.

2.2. Data processing

Total ecosystem C balance ($\Delta_{C\text{-System}}$) at **I** was calculated as:

$$\Delta_{C\text{-System}} = \text{Dep}_{\text{DOC}} + \text{NPP} - \text{Soilresp}_{\text{Het}} - \text{C} - \text{Leach}_{\text{DOC}} \quad (1)$$

where Dep_{DOC} is deposition of carbon into soil as dissolved organic carbon ($\text{g C m}^{-2} \text{y}^{-1}$), NPP is net uptake of carbon by plants ($\text{g C m}^{-2} \text{y}^{-1}$), $\text{Soilresp}_{\text{Het}}$ is carbon used in heterotrophic soil respiration ($\text{g C m}^{-2} \text{y}^{-1}$), $\text{Leach}_{\text{DOC}}$ accounts for losses in dissolved organic carbon through leaching ($\text{g C m}^{-2} \text{y}^{-1}$). See more details in **I**.

Effective projected leaf area (green LAI_e) of shrublands at **II** was calculated as the average number of contacts with green parts (leaves and green branches) of the plants per pin. See further details in **II**.

The vegetation reflectance indices at **II** were calculated as follows:

$$\text{NDVI}_{680} = (\text{R}780 - \text{R}680) / (\text{R}780 + \text{R}680) \quad (2)$$

$$\text{NDVI}_{570} = (\text{R}780 - \text{R}570) / (\text{R}780 + \text{R}570) \quad (3)$$

$$\text{PRI} = (\text{R}531 - \text{R}570) / (\text{R}570 + \text{R}531) \quad (4)$$

R_x in the equations is the reflectance at x nm. For more details of canopy reflectance measurements see details in **II**.

Leaf-level reflectance parameters for alternate leaf sides of deciduous trees at **III** were averaged from the red spectral region (655–665 nm) and the green spectral region (550–560 nm). Sample absorption at the same wavelength ranges was calculated from the equation:

$$\text{absorbance} = (1 - \text{reflectance} - \text{transmittance}) \quad (5)$$

Leaf-level chlorophyll fluorescence parameters, such as quantum yield of PSII of dark-adapted leaves (F_v/F_m) and light-adapted leaves (Φ_{II}), photochemical quenching (qP) and non-photochemical quenching (NPQ and qN) in **II–IV** were calculated from equations:

$$\Phi_{II} = (F_m' - F) / F_m' \quad (6)$$

$$F_v/F_m = (F_m - F_o) / F_m \quad (7)$$

$$qP = (F_m' - F) / (F_m' - F_o') \quad (8)$$

$$\text{NPQ} = (F_m - F_m') / F_m' \quad (9)$$

$$qN = (F_m - F_m') / (F_m - F_o) \quad (10)$$

where F_o is minimum fluorescence yield and F_m maximal fluorescence yield of dark-adapted leaves. F_m' is maximal fluorescence yield and F_o' minimum fluorescence yield of light-adapted leaves. See further details in **II–IV**.

The relative absorption cross-section of PSII (a_{II}) and PSI (a_I) and antenna sizes of PSII (PSUII) and PSI (PSUI) in **IV** were calculated as follows:

$$a_{II} = Yc/\Phi_{II} \quad (11)$$

$$a_I = Yc/Y_p \quad (12)$$

$$Y_p = P_o - P_s/P_m \quad (13)$$

$$PSUI = (a_I/a_I + a_{II})Chl/N_I \quad (14)$$

$$PSUII = (a_{II}/a_I + a_{II})Chl/N_{II} \quad (15)$$

where Yc is quantum yield of linear electron transport, calculated from the photosynthetic carbon metabolism, P_m is the transmittance signal at 820 nm that corresponds to totally oxidized PSI, P_o is the pulse-oxidizable fraction of PSI, P_s is the transmittance signal corresponding to the steady-state oxidation of PSI, N_I is the density of PSI, and N_{II} is the density of PSII. N_{II} was determined as four times the oxygen evolution from a saturating single-turnover flash. Two different kinetic methods were used for determining N_I , see **IV** for details. Chl is the fraction of total chlorophyll that is bound to a certain photosystem. Total Chl was assumed to be distributed between the photosystems proportionally with a_{II} and a_I . The mass of nitrogen in the cores of PSI and PSII in **IV** was calculated from the measured N_I and N_{II} .

For more information about the calculation of different plant parameters see details in **I–IV**.

Light use efficiency:

For this thesis we calculated the light use efficiency, LUE ($g\ C\ MJ^{-1}$), of six European shrublands from a simple equation (Monteith 1972, Monteith and Moss 1977):

$$LUE = NPP/APAR \quad (16)$$

$$APAR = PAR \times fAPAR \quad (17)$$

We used net primary production NPP ($kg\ C\ m^{-2}\ yr^{-1}$) estimations for different European shrublands, which were calculated as described in **I**. APAR was absorbed photosynthetically active radiation ($MJ\ m^{-2}\ yr^{-1}$), which was calculated from PAR (photosynthetically active radiation, $MJ\ m^{-2}\ yr^{-1}$) multiplied by green $fAPAR$ (fraction of PAR absorbed by green vegetation canopy). $fAPAR$ was calculated as a function of LAI (Ruimy et al. 1999):

$$fAPAR = 0.95 \times (1 - e^{-k \times LAI}) \quad (18)$$

For the light extinction coefficient (k) a default value of 0.5 (Campbell and Norman 1998) was used. LAI was measured by the pin-point method (Jonasson 1988, description at **II**) where we assumed LAI to be robustly half of the average number of contacts with a plant per pin (LAI_e):

$$LAI = 0.5 \times LAI_e \quad (19)$$

A different index for estimating LUE of shrublands was derived from a paper by Green et al. (2003), where the strongest predictor of LUE across all data appeared to be an index using nitrogen content of leaves:

$$\text{Index}_{\text{LUE}} = (\text{LAI} \times \text{N}_m / 100) / f\text{APAR} \quad (20)$$

where N_m was obtained from the average nitrogen content of the leaves of the dominant species.

Statistical analysis:

For statistical analysis we used STATISTICA software (StatSoft Inc. USA). Linear and nonlinear regression analysis were used in **I**, **II** and **IV** for analysing the effect of differences in climate and irradiance on mean ecosystem parameters of different study-sites, for estimating light use efficiency of leaves from reflectance parameters and for finding the best fit of relationships between different ecosystem, canopy and leaf parameters. Multiple regression analysis was performed to estimate different light use parameters from reflectance indices measured above plants in **II**. Pearson's correlation in **II** and **IV** or Spearman correlation coefficients in **III** (in the case of heteroscedasticity) were calculated between various leaf and canopy parameters. The normality and homoscedasticity of data was tested. Logarithmic transformation was performed if deviation from normal distribution was found. General linear models (GLM) were constructed in **II**, considering different categorical and continuous predictors. The squared component of a continuous factor was included in the models if the tested variable was nonlinearly related to a given factor. The significance of differences between separate groups was investigated using the Fisher LSD post-hoc test in **II** and **IV**. Non-parametrical Kruskal–Wallis ANOVA by ranks was used for analysing the data in **III**, due to heteroscedasticity. For testing the significance of differences between species in **III**, Wilcoxon signed rank test was used. In **IV** the significance of differences between shade and sun leaves was tested using the *t*-test for independent groups. See more details about statistical analysis in **I–IV**.

3. RESULTS AND DISCUSSION

3.1. Primary production of shrublands

In order to remotely estimate and model changes in shrubland production we first need data on a varying range of carbon storage rates/fluxes and biochemical cycles of different shrublands, since not many papers about the carbon budgets of shrublands have been published. Our studies at six different sites in Europe revealed that apparently similar shrubland ecosystems range from being carbon sinks in moist areas (system uptake $126 \text{ g C m}^{-2} \text{ y}^{-1}$ in Wales) to carbon sources in more arid regions (system emission up to $536 \text{ g C m}^{-2} \text{ y}^{-1}$ in non-steady-state Italian shrubland) (Fig. 4 in I).

Models of the global carbon cycle have demonstrated that in forest ecosystems, belowground carbon allocation is one of the largest fluxes of C aside from canopy assimilation (Davidson et al. 2002). Our study (I) revealed that steady-state shrubland systems appear to allocate even larger amounts of carbon belowground relative to litterfall (an average of 5:1) compared to most forest systems (2:1) as shown by Davidson et al. (2002). We also found that differences in carbon allocation of various European shrublands could not be explained by differences in annual temperature or precipitation, even if the non-steady-state shrubland of Italy was left out of the analysis (Fig. 1). Neither did mean annual radiation explain differences in carbon allocation (data not shown).




Figure 1. Ratio between belowground and aboveground carbon content related to (a) mean annual temperature and (b) mean annual precipitation at six European shrublands. High ratios in Italy result from the early successional stage of the shrubland after clearcut.

As a result, we expected that large belowground C fluxes of shrublands make it difficult to detect changes in the carbon balance of shrublands by remote-sensing tools, as this method deals only with parameters that can be measured aboveground. However, the aboveground carbon of different shrublands showed a significant exponential relationship to mean annual precipitation (Fig.

2b in **I**). Higher precipitation increased also the quantum absorbing area of vegetation, green LAI_e (Fig. 1a in **II**). Despite site-specific allocational differences in carbon pools, our measurements at five European shrublands suggest that at a large scale, aboveground carbon pools of shrublands are likely to be related (though according to five studied shrublands the relationship might be non-linear) to belowground carbon pools in steady-state shrublands (Fig. 1a in **I**), assuring that robust assumptions about belowground carbon pools can be made based on aboveground or above canopy measurements.

3.2. Green LAI of shrubland communities, estimated by NDVI

If we intend to estimate carbon pools and fluxes from a distance, we can not deal just with allocational differences, but must also consider that canopy structure and background properties affect our estimations (Barton and North 2001, Grace et al. 2007, Hall et al. 2008, Hilker et al. 2010). Our measurements of reflectance index NDVI₆₈₀ at six European shrublands revealed a positive relationship between NDVI and green LAI_e (effective green leaf area index). However this relationship tended to saturate at medium to high green LAI_e conditions (Fig. 4 in **II**, Table 4 in **II**). Similar levelling off of NDVI at dense vegetation has been reported in different vegetation types at multiple spatial scales (Gamon et al. 1995, Myneni et al. 2002, Haboudane et al. 2004, Lausch 2013). Thus, our result indicates that NDVI₆₈₀, which is calculated using red spectral region near maximum absorbance of chlorophyll, cannot be used for detecting changes in green LAI in dense shrublands.

In order to lessen the saturation effect of NDVI₆₈₀, we additionally calculated NDVI₅₇₀, which used green spectral region (less absorbed by chlorophyll). Although, at northern sites, relationships between NDVI₅₇₀ and green LAI_e or fluorescence parameters were slightly stronger than the same relationships with NDVI₆₈₀ (Table 5 in **II**), saturation of NDVI₅₇₀ still occurred in denser vegetation (Fig. 4 in **II**). However, a recent multi-scale study with different spectrometers gives hope that NDVI calculated from the green spectral region can also be used at higher LAI conditions if the methodology is improved (Lausch 2013).

While testing NDVI for detecting the effect of drought and night-time warming on shrubland vegetation, we found that generally lower values of NDVI were recorded in drought treatment plots (mean effect at Fig. 2 in **II**). On the other hand, if we searched for treatment effects at different study-sites separately, we found no significant drought effect on NDVI, and the warming treatment significantly increased NDVI only in the Italian site (Fig. 2 in **II**). As the treatments simulated realistic climate changes (being realistically small), the possible changes in green LAI_e were not large either and we found green LAI to be significantly affected by manipulations just in Denmark, where artificial drought significantly decreased green LAI_e (Fig. 2 in **II**). In fact, green LAI and related parameter *f*APAR may not react too quickly to small changes in en-

vironmental conditions. The additional component of primary production – LUE, on the other hand, is highly variable and has been shown to respond strongly to different environmental changes (Gamon et al. 1997, Guo and Trotter 2004, Ahl et al. 2004, Hilker et al. 2012).

3.3. LUE of shrubland communities, estimated by PRI

Recently Garbulsky et al. (2011) reviewed studies where the relationship between remotely sensed vegetation index PRI and vegetation LUE was tested, and found that, in general, PRI can be used at different spatial scales, from leaves to ecosystems, for estimating changes in vegetation light use efficiency. They showed that PRI may be useful in remote sensing assessment of LUE, since it accounted for between 42 and 67% of the total variance of LUE at all spatial scales from leaves to ecosystems. However, they found the lowest R^2 values for PRI vs LUE relationships for shrubland ecosystems.

Our measurements, on the other hand, suggested that PRI can be used for detecting changes in leaf-level quantum use (Fig. 5 in **II**, Table 7 in **II**) and changes in green LAI_e (Fig. 4c in **II**) of dense shrublands, however additional data on shrublands with different productivity would be useful for more conclusive results. We also calculated two different estimations of LUE using data from direct measurements on the ground for separate shrublands and we found that PRI correlated with an index of LUE (Fig. 2c), which is a function of canopy nitrogen content and LAI (Green et al. 2003). Nevertheless, a significant relationship was not found between LUE (based on NPP estimations) and PRI, however, we had too few shrublands with medium to large LUE and we did not use site-specific light extinction coefficients. As the assumption of spherical leaf angle distribution is often not valid, (an example of actual leaf angle distributions of forest canopy is given in Fig. 2 in **III**), site specific light extinction estimations should be used when calculating LUE (Stadt and Lieffers 2000, Sarlikioti et al. 2011). Both LUE estimations were significantly correlated with each other (Fig. 2c). The low R^2 of LUE-PRI relationship in shrublands in the paper by Garbulsky et al. (2011) is based on data from southern shrublands, as only a few studies on PRI and LUE relationships in shrublands are available.

Our measurements of southern shrublands also revealed that, in sparse vegetation (Fig. 4c in **II**) and brighter soil (Tables 6 and 7 in **II**), PRI is not sensitive enough for detecting differences in vegetation. Indeed, Filella et al. (2004) showed in the sparse southern shrubland of Spain that NDVI followed the seasonal fluctuations in photosynthesis much better than PRI. PRI has also been reported to be a less reliable estimator of photosynthetic activity in severe stress, including drought conditions (Sims et al. 2006, Ripullone et al. 2011, Porcar-Castell et al. 2012). However, in our experiment with modest drought treatment no significant differences in variances of PRI vs quantum use relations were detected when compared to control plots (Fig. 5 in **II**). Even though we did not

detect any effect of artificial drought on PRI (Fig. 2b in **II**), we found that warming treatment generally increased PRI values in shrublands (Fig. 2a in **II**).




Figure 2. Relationship between two estimations of canopy light-use efficiency and a reflectance index PRI. (a) LUE was estimated using calculations of NPP and LAI (Monteith 1972, Montieih and Moss 1977). (b) Index_{LUE} is a function of nitrogen content of leaves and LAI (Green et al. 2003). p values for significant differences are shown, ns means $p > 0.05$, $N = 6$.

Our study suggests that in modest stress conditions, PRI is more sensitive to changes in the physiological processes of shrubland vegetation than the previously discussed NDVI, since the PRI vs green LAI_e relationship was affected by treatments (Table 4 in **II**), unlike the NDVI vs green LAI_e relationship. More challenging is the interpretation of the exact physiological mechanism, that influences PRI, as we found no significant treatment effect on leaf-level measurements of quantum capture of PSII (Fig. 5 in **II**), even though PRI was significantly correlated to fluorescence-based quantum use efficiency, since the largest part of the variation in PRI among fluorescence parameters was described by changes in non-photochemical quenching (Fig. 5 in **II**, Table 5 in **II**). It is probable that treatments influenced the carotenoid/chorophyll relationship of plants, which has been shown to be another parameter that affects the PRI signal (Filella et al. 2009). However, as the relationships between PRI and plant physiological traits appear to be species-specific, much work remains to be done to understand the

short and long-term factors influencing the values of PRI for different plant communities (Peñuelas et al. 1995, Ahl et al. 2004, Filella et al. 2004, Liu et al. 2013).

3.4. Effect of canopy structure on light use efficiency of canopies and single leaves

Our results in European shrublands showed that the structure of vegetation (approximated as the variation in green LAI_v) clearly explained the largest part of the variation in the reflectance index PRI (Table 5 in **II**), even if PRI was significantly related to the physiological traits of vegetation. In fact, it is shown that not only LAI but also the vertical structure of the canopy and leaf orientation affects values of PRI (Nichol et al. 2000, Barton and North 2001, Damm et al. 2010). Barton and North (2001) demonstrated that angle between leaf normal and the sun determines the intensity of light received by the leaf and hence the LUE and PRI of the leaf, but also it determines the signal strength from that leaf and therefore its relative contribution to canopy PRI. Thus, as real canopies vary in leaf angle distributions, and this changes during plant growth and due to changes in environmental conditions, it is very problematic to draw a single function which transfers PRI or fluorescence yield into LUE of the whole canopy without considering differences in canopy structure (Barton and North 2001, Damm et al. 2010). In the case of forests the effect of structure and resultant leaf-level light-acclimation differences on parameters of remote sensing are even more severe, as the fraction of leaves of the whole canopy that can actually be remotely “sensed“ from a single view angle is even smaller. Fortunately the first steps for incorporating estimations of shadow fraction into remote estimations of LUE have recently been taken (Hall et al. 2011, 2012, Hilker et al. 2012) and an increasing network of carbon dioxide flux measurement towers (Baldocchi, 2008) further helps us to understand the deviations between remotely sensed vegetation indices and carbon fluxes of whole communities.

In **III** we focused on the impact of canopy structure on light capture efficiency of forest ecosystems, more specifically on leaf- and chloroplast-level light acclimation differences between two contrasting tree species. An often used equivalent for leaf-level light use efficiency is quantum capture efficiency of a leaf, estimated by changes in fluorescence emission. Differences in quantum efficiencies of leaves have often been demonstrated along the vertical light gradient of the forest canopy (Fig. 1 d–f in **III**). Less noticed is the within-leaf chloroplast acclimation profile, which also adapts to changes in light environment. In fact, similarly to other optical methods, leaf-level fluorescence measurements also estimate light capture efficiency remotely, and our aim was to evaluate the dependence of fluorescence-based light capture efficiency estimations on canopy structure, focusing on leaf angle distribution (**III**).

Our measurements of chlorophyll fluorescence in trees showed that the ratio of Φ_{II} , qP and NPQ between alternate leaf sides, as an approximation of intra-leaf light acclimation differences, significantly correlated with leaf angles (Fig.

3 in **III**). The ratio between quantum use efficiency of alternate leaf sides was in general larger in more horizontal leaves, as opposed to vertically oriented leaves. However, our results also indicated that the relationship between adaxial/abaxial quantum use efficiency of leaves and leaf angles was species specific (Fig. 3 in **III**), being significant in the case of light-demanding *Populus tremula* and not significant in shade-tolerant *Tilia cordata*. Considering that the chlorophyll fluorescence of only the upper side of the leaf is often used for estimating the light capture efficiency of the whole leaf, our results indicate that, depending on species, the actual light capture efficiency of the whole leaf can be a function of local light conditions that are determined by leaf angle.

Tsuyama et al. (2003) suggested that the measurements of gas exchange and chlorophyll fluorescence detect signals from different populations of chloroplasts in a leaf. If so, the disproportion between fluorescence measurements of leaf abaxial side and gas exchange measurements might also be different for leaves with different orientation and resultant light conditions. This is because variations in the light conditions of different subpopulations of chloroplasts can change the correlation between whole-leaf photosynthetic properties and the photosynthetic properties of individual chloroplasts near the leaf surface, from where the fluorescence signal is mainly derived (Peguero-Pina et al. 2009). At the same time the region from which the fluorescence signal originated also depends on leaf structure, chlorophyll content and excitation wavelength (Cui et al. 1991, Evans 1999, Vogelmann and Han 2000, Buschmann 2007, Peguero-Pina et al. 2009), all of which change depending on leaf orientation and species. Therefore, for parameterization of vegetation structure so that leaf-scale properties may be related to the properties of the whole ecosystem, different adaptation strategies of species must be taken into account. Unfortunately only a few models currently include representations of specific plant responses, probably because available empirical studies are often not at scales relevant for models (Smith and Dukes 2013).

3.5. Chlorophyll distribution within leaves and canopy

Although leaf angles correlated with the gradient of within-leaf physiology, leaf orientation is merely controlling the amount of photochemically active radiation that is received by both leaf sides at different depths in the forest canopy. Thus, to understand the mechanisms behind the above discussed differences in leaf light use efficiency and optical estimations of light use efficiency, we performed additional studies. In **III** and **IV** we investigated the physiological traits that respond to differences in local light environment and thus produce the different fluorescence figures for alternate leaf sides. First we studied leaf reflectance in red and green spectral regions, which, similarly to fluorescence signals are influenced by a combination of leaf anatomy and biochemistry (Govaerts et al. 1996, Peguero-Pina et al. 2009). Leaf reflectance and absorbance measurements from alternate leaf sides indicated that light demanding *P.*

tremula enhanced the photosynthetic potential of leaf biochemistry by decreasing the interception of excess irradiance on the abaxial side of the leaf by increased reflectance (Table 1 in **III**), resulting in smaller differences in fluorescence parameters measured from leaf adaxial and abaxial surfaces (Fig. 3 in **III**). Shade-tolerant *T. cordata* on the other hand increased the light absorptance of leaves (Table 1 in **III**).

Based on earlier studies, which have determined leaf-area-based chlorophyll content (Chl_s) and leaf mass per area (LMA) as being the best proxies for light adaption of the photosynthetic apparatus of leaves (Tsuyama et al. 2003), we investigated Chl_s and LMA as parameters correlating to most of the optical and photochemical differences between alternate leaf sides. Indeed, LMA and chlorophyll content correlated with optical properties for a single species (Fig. 4 in **III**). But when we pooled together data from both species with different strategies (shade-tolerant and light-demanding) the correlative relationship often failed (Fig. 4 in **III**). Furthermore, leaf reflectance and absorptance in the green spectral region were more strongly correlated to variations in leaf chlorophyll content and LMA than reflectance and absorptance measured from the strongly absorbing red spectral region (Table 2 in **III**). Within one species, LMA explained more of the variance in leaf optics than Chl_s (Table 2 in **III**), possibly because the arrangement of chloroplasts can be a more important determinant of light capture than the quantity of chlorophyll *per se* (Souza and Valio 2003).

The importance of leaf structural properties in regulating light use efficiency became especially evident with the finding that *P. tremula* with leaves of significantly more mass per area than *T. cordata* (Fig. 1b in **III**), revealed lower values of absorption (Table 1 in **III**) and smaller differences between light dissipation regulations of alternate leaf sides (NPQ, Fig. 3 in **III**). However, for light-demanding *P. tremula*, leaf angles appeared to be better estimators of differences between quantum use efficiency of alternate leaf sides (Fig. 3 in **III**) than LMA or Chl_s , contrary to *T. cordata*, where the ratio of adaxial/abaxial NPQ was significantly correlated to LMA ($r=0.52$, $p<0.05$). The results given in **IV** confirmed our findings of species specific differences in chlorophyll distribution, as similar shade adjustment of LMA was accompanied with modest changes in chlorophyll concentration, while the densities of different photosynthetic units adjusted to light differences more extensively (Table 1 in **IV**).

In order to investigate more precisely the rearrangement of pigments and to find which parts of the photosynthetic machinery are most responsive to differences in light conditions, we performed an additional investigation (**IV**). Main chloroplast parameters, which have shown to influence the efficiency of light harvesting in photosystem II are the number of LHCII subunits, the arrangement of pigments within them, the interaction between subunits and their distance from the reaction centre complex (Horton 2012). Our results on chlorophyll a/b ratio (Chl a/b) showed significant redistribution of chlorophylls due to acclimation to shade in the canopies of the measured tree species (Fig. 3). The parameter Chl a/b is often used as a measure of the distribution of chlorophyll between antenna and core complexes of the photosynthetic machinery, as most

studies show that Chl b is present mainly in the antenna complex of photosystem II (LHCII) (Evans 1989, Green & Durnford 1996, Kitajima and Hogan 2003). However, recent studies have shown significant migration of LHCII to PSI complexes during state transition (Minagawa 2011) and in mutant plants lacking state transitions, LHCI itself can be enriched in chlorophyll b (Ruban et al. 2003, Ruban and Johnson 2009). Thus it may be difficult to deduce the actual quantity of LHCII of PSII complex from the changes in Chl a/b in natural canopies.




Figure 3. Ratio of leaf chlorophyll a and b for different deciduous forest species, measured in different study years. Different letters at the top of columns indicate significant differences between shade and sun leaves of the same species and same year according to Wilcoxon signed rank test, $p < 0.001$.

Our calculations of realistic shade acclimation in natural forest systems (IV) showed that sun leaves of *B. pendula* on average attached 2.2 trimers of LHCII to PSII, while shade leaves of the same species on average added one more trimer of LHCII to the PSII core, resulting in 3.2 LHCII trimers per PSII on average at shade. The addition of one trimer of LHCII was accompanied by a decrease in Chl a/b ratio of *B. pendula* by 0.4 units in shade, which is consistent with earlier studies (Walters et al. 1999). At the same time the similar decrease of 0.4 units of Chl a/b in the shade-adapted leaves of *T. cordata* did not mean a similar increase in LHCII per PSII. In sun leaves of *T. cordata*, on average 2.6, and in those of *Solidago virgaurea*, 3.9 LHCII trimers were attached to PSII, but the average number of trimers did not change due to shade adjustment in those two species (Table 1 in IV).

According to our data, the main difference between *B. pendula* and *T. cordata* lied in different densities of PSI and PSII and in the size of PSI antenna (Table 1 in IV). The light adjustment of LHCII is widely known, but our results indicate significantly larger light adjustment of PSI, which increased by about

two LHC trimers due to shade adjustment in *Betula pendula* and was larger on average than the PSII antenna of same leaves (Table 1 in IV). A similar doubling of the light harvesting complex of PSI has been observed by Bailey et al. (2001), while Ballottari et al. (2007) recorded no adjustment of numbers of LHCI in *Arabidopsis thaliana* and revealed that the increase in light capture ability of PSI at low light conditions was achieved through migration of LHCII to PSI complex. It is most probable that the increase in PSI LHC due to shade adjustment in IV was also caused by the migration of two trimers of LHCII to PSI. Recently a lot of work has been done in order to understand the environmental triggers for the photosynthetic state transitions and accompanying migration of LHCII to PSI-LHCI systems (Kovács et al. 2006, Ruban and Johnson 2009, Minagawa 2011, Kouřil et al. 2013), but a lot more needs to be done to understand this mechanism at the level needed for ecological applications. Most probably the capacity for state transitions might be one of the parameters determining the light use efficiency of different species.

Our investigations showed that an increase in PSI antenna occurred not due to increasing chlorophyll concentration, but due to decreasing PSI density in all studied species (Table 1 in IV) and therefore an increasing PSII/PSI ratio with decreasing irradiation was found in all studied species. Most probably the decreasing density of PSI with shade can be associated with a higher proportion of far-red irradiation, which has been shown to be characteristic of lower layers of the forest canopy (Lieffers et al. 1999), since previous studies have shown that changes in red/far-red ratio of irradiation may cause changes in photosystem stoichiometry (Chow et al. 1990, Murchie and Horton 1998). A paper by Ruban et al. (2006) showed that the LHCI antenna itself can also be considerably upregulated when state transition is absent, however no data exists on wild-type species with the absence of photosynthetic state transition. Several contradictions in papers on light acclimation of photosystems suggest that the acclimational changes of PSI might be species specific, and thus, studies on single model-species cannot be applied to a larger set of species. Very little is yet known also about the effect of spectral differences at the top and bottom of forest canopies on the differences in the photosystem composition and how this relates to the quantum yield of photosynthesis (Hogewoning et al. 2012). Our findings of different stoichiometrical changes in chlorophyll distribution between different species as a result of shade explain the reports of species specific relationship between PRI and plant physiology (Peñuelas et al. 1995, Ahl et al. 2004, Filella et al. 2004, Liu et al. 2013) as canopy reflectance measurements often reflect more strongly certain aspects of light use efficiency, such as PRI is significantly influenced by the xanthophyll cycle of nonphotochemical quenching, underestimating the possibility that similar light use efficiency may be achieved through alternative mechanisms of light capture and re-funnelling of excess light.

3.6. Nitrogen distribution within leaves and canopy

The high density of chlorophyll in light harvesting complexes is crucial for efficient use of leaf nitrogen as nitrogen availability is often limiting photosynthesis. Remarkably small concentration of nitrogen-containing proteins binds chlorophylls in light harvesting complexes in a way, which enables the efficient capture of light (Beddard and Porter 1976). Despite of nitrogen-efficient molecules of light capture plants appear to optimize nitrogen use at every step of light acclimation. Besides modest changes in chlorophyll concentration, we found concentrations of leaf nitrogen and the nitrogen-rich protein ribulose-bisphosphate carboxylase-oxygenase (Rubisco) per leaf area to be more responsive to changes in light conditions, as nitrogen of leaves decreased by a factor of three and Rubisco decreased by a factor of four as a response to shade if measured on the basis of leaf area (Fig. 7 in IV). Fig. 4 shows that a similar decrease in leaf-area-based nitrogen (N_s) was present both in the vertical light gradient of the forest canopy of III and in the case of decreasing annual irradiance associated with the vertical distance between different shrublands in Europe (I and II). However, the significant shade acclimation for nitrogen and protein concentration came mainly from differences in LMA, since mass-based leaf nitrogen (N_m) was not significantly related to different light regimes.




Figure 4. Leaf-area-based nitrogen content of (a) lower canopy species *T. cordata* and upper canopy species *P. tremula* in a deciduous forest in Järvelja, Estonia and (b) dominant species of six European shrublands. Mean \pm St. Err. are shown, different letters on top of mean values of forest species denote significant differences between sites according to Fisher LSD post-hoc test.

According to our results on forest species with similar nitrogen availability, N_s investment into light-harvesting components of photosynthesis increased as a result of shade acclimation, while N_s invested into Rubisco decreased (Fig. 8 in IV). Our results indicated that the larger investment of nitrogen into light-harvesting proteins in shade might not be enough to keep the excitation rate of a

PSII centre invariant. As PSI is in close contact with the carbon assimilation system, the density of PSI shade-adjusted almost proportionally to changes in Rubisco, while antenna of PSI correspondingly increased (Fig. 8 in **IV**). The general changes in Rubisco and light capture compartments did not differ considerably between the herbaceous species *Solidago virgaurea* and tree species (Fig. 7 in **IV**), in contrast to what was theoretically predicted by a nitrogen allocation model by Xu et al. (2012). The model (Xu et al. 2012) predicted that lower levels of radiation will have a much stronger effect on the allocation of nitrogen to carboxylation for herbaceous plants than for trees, based on the assumption of proportionally smaller investment of nitrogen into the cell walls of herbaceous leaves. However, the non-photosynthetic nitrogen is not just in cell-walls but also in amino acids, nucleic acids, cytosolic proteins, ribosomes and mitochondria (Evans and Seemann 1989, Hirose et al. 1989). Additionally it has been shown that part of the Rubisco can be inactive and is functioning as reserve protein (Eichelmann and Laisk 1999, Warren et al. 2000).

Our study, on the other hand, excluded also the inactive Rubisco from the calculations of non-photosynthetic nitrogen. Nevertheless, we found that leaf acclimation to shade was followed by a considerable decrease in the reserve of non-photosynthetic fraction of nitrogen (Fig. 7 in **IV**) in tree species and also in the herbaceous *Solidago virgaurea*. Recently the effect of variation in the non-photosynthetic component of N on nitrogen allocation to photosynthesis has been further investigated and it was found that nitrogen investment to cell walls does not always affect the content of photosynthetic N (Hikosaka and Shigeno 2009) although N in cell walls is strongly related with LMA as was expected previously (Hikosaka 2004). However, as the non-photosynthetic N fraction does change within a single canopy, extrapolation of nonphotosynthetic N content from the nitrogen-photosynthesis relationship, where photosynthesis has reached zero, might be questionable (Wright et al. 2004).

Thus investigations on a larger variety of species are needed to understand the origin and variability of the non-photosynthetic fraction of nitrogen in different functional groups, since our study indicated the importance of light acclimation of non-photosynthetic N also in short-lived herbaceous species. Similarly the variability in Rubisco has not yet been studied across a range of different species, although there are indications that the relationship between leaf maximum carboxylation capacity and leaf-area-based photosynthetic nitrogen differ between plant functional groups (Maire et al. 2012). Significant differences in the light-acclimation related variability in photosynthetic compartments or non-photosynthetic N may have considerable effect on modelling vegetation production using C/N relations. This might become even more relevant, if canopy nitrogen content is acquired from remotely sensed data that does not consider within-canopy variation in the fraction of non-photosynthetic nitrogen in leaves.

3.7. Nitrogen budget in relation with carbon production of vegetation

The availability of nitrogen has been shown to be one of the main constraints in the carbon cycling of terrestrial ecosystems (LeBauer and Treseder 2008), and models of ecosystem functioning need input info about nitrogen pools and fluxes of different ecosystems. Nevertheless, very fragmented data is available on the nitrogen cycle of shrublands. Thus in **I** we constructed an integrated scheme of aboveground and belowground nitrogen distribution and fluxes at six European shrubland ecosystems (Fig. 7 in **I**). The largest nitrogen resources (468 g N m^{-2}) were found in soil organic matter and roots of the wettest and coldest shrubland in the UK, while lowest values of belowground nitrogen (93 g N m^{-2}) were found on Hungarian dry and sandy soils (Fig. 7 in **I**) however, the ratio of C/N in the top soil was also largest at the UK site and lowest at the Spanish site (Table 2 in **I**). We found that the size of the aboveground nitrogen pool of the studied shrublands was 2–6% of the belowground nitrogen concentration (Fig. 7 in **I**).

On the other hand, the aboveground nitrogen content, measured as average nitrogen content per leaf area generally increased with increasing light availability according to our study (Fig. 4), and nitrogen has been shown to relate significantly also with photosynthesis across various species and vegetation types (Reich et al. 1997, Niinemets et al. 2001, Green 2003, Niinemets 2010). Therefore, relationships between canopy irradiance, nitrogen and photosynthesis are used for modelling global carbon exchange between vegetation and the atmosphere (Friend 2001, Mäkelä et al. 2008), but these known relationships are mainly scaled up from leaf-level measurements. Recently also a strong relationship between remotely sensed nitrogen content of aboveground vegetation and photosynthetic capacity of whole canopy has been reported for forest ecosystems (Ollinger et al. 2008). However, Ollinger et al. (2008) did not include respiration into their calculations and thus they were not able to conclude, whether changes in nitrogen content and photosynthesis will always lead to a variation in net carbon sequestration. Also the prediction of carbon sequestration using data on canopy nitrogen content may be problematic in changing climate conditions, as a review paper by Sardans et al. (2012) showed significant differences in C/N relationships of plants as a response to changed climate and nitrogen deposition.

Nevertheless, our results on six shrublands suggested that canopy nitrogen content of shrublands possibly gives information about the most variable component of primary production, the light use efficiency (LUE), since an index of LUE, which was calculated as a function of leaf nitrogen content (see equation 20. of thesis; Green et al. 2003), was significantly correlated with calculations of LUE and remotely sensed PRI (Fig. 2). For a large variety of forest tree species, maximal light use efficiency has also been found to depend linearly on nitrogen content (Kergoat et al. 2008, Peltoniemi et al. 2012) at all levels of incident PPFD, while Peltoniemi et al. (2012) found that canopy LUE increased at all forest sites by 23% when mean canopy N increased from 1 % to 2%.




Figure 5. Relationship between mean carbon and nitrogen content of six European shrubland ecosystems.

According to six shrublands of present study, the aboveground nitrogen content and aboveground carbon pools appeared to be linearly related (Fig. 5), suggesting that using nitrogen content of vegetation for estimating vegetation productivity could be valid also for European shrublands. However, for more conclusive results additional data on shrublands with medium to high biomass would be needed. Also, further investigations on the same shrublands by Beier et al. (2008) revealed that C and N cycles respond asymmetrically to warming treatment and concluded that global warming effects on shrublands may lead to progressive nitrogen limitation and resultant changes in plant production. As the effect of climate on C/N relationships in plants is shown to be quite universal (Sardans et al. 2012), estimations of carbon sequestration using data on canopy nitrogen content is even more complicated. On a global scale the inevitable factor determining the C/N ratio of the vegetation is the proportion of wood in plants (Wania et al. 2012). We found the average C/N ratio to be 68 in European shrublands (carbon/nitrogen in aboveground plant parts and roots), which is consistent with that predicted for shrubs by a model of Wania et al. (2012), predicting an average C/N for shrublands to be 75 ± 37 . However, at large scale there is a discrepancy between calculations of global C/N ratio, since very different estimations of global nitrogen concentrations (3–18 Pg N) have been reported for the same years (Lin et al. 2000, Yang et al. 2009, Zaehle et al. 2010, Wania et al. 2012), resulting in average global C/N ratio to vary extensively from 30–200 (Wania et al. 2012). Thus the relationships between canopy-scale nitrogen concentration and plant production still need to be tested in different ecosystem types and climatic conditions on a canopy scale, and information on the nonphotosynthetic proportion of canopy and root N is evidently lacking.

4. CONCLUSIONS

1. We conclude that in remote estimations of carbon budget of shrublands, considerable accumulation of carbon into belowground must be taken into account. In case of studied shrublands the average ratio of belowground/litterfall carbon content in general exceeded the respective ratio of carbon reported for forest systems. Nevertheless, the sizes of above- and belowground carbon pools of different shrublands were related to each-other. European shrublands ranged from being carbon sinks to carbon sources and differed in soil nitrogen content and availability.

2. Differences in green leaf area index and light use efficiency of vegetation were efficiently detected by remotely measured canopy reflectance indexes NDVI and PRI above several European shrublands. However, the relationship between NDVI and green leaf area index tended to saturate at dense shrublands, even if the green spectral region was used in NDVI calculations instead of red. Nevertheless, in dense northern shrublands the relationship was found to be stronger if NDVI was calculated using the green spectral region. Reflectance index PRI appeared to be closely related to changes in light use efficiency of plants at dense northern shrublands, but this relationship was absent in sparse southern shrublands. The treatment effects on green leaf area index and light use efficiency, estimated by reflectance indexes, were minimal and differed in directions of change between shrublands.

3. This study revealed strong effect of canopy structure on the canopy- and leaf-level estimations of light use efficiency. The reflectance parameter PRI was significantly influenced by variations in green leaf area index of shrublands. Measurements of leaf fluorescence revealed, that for light-demanding *Populus tremula*, leaf angles appeared to estimate the differences between light use efficiency of alternate leaf sides (suggesting within-leaf differences in chloroplast acclimation to light), compared to *Tilia cordata*, where the differences in chloroplast light acclimation were better correlated to leaf mass per area.

4. Our investigations of within-leaf mechanisms that influence leaf light use efficiency revealed that differences in leaf chlorophyll a/b ratios, which have long been used as a proxy for chlorophyll distribution between light harvesting complex of photosystem II and core complexes of photosystems, reflected different stoichiometrical changes in different species. Unlike many earlier studies, we found that the size of light harvesting complex of photosystem I and the abundance of photosystem I were more responsive to differences in irradiance than the size of light harvesting complex of photosystem II.

5. We demonstrated that the main leaf trait changing due to light adjustment in all studied species appeared to be the non-photosynthetic fraction of leaf nitro-

gen, variation of which may have significant effect on modelling vegetation production using C/N relation and remotely sensed nitrogen estimations.

6. The finding of significant correlation between light use efficiency and nitrogen based index of light use efficiency according to six shrublands suggests that at European shrublands it may be possible to calculate canopy light use efficiency as a function of canopy nitrogen, since the aboveground nitrogen and carbon pools of studied shrublands were also related, however reports of significant changes in carbon/nitrogen ratios of plant tissues as a result of changing climate may complicate the extrapolation over larger areas and over time.

5. REFERENCES

- Ahl DE, Gower ST, Mackay DS, Burrows SN, Norman JM, Diak GR. 2004. Heterogeneity of light use efficiency in a northern Wisconsin forest: implications for modeling net primary production with remote sensing. *Remote Sensing of Environment* **93**: 168–178.
- Aiba SI, Hanya G, Tsujino R, Takyu M, Seino T, Kimura K, Kitayama K. 2007. Comparative study of additive basal area of conifers in forest ecosystems along elevational gradients. *Ecological Research* **22**: 439–450.
- Bailey S, Walters RG, Jansson S, Horton P. 2001. Acclimation of *Arabidopsis thaliana* to the light environment: the existence of separate low light and high light responses. *Planta* **213**: 794–801.
- Baldocchi D. 2008. TURNER REVIEW No. 15. 'Breathing' of the terrestrial biosphere: lessons learned from a global network of carbon dioxide flux measurement systems. *Australian Journal of Botany* **56**: 1–26.
- Baldocchi DD, Wilson K. 2001. Modeling CO₂ and water vapor exchange of a temperate broadleaved forest across hourly to decadal time scales. *Ecological Modelling* **142**: 155–184.
- Ballottari M, Dall'Osto L, Morosinotto T, Bassi R. 2007. Contrasting behavior of higher plant photosystem I and II antenna systems during acclimation. *Journal of Biological Chemistry* **282**: 8947–8958.
- Barton CVM, North PRJ. 2001. Remote sensing of canopy light use efficiency using the photochemical reflectance index: Model and sensitivity analysis. *Remote Sensing of Environment* **78**: 264–273.
- Beddard GS, Porter G. 1976. Concentration quenching in chlorophyll. *Nature* **260**: 366–367.
- Beier C, Emmett B, Gundersen P, Tietema A, Peñuelas J, Estiarte M, Gordon C, Gorissen A, Llorens L, Roda F, Williams D. 2004. Novel approaches to study climate change effects on terrestrial ecosystems in the field: drought and passive nighttime warming. *Ecosystems* **7**: 583–597.
- Beier C, Emmett BA, Peñuelas J, Schmidt IK, Tietema A, Estiarte M, Gundersen P, Llorens L, Riis-Nielsen T, Sowerby A, Gorissen A. 2008. Carbon and nitrogen cycles in European ecosystems respond differently to global warming. *Science of the Total Environment* **407**: 692–697.
- Bokhorst S, Huiskes A, Convey P, Sinclair BJ, Lebouvier M, Van de Vijver B, Wall DH. 2011. Microclimate impacts of passive warming methods in Antarctica: implications for climate change studies. *Polar Biology* **34**: 1421–1435.
- Buschmann C. 2007. Variability and application of the chlorophyll fluorescence emission ratio red/far-red of leaves. *Photosynthesis Research* **92**: 261–271.
- Campbell GS, Norman JM. 1998. An introduction to environmental biophysics, Springer-Verlag, New York.
- Caylor KK, Dowty PR, Shugart HH. 2003. Relationship between small-scale structural variability and simulated vegetation productivity across a regional moisture gradient in southern Africa. *Global Change Biology*, **10**: 374–382.
- Cescatti A, Niinemets Ü. 2004. Sunlight capture. Leaf to landscape. In: Smith WK, Vogelmann TC, Chritchley C (eds.), *Photosynthetic adaptation: chloroplast to landscape*. Springer, Berlin, pp 42–85.

- Chen JM, Liu J, Cihlar J, Goulden ML. 1999.** Daily canopy photosynthesis model through temporal and spatial scaling for remote sensing applications. *Ecological Modelling* **124**: 99–119.
- Chow WS, Hope AB, Anderson JM. 1990.** A reassessment of the use of herbicide binding to measure photosystem 2 reaction centres in plant thylakoids. *Photosynthesis Research* **24**: 109–113.
- Cui M, Vogelmann TC, Smith WK. 1991.** Chlorophyll and light gradients in sun and shade leaves of *Spinacia oleracea*. *Plant, Cell and Environment* **14**: 493–500.
- Damm A, Elbers J, Erler A, Gioli B, Hamdi K, Hutjes R, Kosvancova M, Meroni M, Miglietta F, Moersch A, Moreno J, Schickling A, Sonnenschein R, Udelhoven T, Van Der Linden S, Hostert P, Rascher U. 2010.** Remote sensing of sun-induced fluorescence to improve modeling of diurnal courses of gross primary production (GPP). *Global Change Biology* **16**: 171–186.
- Davidson EA, Savage K, Bolstad P, Clark DA, Curtis PS, Ellsworth DS, Hanson PJ, Law BE, Luo Y, Pregitzer KS, Randolph JC, Zak, D. 2002.** Belowground carbon allocation in forests estimated from litterfall and IRGA-based soil respiration measurements. *Agricultural and Forest Meteorology* **113**: 39–51.
- de Pury DGG, Farquhar GD. 1997.** Simple scaling of photosynthesis from leaves to canopies without the errors of big-leaf models. *Plant, Cell & Environment* **20**: 537–557.
- Drolet GG, Middleton EM, Huemmrich KF, Hall FG, Amiro BD, Barr AG, Black TA, McCaughey JH, Margolis HA. 2008.** Regional mapping of gross light-use efficiency using MODIS spectral indices. *Remote Sensing of Environment* **112**: 3064–3078.
- Eichelmann H, Laisk A. 1999.** Ribulose-1,5-bisphosphate carboxylase/oxygenase content, assimilatory charge and mesophyll conductance in leaves. *Plant Physiology* **119**: 179–189.
- Evans JR. 1989.** Photosynthesis and nitrogen relationships in leaves of C3 plants. *Oecologia*, **78**: 9–19.
- Evans JR. 1999.** Leaf anatomy enables more equal access to light and CO₂ between chloroplasts. *New Phytologist* **143**: 93–104.
- Evans JR, Seemann JR. 1989.** The allocation of protein nitrogen in the photosynthetic apparatus: costs, consequences and control. In: Briggs WR (ed.), *Photosynthesis*. Liss, New York. pp 183–205.
- Fensholt R, Sandholt I, Rasmussen MS. 2004.** Evaluation of MODIS LAI, fAPAR and the relation between fAPAR and NDVI in a semi-arid environment using in situ measurements. *Remote Sensing of Environment* **91**: 490–507.
- Field CB, Chapin FS, Matson PA, Mooney HA. 1992.** Responses of terrestrial ecosystems to the changing atmosphere: a resource-based approach. *Annual Review of Ecology and Systematics* **23**: 201–235.
- Filella I, Peñuelas J, Llorens L, Estiarte M. 2004.** Reflectance assessment of seasonal and annual changes in biomass and CO₂ uptake of a Mediterranean shrubland submitted to experimental warming and drought. *Remote Sensing of Environment* **90**: 308–318.
- Filella I, Porcar-Castell A, Munné-Bosch S, Bäck J, Garbulsky MF, Peñuelas J. 2009.** PRI assessment of long-term changes in carotenoids/chlorophyll ratio and short-term changes in de-epoxidation state of the xanthophyll cycle. *International Journal of Remote Sensing* **30**: 4443–4455.

- Friend AD. 2001.** Modelling canopy CO₂ fluxes: are ‘big-leaf’ simplifications justified? *Global Ecology and Biogeography* **10**: 603–619.
- Gamon JA, Peñuelas J, Field CB. 1992.** A narrow-waveband spectral index that tracks diurnal changes in photosynthetic efficiency. *Remote Sensing of Environment* **41**: 35–44.
- Gamon JA, Field CB, Goulden ML, Griffin KL, Hartley AE, Joel G, Peñuelas J, Valentini R. 1995.** Relationship between NDVI, canopy structure and photosynthesis in three Californian vegetation types. *Ecological Applications* **5**: 28–41.
- Gamon JA, Serrano L, Surfus JS. 1997.** The photochemical reflectance index: an optical indicator of photosynthetic radiation use efficiency across species, functional types, and nutrient levels. *Oecologia* **112**: 492–501.
- Garbulsky MF, Peñuelas J, Gamon J, Inoue Y, Filella I. 2011.** The photochemical reflectance index (PRI) and the remote sensing of leaf, canopy and ecosystem radiation use efficiencies: a review and meta-analysis. *Remote Sensing of Environment* **115**: 281–297.
- Givnish TJ. 1988.** Adaptation to sun and shade: a whole-plant perspective. *Functional Plant Biology* **15**: 63–92.
- Govaerts YM, Jacquemoud S, Verstraete MM, Ustin SL. 1996.** Three-dimensional radiation transfer modeling in a dicotyledon leaf. *Applied Optics* **35**: 6585–6598.
- Grace J, Nichol C, Disney M, Lewis P, Quaife T, Bowyer P. 2007.** Can we measure terrestrial photosynthesis from space directly, using spectral reflectance and fluorescence? *Global Change Biology* **13**: 1484–1497.
- Green BR, Durnford DG. 1996.** The chlorophyll-carotenoid proteins of oxygenic photosynthesis. *Annual Review of Plant Biology* **47**: 685–714.
- Green DS, Erickson JE, Kruger EL. 2003.** Foliar morphology and canopy nitrogen as predictors of light-use efficiency in terrestrial vegetation. *Agricultural and Forest Meteorology* **115**: 163–171.
- Guo J, Trotter CM. 2004.** Estimating photosynthetic light-use efficiency using the photochemical reflectance index: variations among species. *Functional Plant Biology* **31**: 255–265.
- Haboudane D, Miller JR, Pattey E, Zarco-Tejada PJ, Strachan IB. 2004.** Hyper-spectral vegetation indices and novel algorithms for predicting green LAI of crop canopies: Modeling and validation in the context of precision agriculture. *Remote Sensing of Environment* **90**: 337–352.
- Hall FG, Hilker T, Coops NC, Lyapustin A, Huemmrich KF, Middleton E, Margolis H, Drolet G, Black TA. 2008.** Multi-angle remote sensing of forest light use efficiency by observing PRI variation with canopy shadow fraction. *Remote Sensing of Environment* **112**: 3201–3211.
- Hall FG, Hilker T, Coops NC. 2011.** PHOTOSYNSAT, photosynthesis from space: Theoretical foundations of a satellite concept and validation from tower and spaceborne data. *Remote Sensing of Environment* **115**: 1918–1925.
- Hall FG, Hilker T, Coops NC. 2012.** Data assimilation of photosynthetic light-use efficiency using multi-angular satellite data: I. Model formulation. *Remote Sensing of Environment* **121**: 301–308.
- Heinsch FA, Reeves M, Bowker CF, Votava P, Kang S, Milesi C, Zhao M, Glassy J, Jolly WM, Loehman R, Bowker CF, Nemani RS, Running SW. 2003.** GPP and NPP (MOD17A2/A3) Products NASA MODIS Land Algorithm. *MOD17 User's Guide*, pp 1–57.

- Hikosaka K. 2004.** Interspecific difference in the photosynthesis-nitrogen relationship: patterns, physiological causes, and ecological importance. *Journal of Plant Research* **117**: 481–494.
- Hikosaka K, Shigeno A. 2009.** The role of Rubisco and cell walls in the interspecific variation in photosynthetic capacity. *Oecologia* **160**: 443–451.
- Hilker T, Coops NC, Wulder MA, Black TA, Guy RD. 2008.** The use of remote sensing in light use efficiency based models of gross primary production: A review of current status and future requirements. *Science of the Total Environment* **404**: 411–423.
- Hilker T, Hall FG, Tucker CJ, Coops NC, Black TA, Nichol CJ, Sellers PJ, Barre A, Hollinger DY, Munger JW. 2012.** Data assimilation of photosynthetic light-use efficiency using multi-angular satellite data: II Model implementation and validation. *Remote Sensing of Environment* **121**: 287–300.
- Hilker T, Hall FG, Coops NC, Lyapustin A, Wang Y, Nesic Z, Grant N, Black TA, Wulder MA, Kljun N, Hopkinson C, Chasmer L. 2010.** Remote sensing of photosynthetic light-use efficiency across two forested biomes: spatial scaling. *Remote Sensing of Environment* **114**: 2863–2874.
- Hikosaka K. 2004.** Interspecific difference in the photosynthesis-nitrogen relationship: patterns, physiological causes, and ecological importance. *Journal of Plant Research* **17**: 481–494.
- Hirose T. 2005.** Development of the Monsi–Saeki theory on canopy structure and function. *Annals of Botany* **95**: 483–494.
- Hirose T, Werger MJA. 1987.** Maximizing daily canopy photosynthesis with respect to the leaf nitrogen allocation pattern in the canopy. *Oecologia* **72**: 520–526.
- Hirose T, Werger MJA, van Rheenen JWA. 1989.** Canopy development and leaf nitrogen distribution in a stand of *Carex acutiformis*. *Ecology* **70**: 1610–1618.
- Hogewoning SW, Wientjes E, Douwstra P, Trouwborst G, van Ieperen W, Croce R, Harbinson J. 2012.** Photosynthetic quantum yield dynamics: From photosystems to leaves. *The Plant Cell* **24**: 1921–1935.
- Horton P. 2012.** Optimization of light harvesting and photoprotection: molecular mechanisms and physiological consequences. *Philosophical Transactions of the Royal Society B: Biological Sciences* **367**: 3455–3465.
- Houborg R, Anderson MC, Norman JM, Wilson T, Meyers T. 2009.** Intercomparison of a ‘bottom-up’ and ‘top-down’ modeling paradigm for estimating carbon and energy fluxes over a variety of vegetative regimes across the US. *Agricultural and Forest Meteorology* **149**: 1875–1895.
- Huemmerich KF, Gamon JA, Tweedie CE, Oberbauer SF, Kinoshita G, Houston S, Kuchy A, Hollister RD, Kwon H, Mano M, Harazono Y, Webber PJ, Oechel WC. 2010.** Remote sensing of tundra gross ecosystem productivity and light use efficiency under varying temperature and moisture conditions. *Remote Sensing of Environment* **114**: 481–489.
- IPCC. 2007.** Climate change 2007: The physical science basis. Contribution of working group I. In: Solomon S, Qin D, Manning M, Chen Z, Marquis M, Averyt KB, Tignor M, Miller HL (eds.), *Fourth Assessment Report of the Intergovernmental Panel on Climate Change*. Cambridge, UK, Cambridge University Press.
- Ishii H, Asano S. 2010.** The role of crown architecture, leaf phenology and photosynthetic activity in promoting complementary use of light among coexisting species in temperate forests. *Ecological Research* **25**: 715–722.

- Jonasson S. 1988.** Evaluation of the point intercept method for the estimation of plant biomass. *Oikos* **52**: 101–106.
- Kergoat L, Lafont S, Arneth A, Le Dantec V, Saugier B. 2008.** Nitrogen controls plant canopy light-use efficiency in temperate and boreal ecosystems. *Journal of Geophysical Research* **113**, G04017.
- Kitajima K, Hogan KP. 2003.** Increases of chlorophyll a/b ratios during acclimation of tropical woody seedlings to nitrogen limitation and high light. *Plant, Cell and Environment* **26**: 857–865.
- Kouřil R, Wientjes E, Bultema JB, Croce R, Boekema EJ. 2013.** High-light vs. low-light: Effect of light acclimation on photosystem II composition and organization in *Arabidopsis thaliana*. *Biochimica et Biophysica Acta (BBA)-Bioenergetics* **1827**: 411–419.
- Kovács L, Damkjær J, Kerešič S, Iliaia C, Ruban AV, Boekema EJ, Jansson S, Horton P. 2006.** Lack of the light-harvesting complex CP24 affects the structure and function of the grana membranes of higher plant chloroplasts. *The Plant Cell Online* **18**: 3106–3120.
- Kull O. 2002.** Acclimation of photosynthesis in canopies: models and limitations. *Oecologia* **133**: 267–279.
- Kull O, Jarvis PG. 1995.** The role of nitrogen in a simple scheme to scale up photosynthesis from leaf to canopy. *Plant, Cell and Environment* **18**: 1174–1182.
- Lambers H, Raven JA, Shaver GR, Smith SE. 2008.** Plant nutrient-acquisition strategies change with soil age. *Trends in Ecology & Evolution* **23**: 95–103.
- Lausch A, Pause M, Merbach I, Zacharias S, Doktor D, Volk M, Seppelt R. 2013.** A new multiscale approach for monitoring vegetation using remote sensing-based indicators in laboratory, field, and landscape. *Environmental Monitoring and Assessment*, **185**: 1215–1235.
- Lavorel S, Grigulis K, McIntyre S, Williams NS, Garden D, Dorrough J, Berman S, Quéfier F, Thébault A, Bonis A. 2008.** Assessing functional diversity in the field –methodology matters! *Functional Ecology* **22**: 134–147.
- LeBauer DS, Treseder KK. 2008.** Nitrogen limitation of net primary productivity in terrestrial ecosystems is globally distributed. *Ecology* **89**: 371–379.
- Lieffers VJ, Messier C, Stadt KJ, Gendron F, Comeau PG. 1999.** Predicting and managing light in the understory of boreal forests. *Canadian Journal of Forest Research* **29**: 796–811.
- Lin, BL, Sakoda A, Shibasaki R, Goto N, Suzuki M. 2000.** Modelling a global biogeochemical nitrogen cycle in terrestrial ecosystems, *Ecological Modelling* **135**: 89–110.
- Liu L, Zhang Y, Jiao Q, Peng D. 2013.** Assessing photosynthetic light-use efficiency using a solar-induced chlorophyll fluorescence and photochemical reflectance index. *International Journal of Remote Sensing* **34**: 4264–4280.
- Liu X, Smith WL, Zhou DK, Larar A. 2006.** Principal component-based radiative transfer model for hyperspectral sensors: theoretical concept. *Applied Optics* **45**: 201–209.
- Maire V, Martre P, Kattge J, Gastal F, Esser G, Fontaine S, Soussana JF. 2012.** The Coordination of Leaf Photosynthesis Links C and N Fluxes in C3 Plant Species. *PLoS one* **7**: e38345.
- Martin ME, Plourde LC, Ollinger SV, Smith ML, McNeil BE. 2008.** A generalizable method for remote sensing of canopy nitrogen across a wide range of forest ecosystems. *Remote Sensing of Environment* **112**: 3511–3519.

- Mäkelä A, Pulkkinen M, Kolari P, Lagergren F, Berbigier P, Lindroth A, Loustau D, Nikinmaa E, Vesala T, Hari P. 2008.** Developing an empirical model of stand GPP with the LUE approach: analysis of eddy covariance data at five contrasting conifer sites in Europe. *Global Change Biology* **14**: 92–108.
- Meir P, Kruijt B, Broadmeadow M, Barbosa E, Kull O, Carswell F, Nobre A, Jarvis PG. 2002.** Acclimation of photosynthetic capacity to irradiance in tree canopies in relation to leaf nitrogen concentration and leaf mass per unit area. *Plant, Cell and Environment* **25**: 343–357.
- Meroni M, Colombo R. 2006.** Leaf level detection of solar induced chlorophyll fluorescence by means of a subnanometer resolution spectroradiometer. *Remote Sensing of Environment* **103**: 438–448.
- Minagawa J. 2011.** State transitions—The molecular remodeling of photosynthetic supercomplexes that controls energy flow in the chloroplast. *Biochimica et Biophysica Acta (BBA)-Bioenergetics* **1807**: 897–905.
- Monteith JL. 1972.** Solar radiation and productivity in tropical ecosystems. *Journal of Applied Ecology* **9**: 747–766.
- Monteith JL, Moss CJ. 1977.** Climate and the efficiency of crop production in Britain [and discussion]. *Philosophical Transactions of the Royal Society of London. B, Biological Sciences* **281**: 277–294.
- Moreau D, Allard V, Gaju O, Le Gouis J, Foulkes MJ, Martre P. 2012.** Acclimation of leaf nitrogen to vertical light gradient at anthesis in wheat is a whole-plant process that scales with the size of the canopy. *Plant Physiology* **160**: 1479–1490.
- Mott KA, Woodrow IE. 2000.** Modelling the role of Rubisco activase in limiting non-steady-state photosynthesis. *Journal of Experimental Botany* **51**: 399–406.
- Moya I, Camenen L, Evain S, Goulas Y, Cerovic ZG, Latouche G, Flexas J, Ounis A. 2004.** A new instrument for passive remote sensing: 1. Measurements of sunlight-induced chlorophyll fluorescence. *Remote Sensing of Environment* **91**: 186–197.
- Murchie EH, Horton P. 1998.** Contrasting patterns of photosynthetic acclimation to the light environment are dependent on the differential expression of the responses to altered irradiance and spectral quality. *Plant, Cell and Environment* **21**: 139–148.
- Myers DA, Jordan DN, Vogelmann TC. 1997.** Inclination of sun and shade leaves influences chloroplast light harvesting and utilization. *Physiologia Plantarum* **99**: 395–404.
- Myneni RB, Hoffman S, Knyazikhin Y, Privette JL, Glassy J, Tian Y, Wang Y, Song X, Zhang Y, Smith GR, Lotsch A, Friedl M, Morisette JT, Votava P, Nemani RR, Running SW. 2002.** Global products of vegetation leaf area and fraction absorbed PAR from year one of MODIS data. *Remote Sensing of Environment* **83**: 214–231.
- Nichol CJ, Huemmrich KF, Black TA, Jarvis PG, Walthall CL, Grace J, Hall FG. 2000.** Remote sensing of photosynthetic-light-use efficiency of boreal forest. *Agricultural and Forest Meteorology* **101**: 131–142.
- Niinemets Ü. 2010.** A review of light interception in plant stands from leaf to canopy in different plant functional types and in species with varying shade tolerance. *Ecological Research* **25**: 693–714.
- Niinemets Ü, Ellsworth DS, Lukjanova A, Tobias M. 2001.** Site fertility and the morphological and photosynthetic acclimation of *Pinus sylvestris* needles to light. *Tree Physiology* **21**: 1231–1244.

- Niinemets Ü, Tenhunen JD. 1997.** A model separating leaf structural and physiological effects on carbon gain along light gradients for the shade-tolerant species *Acer saccharum*. *Plant, Cell and Environment* **20**: 845–866.
- Ollinger SV, Richardson AD, Martin ME, Hollinger DY, Frolking SE, Reich PB, Plourde LC, Katul GG, Munger JW, Oren R, Smith M-L, Paw U KT, Bolstad PV, Cook BD, Daya MC, Martin TA, Monson RK, Schmid HP. 2008.** Canopy nitrogen, carbon assimilation, and albedo in temperate and boreal forests: functional relations and potential climate feedbacks. *Proceedings of the National Academy of Sciences* **105**: 19336–19341.
- Parmesan C, Yohe G. 2003.** A globally coherent fingerprint of climate change impacts across natural systems. *Nature* **421**: 37–42.
- Peguero-Pina JJ, Gil-Pelegrín E, Morales F. 2009.** Photosystem II efficiency of the palisade and spongy mesophyll in *Quercus coccifera* using adaxial/abaxial illumination and excitation light sources with wavelengths varying in penetration into the leaf tissue. *Photosynthesis Research* **99**: 49–61.
- Peltoniemi M, Pulkkinen M, Kolari P, Duursma RA, Montagnani L, Wharton S, Lagergren F, Takagi K, Verbeek H, Christensen T, Vesala T, Falk M, Loustau D, Mäkelä A. 2012.** Does canopy mean nitrogen concentration explain variation in canopy light use efficiency across 14 contrasting forest sites? *Tree physiology* **32**: 200–218.
- Peng D, Zhang B, Liu L, Fang H, Chen D, Hu Y, Liu L. 2012.** Characteristics and drivers of global NDVI-based FPAR from 1982 to 2006. *Global Biogeochemical Cycles* **26**.
- Peñuelas J, Filella I, Gamon JA. 1995.** Assessment of photosynthetic radiation-use efficiency with spectral reflectance. *New Phytologist* **131**: 291–296.
- Peñuelas J, Prieto P, Beier C, Cesaraccio C, De Angelis P, de Dato G, Emmett BA, Estiarte M, Garadnai J, Gorissen A, Kovács Láng E, Kröel-Dulay G, Llorens L, Pellizzaro G, Riis-Nielsen T, Schmidt IK, Sirca C, Sowerby A, Spano D, Tietema, A. 2007.** Response of plant species richness and primary productivity in shrublands along a north–south gradient in Europe to seven years of experimental warming and drought: reductions in primary productivity in the heat and drought year of 2003. *Global Change Biology* **13**: 2563–2581.
- Porcar-Castell A, Garcia-Plazaola JI, Nichol CJ, Kolari P, Olascoaga B, Kuusinen N, Fernández-Marín B, Pulkkinen M, Juurola E, Nikinmaa E. 2012.** Physiology of the seasonal relationship between the photochemical reflectance index and photosynthetic light use efficiency. *Oecologia* **170**: 313–323.
- Porcar-Castell A, Palmroth S. 2012.** Modelling photosynthesis in highly dynamic environments: the case of sunflecks. *Tree Physiology* **32**: 1062–1065.
- Portsmouth A, Niinemets Ü. 2007.** Structural and physiological plasticity in response to light and nutrients in five temperate deciduous woody species of contrasting shade tolerance. *Functional Ecology* **21**: 61–77.
- Potter C, Klooster S, Genovese V. 2012.** Net primary production of terrestrial ecosystems from 2000 to 2009. *Climatic Change* **115**: 365–378.
- Potter CS, Randerson JT, Field CB, Matson PA, Vitousek PM, Mooney HA, Klooster SA. (1993).** Terrestrial ecosystem production: a process model based on global satellite and surface data. *Global Biogeochemical Cycles* **7**: 811–841.
- Reich PB, Grigal DF, Aber JD, Gower ST. 1997.** Nitrogen mineralization and productivity in 50 hardwood and conifer stands on diverse soils. *Ecology* **78**: 335–347.

- Ripullone F, Rivelli AR, Baraldi R, Guarini R, Guerrieri R, Magnani F, Peñuelas J, Raddi DS, Borghetti M. 2011.** Effectiveness of the photochemical reflectance index to track photosynthetic activity over a range of forest tree species and plant water statuses. *Functional Plant Biology* **38**: 177–186.
- Rouse Jr JW, Haas RH, Deering DW, Schell JA, Harlan JC. 1974.** Monitoring the vernal advancement and retrogradation (green wave effect) of natural vegetation. *NASA/GSFC, Type III, Final Report*. Greenbelt, MD, USA, pp 1–371.
- Ruban AV, Johnson MP. 2009.** Dynamics of higher plant photosystem cross-section associated with state transitions. *Photosynthesis Research* **99**: 173–183.
- Ruban AV, Solovieva S, Lee PJ, Iliaia C, Wentworth M, Ganeteg U, Klimmek F, Chow WC, Anderson JM, Jansson S, Horton P. 2006.** Plasticity in the composition of the light harvesting antenna of higher plants preserves structural integrity and biological function. *Journal of Biological Chemistry* **281**: 14981–14990.
- Ruban AV, Wentworth M, Yakushevska AE, Keegstra W, Lee PJ, Dekker JP, Jansson S, Boekema E, Horton P. 2003.** Plants lacking the main light harvesting complex retain photosystem II macro-organisation. *Nature* **421**: 648–652.
- Ruimy A, Kergoat L, Bondeau A. 1999.** Comparing global models of terrestrial net primary productivity (NPP): analysis of differences in light absorption and light-use efficiency. *Global Change Biology* **5**: 56–64.
- Sardans J, Rivas-Ubach A, Peñuelas J. 2012.** The C: N: P stoichiometry of organisms and ecosystems in a changing world: A review and perspectives. *Perspectives in Plant Ecology, Evolution and Systematics* **14**: 33–47.
- Sarlikioti V, De Visser PHB, Marcelis LFM. 2011.** Exploring the spatial distribution of light interception and photosynthesis of canopies by means of a functional–structural plant model. *Annals of Botany* **107**: 875–883.
- Schimel JP, Bennett J. 2004.** Nitrogen mineralization: challenges of a changing paradigm. *Ecology* **85**: 591–602.
- Sims DA, Luo H, Hastings S, Oechel WC, Rahman AF, Gamon JA. 2006.** Parallel adjustments in vegetation greenness and ecosystem CO₂ exchange in response to drought in a Southern California chaparral ecosystem. *Remote Sensing of Environment* **103**: 289–303.
- Smith NG, Dukes JS. 2013.** Plant respiration and photosynthesis in global-scale models: incorporating acclimation to temperature and CO₂. *Global Change Biology* **19**: 45–63.
- Souza RP, Válio IF. 2003.** Leaf optical properties as affected by shade in saplings of six tropical tree species differing in successional status. *Brazilian Journal of Plant Physiology* **15**: 49–54.
- Stadt KJ, Lieffers VJ. 2000.** MIXLIGHT: a flexible light transmission model for mixed-species forest stands. *Agricultural and forest meteorology* **102**: 235–252.
- Street LE, Shaver GR, Rastetter EB, van Wijk MT, Kaye BA, Williams M. 2012.** Incident radiation and the allocation of nitrogen within Arctic plant canopies: implications for predicting gross primary productivity. *Global Change Biology* **18**: 2838–2852.
- Terashima I, Hanba YT, Tazoe Y, Vyas P, Yano S. 2006.** Irradiance and phenotype: comparative eco-development of sun and shade leaves in relation to photosynthetic CO₂ diffusion. *Journal of Experimental Botany* **57**: 343–354.
- Tsuyama M, Shibata M, Kobayashi Y. 2003.** Leaf factors affecting the relationship between chlorophyll fluorescence and the rate of photosynthetic electron transport as determined from CO₂ uptake. *Journal of Plant Physiology* **160**: 1131–1139.

- Valladares F, Niinemets Ü. 2008.** Shade tolerance, a key plant feature of complex nature and consequences. *Annual Review of Ecology, Evolution, and Systematics* **39**: 237–257.
- Vogelmann TC, Han T. 2000.** Measurement of gradients of absorbed light in spinach leaves from chlorophyll fluorescence profiles. *Plant, Cell and Environment* **23**: 1303–1311.
- Walters RG, Rogers JJ, Shephard F, Horton P. 1999.** Acclimation of *Arabidopsis thaliana* to the light environment: the role of photoreceptors. *Planta* **209**: 517–527.
- Wania R, Meissner KJ, Eby M, Arora V, Ross I, Weaver AJ. 2012.** Carbon-nitrogen feedbacks in the UVic ESCM. *Geoscientific Model Development Discussions* **5**: 67–118.
- Warren CR, Adams MA, Chen Z. 2000.** Is photosynthesis related to concentrations of nitrogen and Rubisco in leaves of Australian native plants? *Australian Journal of Plant Physiology* **27**: 407–416.
- Way DA, Pearcy RW. 2012.** Sunflecks in trees and forests: from photosynthetic physiology to global change biology. *Tree Physiology* **32**: 1066–1081.
- White MA, Thornton PE, Running SW, Nemani RR. 2000.** Parameterization and sensitivity analysis of the BIOME-BGC terrestrial ecosystem model: net primary production controls. *Earth Interactions* **4**: 1–85.
- Wright IJ, Reich PB, Westoby M, Ackerly DD, Baruch Z, Bongers F, Cavender-Bares J, Chapin T, Cornelissen JHC, Diemer M, Flexas J, Garnier E, Groom PK, Gulias J, Hikosaka K, Lamont BB, Lee T, Lee W, Lusk C, Midgley JJ, Navas M-L, Niinemets Ü, Oleksyn J, Osada N, Poorter H, Poot P, Prior L, Pyankov VI, Roumet C, Thomas SC, Tjoelker MG, Veneklaas EJ, Villar R. 2004.** The worldwide leaf economics spectrum. *Nature* **428**: 821–827.
- Xu C, Fisher R, Wullschleger SD, Wilson CJ, Cai M, McDowell NG. 2012.** Toward a mechanistic modeling of nitrogen limitation on vegetation dynamics. *PloS one*, **7**: e37914.
- Yang X, Wittig V, Jain AK, Post W. 2009.** Integration of nitrogen cycle dynamics into the Integrated Science Assessment Model for the study of terrestrial ecosystem responses to global change. *Global Biogeochemical Cycles* **23** GB4029.
- Yuan W, Liu S, Zhou G, Zhou G, Tieszen LL, Baldocchi D, Bernhofer C, Gholz H, Goldstein AH, Goulden ML, Hollinger DY, Hu Y, Law BE, Stoy PC, Vesala T, Wofsy SC. 2007.** Deriving a light use efficiency model from eddy covariance flux data for predicting daily gross primary production across biomes. *Agricultural and Forest Meteorology* **143**: 189–207.
- Zaehle S, Friend AD, Friedlingstein P, Dentener F, Peylin P, Schulz M. 2010.** Carbon and nitrogen cycle dynamics in the O-CN land surface model: 2. Role of the nitrogen cycle in the historical terrestrial carbon balance. *Global Biogeochemical Cycles* **24** GB1006.
- Zaehle S, Dalmonech D. 2011.** Carbon–nitrogen interactions on land at global scales: current understanding in modelling climate biosphere feedbacks. *Current Opinion in Environmental Sustainability* **3**: 311–320.
- Zarco-Tejada PJ, Miller JR, Mohammed GH, Noland TL, Sampson PH. 2000.** Chlorophyll fluorescence effects on vegetation apparent reflectance: II. Laboratory and airborne canopy-level measurements with hyperspectral data. *Remote Sensing of Environment*, **74**: 596–608.

SUMMARY IN ESTONIAN

Taimede valguskasutuse efektiivsus ning süsiniku ja lämmastiku bilanss: kaugeire kasutamine ja füsioloogilised determinandid

Globaalse kliima pidev muutumine põhjustab muutusi maailma erinevate taimekoosluste primaarproduksioonis, mistõttu uuritakse järjest enam taimkatte süsinikuringet mõjutavaid tegureid. Selleks, et me võiksime piisava täpsusega ennustada muutusi süsiniku sidumises ja primaarproduksioonis, on vaja teada, mil moel keskkonna erinevad omadused primaarproduksiooni mõjutavad.

Laias laastus määrab taimkatte produktsiooni taimede eri osadeni jõudev fotosünteesiliselt aktiivse kiirguse hulk ning valguse kasutamise efektiivsus taimede fotosünteesisüsteemi poolt. Neid kahte primaarproduksiooni osa mõjutavad omakorda erinevad tegurid, teiste hulgas võra ja lehe liigi- ja kooslusespetsiifiline ehitus, fotosünteesiapaaraadi võime kohaneda valgustingimuste erinevustega, mullalämmastiku kättesaadavus ning lämmastiku kasutamise tõhusus, vee kättesaadavus ja temperatuur. Arvestades ülemaailmsete uuringute ulatust, ei ole võimalik muutusi taimede süsinikutasakaalus jälgida vaid paikkondlikult - väga oluline osa taimede ökofüsioloogilistest uuringutest baseerub kaugeire-infol. Seetõttu on kaugeire abil saadud taimkatte peegeldumisindeksite täiustamine ja kasutatavuse hindamine erinevates taimekooslustes üks olulisemaid taimede ökofüsioloogia uurimissuundi tänasel päeval. Et aga kindlaks teha taimkatte peegeldumisindekseid mõjutavad tegurid, on jätkuvalt oluline mõista, mil viisil toimub erinevatele taimeliikidele ja funktsionaalsetele gruppidele omane valguskohanemine nii võra, lehe kui ka kloroplasti tasandil. Seejuures on kloroplasti tasandil toimuva valguskohanemise mehhanismide varieeruvuse kohta reaalses taimekooslustes vähe infot. Sama oluline on uurida lehe valguskasutuse tõhusust mõõtvate tulemuste laiendatavust kogu võrale ja kooslusele, kuna taimevõra valguskasutuse efektiivsuse n-ö alt üles modelleerimisel võib võra struktuurierinevuste või kloroplasti tasandil toimuvate erinevate valguskohanemise strateegiatega arvestamine oluliselt parandada mudeli vastavust reaalsusega.

Käesoleva doktoritöö laiemale eesmärgiks oli uurida, millised faktorid mõjutavad taimkatte produktsiooni ja selle hindamist kaugeire meetodeil, keskendudes valguskasutuse efektiivsusele, mis on primaarproduksiooni üks varieeruvamaid ja raskemini ennustatavaid komponente. Uuriti, millised taimede parameetrid reageerivad kõige ulatuslikumalt muutustele valguskeskkonnas ning kuidas nende parameetrite erinevad väärtused mõjutavad taimkatte valguskasutuse mõõtmisi nii lehe kui võra tasandil.

Töö kitsamad eesmärgid olid järgmised.

1. Kaardistada taimede süsiniku- ja lämmastikuvarusid erinevates Euroopa puhmastikukooslustes.

2. Analüüsida kaugseire meetodil mõõdetud peegeldumisindeksite NDVI ja PRI kasutatavust Euroopa puhmastike primaarproduktiooni komponentide - efektiivse lehepinnaindeksi (green LAI_e) ja valguskasutuse efektiivsuse (LUE) - hindamisel.
3. Uurida võra ja lehestiku struktuuri mõju valguse kasutamise efektiivsuse mõõtmistele.
4. Uurida, kuivõrd varieerub liigiti ja võra erinevates piirkondades klorofüllü jaotus lehes, mis omakorda mõjutab taimede valguse kasutamise efektiivsust.
5. Uurida looduslikku varieeruvust fotosünteesilise ja mittefotosünteesilise lämmastiku hulgas ning jaotuses erinevatel liikidel ja erinevates valgustingimustes.
6. Uurida Euroopa puhmastike näitel, kas taimede lämmastikuisalduse alusel on võimalik kaudselt hinnata süsinikuproduktiooni.

Loetletud eesmärkide saavutamiseks viidi läbi kaks erinevat eksperimenti. Uuriti koosluse süsiniku ja lämmastiku jaotust ning hinnati kaugseire meetodeid kuues erineva valgus-, niiskus- ja temperatuurirežiimiga puhmastikus: Suurbritannias Walesis, Taanis, Hollandis, Ungaris, Hispaanias Kataloonias ja Itaalias Sardiinias. Lisaks pidevale keskkonnaparameetrite salvestamisele mõõdeti ka taimede ja varise süsiniku ning lämmastiku kontsentratsiooni, mulla lämmastikuühendite ja lahustunud orgaaniliste ühendite sisaldust, taimkatte struktuuri parameetreid, mulla hingamist ja dominantsete liikide ülemistel lehtedel klorofüllü fluorestsentsi. Lisaks mõõdeti 1 m kõrguselt taimkatte kohalt peegeldumisindeksid NDVI (spektri punasest piirkonnast NDVI₆₈₀ ja rohelisest piirkonnast NDVI₅₇₀) ja PRI.

Valguskasutuse efektiivsuse ja seda mõjutavate faktorite, klorofüllü ja lämmastiku jaotumise varieeruvust lehes, võras ja liigiti uuriti Eestis Järvselja heitlehises segametsas, kus võrastiku tornid lubasid ligipääsu erinevate valgustingimustega kohanenud lehtedele. Lehe eri külgedelt mõõdeti erinevates valgustingimustes kasvanud lehtede klorofüllü fluorestsentsi ja peegeldumist spektri nähtavas osas ning hinnati lehe ja erinevate fotosünteesisüsteemi komponentide klorofüllü- ja lämmastikuisaldust.

Selleks, et kaugseire meetodeil hinnata muutusi puhmastike primaarproduktioonis, on vaja infot süsiniku jaotumise ja biokeemiliste tsüklite kohta puhmastikukooslustes, kuna erinevalt metsa- ja niidukooslustest on süsiniku ja lämmastiku tasakaalu puhmastikes üsna vähe uuritud. Käesoleva uurimuse tulemusel selgus, et Euroopa puhmastikud varieeruvad oluliselt süsiniku omastamise poolest. Uuritud puhmastike hulgas leidis nii süsinikku neelavaid kooslusi (126 g C m⁻² a⁻¹ niiskes Walesi kanarbiku-mustikapuhmastikus) kui ka süsiniku emiteerijaid (-536 g C m⁻² a⁻¹ mittestabiilses kuivas Itaalia puhmastikus). Niiskes Walesi puhmastikus oli samuti suurim mulla lämmastikuisaldus (468 g N m⁻²), madalaim oli see aga Ungaris kuival liivmullal (93 g N m⁻²). Vaatamata kõrgele süsiniku sidumisele ja mulla lämmastikuisaldusele, oli lämmastiku kättesaadavus üheks taimede kasvu pidurdavaks teguriks Walesi puhmastikus: mulla C/N suhe

Walesi puhmastikus oli 37,4, samas kui vastav suhe kuivas Hispaania puhmastiku mullas oli 12,8. Maapealsete taimeosade süsinikusaldust puhmastikes mõjutas tugevalt keskkonna niiskusrežiim. Kuigi niiskusrežiimi mõju maa-alusele süsinikusaldusele ei väljendunud nii selgelt, olid maapealne ja maa-alune süsinikusaldus omavahel tugevas mittelineaarses seoses. Tänu maapealse ja maa-aluse süsiniku kontsentratsioonide tugevale seotusele on põhjust otsida võimalusi ka kogu koosluse primaarproduktiooni hindamiseks puhmastikukoosluste kohalt kaugseire meetodite abil.

Varasemalt on näidatud, et kaugseire meetodeil saadud peegeldumisindeks NDVI väärtused sõltuvad taimkatte efektiivse lehepinnaindeksi (green LAI_e) väärtustest ja taimede poolt neelatud fotosünteesiliselt aktiivse valguse hulgast. Käesolev uurimus näitas, et NDVI väärtused olid küll oluliselt seotud muutustega puhmastikukoosluste efektiivses lehepinnas, kuid NDVI väärtused küllastusid keskmiste ja suuremate green LAI_e väärtuste juures isegi juhul, kui indeksi arvutamisel kasutati spektri rohelist piirkonda (570 nm) punase (680 nm) asemel. Siiski oli tihedates põhjapoolsetes puhmastikes NDVI ja green LAI_e vaheline seos tugevam juhul, kui NDVI arvutamisel kasutati spektri roheline piirkonda väärtusi. Teise peegeldumisindeksi, PRI väärtusi kasutatakse taimekoosluste valguskasutuse efektiivsuse hindamiseks. Käesoleva uuringu põhjal korreleerusid erinevate puhmastike PRI väärtused taimede lämmastikusalduse baasil arvutatud valguskasutusefektiivsuse indeksiga. Samuti oli PRI tihedates põhjapoolsetes puhmastikes tugevalt seotud taimelehtede kohalt mõõdetud valguskasutuse efektiivsuse väärtustega, kuid lõunapoolsetes puhmastikes taoline seos puudus. Kui uurisime kunstlikult tekitatud põua ja öise temperatuuri tõusu mõju puhmastikele, siis põua tõttu puhmastike NDVI keskmised väärtused enamasti vähenesid ja temperatuuri tõustes PRI väärtused enamasti suurenesid, kuid üksikute puhmastike lõikes oli mõõduka temperatuuritõusu ja põua mõju produktiooni erinevatele komponentidele vähene ja eri maades isesuguse suunaga.

Nii puhmastikukooslustes kui ka Järvelja metsas toimunud uuringud näitasid, et taimede ja koosluse struktuur mõjutab tugevalt nii lehtede (fluorestsentsi mõõtmised) kui koosluse (PRI mõõtmised) valguskasutamise efektiivsuse mõõtmisi. Peegeldumisparameeter PRI näitas valguskasutuse efektiivsust tunduvalt halvemini hõredamates puhmastikes, kus PRI väärtusi mõjutas heleda mulla peegeldumine taimestiku vahelt. Lehe tasandil toimunud mõõtmised näitasid, et ülemise metsarinde liigi, hariliku haava (*Populus tremula*) puhul oli erinevuste järgi lehe nurkades võimalik hinnata lehe üla- ja alaküljelt mõõdetud valguskasutuse efektiivsuste erinevusi (Φ_{II} lehe ülakülg / Φ_{II} lehe alakülg), mis omakorda viitab nähtusele, et erinevused lehe nurkades peegeldavad erinevusi lehtede kloroplastide valguskohanemises. Metsa alumise rinde puuliigi, hariliku pärna (*Tilia cordata*) puhul aga kajastasid lehtede pindtiheduse (LMA) erinevused muutusi lehe eri külgedelt mõõdetud liigse valguse eemale juhtimises (NPQ lehe ülakülg / NPQ lehe alakülg) paremini kui lehe nurgad. Lehe eri külgede teistsugune valguse kasutamine tingib selle, et erinevate nurkade ja pindtihedusega lehtede puhul kirjeldavad fluorestsentsi meetodil lehe ülaküljelt

mõõdetud valguskasutuse efektiivsuse väärtused erinevat fraktsiooni lehe kloroplastidest.

Uurides seejärel põhjalikumalt taimede valguskohanemise mehhanisme, selgus, et kloroplastide valguskohanemise tulemusena muutub lehes klorofüll a ja b jaotus. Metsa ülارينde liigi, arukase (*Betula pendula*) varjulehtedes lisandus fotosüsteem II (PSII) valguspüügi süsteemile (LHC) üks täiendav trimeer ja fotosüsteem I (PSI) valguspüügi aparaadile lisandus varjus kaks trimeeri. Hariliku pärna või hariliku kuldvitsa (*Solidago virgaurea*) puhul sellist selget LHC suurenemist ei ilmnenud. Küll aga vähenes kõigil uuritud liikidel PSI tihedus lehes, mistõttu kõigi liikide varjulehtedes suurenes PSII/PSI suhe. Lehe lämmastikuisaldust mõjutasid veelgi enam erinevused valgustingimustes. Varjulehtedes suurenes sellise lämmastiku hulk, mis oli paigutatud LHC-sse, ning vähem lämmastikku paigutati Rubisco ning PSI tootmisesse. Eriti rõhutas käesolev uuring mittefotosünteesilise lämmastiku osalust taimede valgusega kohanemises, kuna tulemused näitasid, et erinevates valgustingimustes varieerus kõige enam mittefotosünteesilise lämmastiku kogus. Taimede lämmastikuisalduse tugevat seotust valguse kasutamisega näitas ka koosluse valguse kasutuse efektiivsuse (LUE) hea korrelatsioon lehtede lämmastikuisalduse baasil arvatud LUE-indeksiga, kusjuures erinevate Euroopa puhmastike maapealsed lämmastiku- ja süsinikuvarud olid omavahel seotud.

ACKNOWLEDGEMENTS

My greatest thanks go to my supervisors. I am very grateful to Olevi Kull, who managed to bring together my passion for wildlife, heights and far-away places, offering me a research subject that led me to the inner depths of plants, to the very top of trees and to different parts of Europe, and encouraging me during the first painful steps of writing scientific papers. My sincerest gratitude goes to Josep Peñuelas, who without hesitation agreed to take over my supervision after Olevi passed away, helped and encouraged me through the processes of rejections and rewriting and took it very calmly when I told him that the publishing of papers and the writing of the thesis will be delayed because one, another, and another child of mine was about to be born. Many thanks to Anu Sõber, who provided me with facilities and funding, offered me valuable criticism on the thesis and continued to be optimistic and supportive about my doctoral studies.

I am very grateful to all the co-authors of the publications, many thanks for accompanying me into studies of different scales and areas of plant research and providing me with laboratories and valuable datasets. My greatest thanks goes to co-author Lea Hallik, who invested a lot of time and energy helping me at all steps of publishing my first-author papers. Many thanks, to the referees for thorough explanation and criticism during preparation of the publications. I thank Jaan Liira for helping me with statistical analysis and Robert Szava-Kovats and Ilmar Part for reviewing my English. Thank you, Esa Tyystjärvi and Eija Pätsikkä for the opportunity to learn the methodology of extracting chlorophyll-binding proteins from tree leaves at Turku, Finland.

I am really happy to thank my colleagues, it was a great pleasure to work with you indoors and on the study-field, it was a pleasure to discuss whatever questions arose while analysing or writing the papers, it was a pleasure to eat lunch at 20 m height on scaffolding towers in order to avoid mosquitoes, it was nice to pick mushrooms and berries after fieldwork and prepare them together in the evenings, it was a pleasure to sit silently at 4.00 am on a foggy hill in Wales measuring the quantum use efficiency of *Calluna* and watching black grouse playing over our heads, it was nice to crawl with you under the fragrant rosemary bushes of Sardinia in search of tortoises, and finally it was a pleasure to come together for “journal clubs”, “cucumber conferences” and “multi-talented galas”. Thank you, Ingmar Tulva, Lea Hallik, Eve Kaurilind, Helen Karu, Olaf Rääm, Priit Kupper, Mai Kukumägi, Anne Aan, Kersti Püssa, Tsipe Aavik, Maarja Öpik, Aveliina Helm, Virve Sõber, Mari Lepik and Oliver Parrest.

My warmest thanks go to my family for all the love and support and understanding, I could not have finished the thesis without your patience Toomas, Liisa, Mari and Kati. But the most valuable help came from my father and mother – Raivo Mänd and Marika Mänd – they have supported me all the way. My advancements in scientific writing are strongly influenced by my father and my fascination for science comes no doubt from my parents.

The studies have been financially supported by the University of Tartu, Ministry of Education and Research of Estonia, Estonian Science Foundation, Doctoral School of Earth Sciences and Ecology and the European Commission through projects CLIMOOR and VULCAN and through the European Regional Fund (the Centre of Excellence in Environmental Adaptation).

CURRICULUM VITAE

Name: Pille Mänd
Date of birth: 04.02.1978
Citizenship: Estonian
Address: University of Tartu, Institute of Ecology and Earth Sciences,
Department of Botany, Lai 40, 51005 Tartu, Estonia,
E-mail: pille.mand@ut.ee

Education:

1996 Miina Härma Gymnasium
2000 University of Tartu, *BSc* in biology
2002 University of Tartu, *MSc* in plant ecology and ecophysiology

Language skills: Estonian, English, Russian, German

Professional career:

2003–2004 University of Tartu, Institute of Botany and Ecology, technician
2006–2007 University of Tartu, Institute of Botany and Ecology, technician
since 2009 University of Tartu, Institute of Ecology and Earth Sciences,
Department of Botany, researcher

Main fields of research:

Light acclimation of plants, remote sensing of vegetation, responses of plants to changes in climate

List of publications:

- Mänd P**, Hallik L, Peñuelas J, Kull O. 2013. Electron transport efficiency at opposite leaf sides: effect of vertical distribution of leaf angle, structure, chlorophyll content and species in a forest canopy. *Tree Physiology* 33: 202–210
- Mänd P**, Hallik L, Peñuelas J, Nilson T, Duce P, Emmett BA, Beier C, Estiarte M, Garadnai J, Kalapos T, Schmidt IK, Kovács-Láng E, Prieto P, Tietema A, Westerveld JW, Kull O. 2010. Responses of the reflectance indices PRI and NDVI to experimental warming and drought in European shrublands along a north-south climatic gradient. *Remote Sensing of Environment* 114: 626–636
- Beier C, Emmett BA, Tietema A, Schmidt IK, Peñuelas J, Kovács Láng E, Duce P, DeAngelis P, Gorissen A, Estiarte M, de Dato GD, Sowerby A, Kröel-Dulay G, Lellei-Kovács E, Kull O, **Mänd P**, Petersen H, Gjelstrup P, Spano D. 2009. Carbon and nitrogen balances for 6 shrublands across Europe. *Global Biogeochemical Cycles* 23: 1–13
- Riikonen J, Syrjälä L, Tulva I, **Mänd P**, Oksanen E, Poteri M, Vapaavuori E. 2008. Stomatal characteristics and infection biology of *Pyrenopeziza betuli-cola* in *Betula pendula* trees grown under elevated CO₂ and O₃. *Environmental Pollution* 156: 536–543

Eichelmann H, Oja V, Rasulov B, Padu E, Bichele I, Pettai H, **Mänd P**, Kull O, Laisk A. 2005. Adjustment of leaf photosynthesis to shade in a natural canopy: reallocation of nitrogen. *Plant, Cell and Environment* 28: 389–401

Other science publications and editing:

Tulva I, Öpik M, **Mänd P** (eds.). 2006. Theory of Inheritance. *Schola Biotheoretica XXXII*. Estonian Naturalists Society, OÜ Sulemees, Tartu /in Estonian/

Mänd P, Tulva I. 2004. Balancing the global carbon cycle. In: Tulva I, Öpik M, Puura I (eds.), Theory of Equilibrium. *Schola Biotheoretica XXX*. Tartu, OÜ Sulemees, pp. 126–128. /in Estonian/

Kupper P, **Mänd P**, Tulva I. 2003. From dusk to dawn: do plants sleep deeply? In: Tammaru T, Puura I. (eds.), Theory of Rest. *Schola Biotheoretica XXIX*. Tartu, OÜ Sulemees, pp. 21–25. /in Estonian/

Awards and scholarships:

- Scholarships from Doctoral School of Earth Sciences and Ecology for participation in conference at Italia, Rome 2006, in PHYSENSE seminar of Nordic Network at Estonia, Tartu 2007 and in *PhD* courses, Estonia, Häädemeeste 2004 and Järvelja-Vihula, Estonia 2006
- Fellowship for attending a conference at Czech, Olomouc 2006
- Travel scholarships for participation in NordForsk doctoral course at Finland, Joensuu 2004 and Sweden, Umea 2005
- Fellowship for attending the ESP Congress, Norway, Lillehammer 2001.
- Study grant from Rotalia Foundation 1999/2000
- Estonian National Contest for Young Scientists at university level, III Prize, 2000

Other administrative and professional activities:

- Member of organizing team of exposition of fungi at Ulvi, Estonia 2012
- Coordinating the building of websites advertizing the research topics of department of Botany, preparing website of plant ecophysiology and organizing a seminar introducing the research topics of department of Botany for bachelor students, 2007
- Organizing a weekly seminar “Journal club” for graduate-students of biology at 2006/2007
- Member of organizing team of international doctoral school of NordForsk and Doctoral School of Earth Sciences and Ecology, Estonia: Tree canopy – structure and functioning – from below and above. 3–14 September 2006, Järvelja and Vihula, Estonia
- Member of organizing team of spring-schools of “Theoretical Biology” at years 2004–2006
- Member of Estonian Naturalists Society
- Member of Estonian Seminatural Community Conservation Association
- Member of Tartu Students' Nature Protection Circle (1998–1999 in board)

ELULOOKIRJELDUS

Nimi: Pille Mänd
Sünniaeg: 04.02.1978
Kodakondsus: Eesti
Aadress: Tartu Ülikool, Ökoloogia ja Maateaduste Instituut, Botaanika osakond, Lai 40, 51005 Tartu, Eesti,
E-post: pille.mand@ut.ee

Haridus:

Miina Härma Gümnaasium (1996)
Tartu Ülikool, *BSc* bioloogia erialal (2000)
Tartu Ülikool, *MSc* taimeökoloogia ja ökofüsioloogia erialal (2002)

Keelteoskus: eesti, inglise, vene, saksa

Teenistuskäik:

2003–2004 Tartu Ülikool, Botaanika ja Ökoloogia Instituut, tehnik
2006–2007 Tartu Ülikool, Botaanika ja Ökoloogia Instituut, tehnik
alates 2009 Tartu Ülikool, Ökoloogia ja Maateaduste Instituut, Botaanika osakond, teadur

Peamised uurimisvaldkonnad:

Valguskohanemine taimedel, primaarproduktiooni hindamine kaugseire meetodeil, kliimamuutuste mõju taimedele

Publikatsioonide loetelu:

- Mänd P**, Hallik L, Peñuelas J, Kull O. 2013. Electron transport efficiency at opposite leaf sides: effect of vertical distribution of leaf angle, structure, chlorophyll content and species in a forest canopy. *Tree Physiology* 33: 202–210.
- Mänd P**, Hallik L, Peñuelas J, Nilson T, Duce P, Emmett BA, Beier C, Estiarte M, Garadnai J, Kalapos T, Schmidt IK, Kovács-Láng E, Prieto P, Tietema A, Westerveld JW, Kull O. 2010. Responses of the reflectance indices PRI and NDVI to experimental warming and drought in European shrublands along a north–south climatic gradient. *Remote Sensing of Environment* 114: 626–636.
- Beier C, Emmett BA, Tietema A, Schmidt IK, Peñuelas J, Kovács Láng E, Duce P, DeAngelis P, Gorissen A, Estiarte M, de Dato GD, Sowerby A, Kröel-Dulay G, Lellei-Kovács E, Kull O, **Mänd P**, Petersen H, Gjelstrup P, Spano D. 2009. Carbon and nitrogen balances for 6 shrublands across Europe. *Global Biogeochemical Cycles* 23: 1–13.
- Riikonen J, Syrjälä L, Tulva I, **Mänd P**, Oksanen E, Poteri M, Vapaavuori E. 2008. Stomatal characteristics and infection biology of *Pyrenopeziza betulicola* in *Betula pendula* trees grown under elevated CO₂ and O₃. *Environmental Pollution* 156: 536–543.

Eichelmann H, Oja V, Rasulov B, Padu E, Bichele I, Pettai H, **Mänd P**, Kull O, Laisk A. 2005. Adjustment of leaf photosynthesis to shade in a natural canopy: reallocation of nitrogen. *Plant, Cell and Environment* 28: 389–401

Muud teaduslikud publikatsioonid ja osalemine toimetustöös:

Tulva I, Öpik M, **Mänd P** (toim.). 2006. Pärandumise teooria. *Schola Biotheoretica XXXII*. Eesti Loodusuurijate Selts, OÜ Sulemees, Tartu

Mänd P, Tulva I. 2004. Globaalset süsinikuringet rihti ajamas. *Schola Biotheoretica XXX*. Tasakaalu Teooria, toim. Tulva I, Öpik M, Puura I. Tartu, OÜ Sulemees, pp. 126–128

Kupper P, **Mänd P**, Tulva I. 2003. Ehast koiduni: kas taim magab sügavalt? *Schola Biotheoretica XXIX*. Puhkuse teooria, toim. Tammaru T, Puura I. Tartu, OÜ Sulemees, pp. 21–25

Preemiad ja stipendiumid:

- Maateaduste ja ökoloogia doktorikooli stipendiumid osalemaks konverentsil Itaalias, Rooma 2006, PHYSENSE põhjamaade võrgustiku seminaril Eestis, Tartu 2007 ja doktorikoolides Eestis, Häädemeeste 2004 ja Järvselja-Vihula, 2006
- Stipendium osalemaks konverentsil Tšehhis, Olomouc 2006
- Reisistipendiumid osalemaks NordForsk doktorikoolis Soomes, Joensuu 2004 ja Rootsisis, Umea 2005
- Stipendium osalemaks ESP kongressil Norras, Lillehammer 2001
- Õppetipendium Rotalia Fondilt 1999/2000
- Eesti üliõpilaste teadustööde riikliku konkursi III preemia, 2000

Muu teaduslik organisatsiooniline ja erialane tegevus:

- Seenenäituse korraldusmeeskonnas, Ulvi, Rägavere vald, Eesti 2012
- Botaanika osakonna uurimisteesmasid populaarsel viisil tutvustava veebilehe tegemise kordineerimine ja ökofüsioloogia õppetooli uurimissuundi tutvustava veebilehe tegemine ning Botaanika osakonda tutvustava seminari korraldamine bakalaureuse tudengitele, 2007
- Iganädalase bioloogia kraadiõppurite seminari “Journal club” korraldamine 2006/2007
- Põhjamaade võrgustiku Carbon Dynamics in Managed Terrestrial Ecosystems (NordForsk) ja Maateaduste ja ökoloogia doktorikooli PhD kursuse korraldusmeeskonnas. Tree canopy – structure and functioning – from below and above. 3.–14. september 2006, Järvselja ja Vihula, Eesti
- Teoreetilise bioloogia kevadkooli korraldusmeeskonnas aastatel 2004–2006
- Eesti Looduseuurijate Seltsi liige
- Pärandkoosluste Kaitse Ühingu liige
- Tartu Üliõpilaste Looduskaitseringi liige (juhatuses 1998–1999)

DISSERTATIONES BIOLOGICAE UNIVERSITATIS TARTUENSIS

1. **Toivo Maimets**. Studies of human oncoprotein p53. Tartu, 1991, 96 p.
2. **Enn K. Seppet**. Thyroid state control over energy metabolism, ion transport and contractile functions in rat heart. Tartu, 1991, 135 p.
3. **Kristjan Zobel**. Epifüütsete makrosamblike väärtus õhu saastuse indikaatoritena Hamar-Dobani boreaalsetes mägimetsades. Tartu, 1992, 131 lk.
4. **Andres Mäe**. Conjugal mobilization of catabolic plasmids by transposable elements in helper plasmids. Tartu, 1992, 91 p.
5. **Maia Kivisaar**. Studies on phenol degradation genes of *Pseudomonas* sp. strain EST 1001. Tartu, 1992, 61 p.
6. **Allan Nurk**. Nucleotide sequences of phenol degradative genes from *Pseudomonas* sp. strain EST 1001 and their transcriptional activation in *Pseudomonas putida*. Tartu, 1992, 72 p.
7. **Ülo Tamm**. The genus *Populus* L. in Estonia: variation of the species biology and introduction. Tartu, 1993, 91 p.
8. **Jaanus Remme**. Studies on the peptidyltransferase centre of the *E.coli* ribosome. Tartu, 1993, 68 p.
9. **Ülo Langel**. Galanin and galanin antagonists. Tartu, 1993, 97 p.
10. **Arvo Käär**. The development of an automatic online dynamic fluorescence-based pH-dependent fiber optic penicillin flowthrough biosensor for the control of the benzylpenicillin hydrolysis. Tartu, 1993, 117 p.
11. **Lilian Järvekülg**. Antigenic analysis and development of sensitive immunoassay for potato viruses. Tartu, 1993, 147 p.
12. **Jaak Palumets**. Analysis of phytomass partition in Norway spruce. Tartu, 1993, 47 p.
13. **Arne Sellin**. Variation in hydraulic architecture of *Picea abies* (L.) Karst. trees grown under different environmental conditions. Tartu, 1994, 119 p.
13. **Mati Reeben**. Regulation of light neurofilament gene expression. Tartu, 1994, 108 p.
14. **Urmas Tartes**. Respiration rhythms in insects. Tartu, 1995, 109 p.
15. **Ülo Puurand**. The complete nucleotide sequence and infections *in vitro* transcripts from cloned cDNA of a potato A potyvirus. Tartu, 1995, 96 p.
16. **Peeter Hõrak**. Pathways of selection in avian reproduction: a functional framework and its application in the population study of the great tit (*Parus major*). Tartu, 1995, 118 p.
17. **Erkki Truve**. Studies on specific and broad spectrum virus resistance in transgenic plants. Tartu, 1996, 158 p.
18. **Illar Pata**. Cloning and characterization of human and mouse ribosomal protein S6-encoding genes. Tartu, 1996, 60 p.
19. **Ülo Niinemets**. Importance of structural features of leaves and canopy in determining species shade-tolerance in temperature deciduous woody taxa. Tartu, 1996, 150 p.

20. **Ants Kurg.** Bovine leukemia virus: molecular studies on the packaging region and DNA diagnostics in cattle. Tartu, 1996, 104 p.
21. **Ene Ustav.** E2 as the modulator of the BPV1 DNA replication. Tartu, 1996, 100 p.
22. **Aksel Soosaar.** Role of helix-loop-helix and nuclear hormone receptor transcription factors in neurogenesis. Tartu, 1996, 109 p.
23. **Maido Remm.** Human papillomavirus type 18: replication, transformation and gene expression. Tartu, 1997, 117 p.
24. **Tiiu Kull.** Population dynamics in *Cypridium calceolus* L. Tartu, 1997, 124 p.
25. **Kalle Olli.** Evolutionary life-strategies of autotrophic planktonic microorganisms in the Baltic Sea. Tartu, 1997, 180 p.
26. **Meelis Pärtel.** Species diversity and community dynamics in calcareous grassland communities in Western Estonia. Tartu, 1997, 124 p.
27. **Malle Leht.** The Genus *Potentilla* L. in Estonia, Latvia and Lithuania: distribution, morphology and taxonomy. Tartu, 1997, 186 p.
28. **Tanel Tenson.** Ribosomes, peptides and antibiotic resistance. Tartu, 1997, 80 p.
29. **Arvo Tuvikene.** Assessment of inland water pollution using biomarker responses in fish *in vivo* and *in vitro*. Tartu, 1997, 160 p.
30. **Urmas Saarma.** Tuning ribosomal elongation cycle by mutagenesis of 23S rRNA. Tartu, 1997, 134 p.
31. **Henn Ojaveer.** Composition and dynamics of fish stocks in the gulf of Riga ecosystem. Tartu, 1997, 138 p.
32. **Lembi Lõugas.** Post-glacial development of vertebrate fauna in Estonian water bodies. Tartu, 1997, 138 p.
33. **Margus Pooga.** Cell penetrating peptide, transportan, and its predecessors, galanin-based chimeric peptides. Tartu, 1998, 110 p.
34. **Andres Saag.** Evolutionary relationships in some cetrarioid genera (Lichenized Ascomycota). Tartu, 1998, 196 p.
35. **Aivar Liiv.** Ribosomal large subunit assembly *in vivo*. Tartu, 1998, 158 p.
36. **Tatjana Oja.** Isoenzyme diversity and phylogenetic affinities among the eurasian annual bromes (*Bromus* L., Poaceae). Tartu, 1998, 92 p.
37. **Mari Moora.** The influence of arbuscular mycorrhizal (AM) symbiosis on the competition and coexistence of calcareous grassland plant species. Tartu, 1998, 78 p.
38. **Olavi Kurina.** Fungus gnats in Estonia (*Diptera: Bolitophilidae, Keroplattidae, Macroceridae, Ditomyiidae, Diadocidiidae, Mycetophilidae*). Tartu, 1998, 200 p.
39. **Andrus Tasa.** Biological leaching of shales: black shale and oil shale. Tartu, 1998, 98 p.
40. **Arnold Kristjuhan.** Studies on transcriptional activator properties of tumor suppressor protein p53. Tartu, 1998, 86 p.
41. **Sulev Ingerpuu.** Characterization of some human myeloid cell surface and nuclear differentiation antigens. Tartu, 1998, 163 p.

42. **Veljo Kisand.** Responses of planktonic bacteria to the abiotic and biotic factors in the shallow lake Võrtsjärv. Tartu, 1998, 118 p.
43. **Kadri Põldmaa.** Studies in the systematics of hypomyces and allied genera (Hypocreales, Ascomycota). Tartu, 1998, 178 p.
44. **Markus Vetemaa.** Reproduction parameters of fish as indicators in environmental monitoring. Tartu, 1998, 117 p.
45. **Heli Talvik.** Prepatent periods and species composition of different *Oesophagostomum* spp. populations in Estonia and Denmark. Tartu, 1998, 104 p.
46. **Katrin Heinsoo.** Cuticular and stomatal antechamber conductance to water vapour diffusion in *Picea abies* (L.) karst. Tartu, 1999, 133 p.
47. **Tarmo Annilo.** Studies on mammalian ribosomal protein S7. Tartu, 1998, 77 p.
48. **Indrek Ots.** Health state indicies of reproducing great tits (*Parus major*): sources of variation and connections with life-history traits. Tartu, 1999, 117 p.
49. **Juan Jose Cantero.** Plant community diversity and habitat relationships in central Argentina grasslands. Tartu, 1999, 161 p.
50. **Rein Kalamees.** Seed bank, seed rain and community regeneration in Estonian calcareous grasslands. Tartu, 1999, 107 p.
51. **Sulev Kõks.** Cholecystokinin (CCK) — induced anxiety in rats: influence of environmental stimuli and involvement of endopioid mechanisms and erotonin. Tartu, 1999, 123 p.
52. **Ebe Sild.** Impact of increasing concentrations of O₃ and CO₂ on wheat, clover and pasture. Tartu, 1999, 123 p.
53. **Ljudmilla Timofejeva.** Electron microscopical analysis of the synaptosomal complex formation in cereals. Tartu, 1999, 99 p.
54. **Andres Valkna.** Interactions of galanin receptor with ligands and G-proteins: studies with synthetic peptides. Tartu, 1999, 103 p.
55. **Taavi Virro.** Life cycles of planktonic rotifers in lake Peipsi. Tartu, 1999, 101 p.
56. **Ana Rebane.** Mammalian ribosomal protein S3a genes and intron-encoded small nucleolar RNAs U73 and U82. Tartu, 1999, 85 p.
57. **Tiina Tamm.** Cocksfoot mottle virus: the genome organisation and translational strategies. Tartu, 2000, 101 p.
58. **Reet Kurg.** Structure-function relationship of the bovine papilloma virus E2 protein. Tartu, 2000, 89 p.
59. **Toomas Kivisild.** The origins of Southern and Western Eurasian populations: an mtDNA study. Tartu, 2000, 121 p.
60. **Niilo Kaldalu.** Studies of the TOL plasmid transcription factor XylS. Tartu 2000. 88 p.
61. **Dina Lepik.** Modulation of viral DNA replication by tumor suppressor protein p53. Tartu 2000. 106 p.
62. **Kai Vellak.** Influence of different factors on the diversity of the bryophyte vegetation in forest and wooded meadow communities. Tartu 2000. 122 p.

63. **Jonne Kotta.** Impact of eutrophication and biological invasions on the structure and functions of benthic macrofauna. Tartu 2000. 160 p.
64. **Georg Martin.** Phytobenthic communities of the Gulf of Riga and the inner sea the West-Estonian archipelago. Tartu, 2000. 139 p.
65. **Silvia Sepp.** Morphological and genetical variation of *Alchemilla L.* in Estonia. Tartu, 2000. 124 p.
66. **Jaan Liira.** On the determinants of structure and diversity in herbaceous plant communities. Tartu, 2000. 96 p.
67. **Priit Zingel.** The role of planktonic ciliates in lake ecosystems. Tartu 2001. 111 p.
68. **Tiit Teder.** Direct and indirect effects in Host-parasitoid interactions: ecological and evolutionary consequences. Tartu 2001. 122 p.
69. **Hannes Kollist.** Leaf apoplastic ascorbate as ozone scavenger and its transport across the plasma membrane. Tartu 2001. 80 p.
70. **Reet Marits.** Role of two-component regulator system PehR-PehS and extracellular protease PrtW in virulence of *Erwinia Carotovora* subsp. *Carotovora*. Tartu 2001. 112 p.
71. **Vallo Tilgar.** Effect of calcium supplementation on reproductive performance of the pied flycatcher *Ficedula hypoleuca* and the great tit *Parus major*, breeding in Northern temperate forests. Tartu, 2002. 126 p.
72. **Rita Hõrak.** Regulation of transposition of transposon Tn4652 in *Pseudomonas putida*. Tartu, 2002. 108 p.
73. **Liina Eek-Piirsoo.** The effect of fertilization, mowing and additional illumination on the structure of a species-rich grassland community. Tartu, 2002. 74 p.
74. **Krõõt Aasamaa.** Shoot hydraulic conductance and stomatal conductance of six temperate deciduous tree species. Tartu, 2002. 110 p.
75. **Nele Ingerpuu.** Bryophyte diversity and vascular plants. Tartu, 2002. 112 p.
76. **Neeme Tõnisson.** Mutation detection by primer extension on oligonucleotide microarrays. Tartu, 2002. 124 p.
77. **Margus Pensa.** Variation in needle retention of Scots pine in relation to leaf morphology, nitrogen conservation and tree age. Tartu, 2003. 110 p.
78. **Asko Lõhmus.** Habitat preferences and quality for birds of prey: from principles to applications. Tartu, 2003. 168 p.
79. **Viljar Jaks.** p53 — a switch in cellular circuit. Tartu, 2003. 160 p.
80. **Jaana Männik.** Characterization and genetic studies of four ATP-binding cassette (ABC) transporters. Tartu, 2003. 140 p.
81. **Marek Sammul.** Competition and coexistence of clonal plants in relation to productivity. Tartu, 2003. 159 p.
82. **Ivar Ilves.** Virus-cell interactions in the replication cycle of bovine papillomavirus type 1. Tartu, 2003. 89 p.
83. **Andres Männik.** Design and characterization of a novel vector system based on the stable replicator of bovine papillomavirus type 1. Tartu, 2003. 109 p.

84. **Ivika Ostonen.** Fine root structure, dynamics and proportion in net primary production of Norway spruce forest ecosystem in relation to site conditions. Tartu, 2003. 158 p.
85. **Gudrun Veldre.** Somatic status of 12–15-year-old Tartu schoolchildren. Tartu, 2003. 199 p.
86. **Ülo Väli.** The greater spotted eagle *Aquila clanga* and the lesser spotted eagle *A. pomarina*: taxonomy, phylogeography and ecology. Tartu, 2004. 159 p.
87. **Aare Abroi.** The determinants for the native activities of the bovine papillomavirus type 1 E2 protein are separable. Tartu, 2004. 135 p.
88. **Tiina Kahre.** Cystic fibrosis in Estonia. Tartu, 2004. 116 p.
89. **Helen Orav-Kotta.** Habitat choice and feeding activity of benthic suspension feeders and mesograzers in the northern Baltic Sea. Tartu, 2004. 117 p.
90. **Maarja Öpik.** Diversity of arbuscular mycorrhizal fungi in the roots of perennial plants and their effect on plant performance. Tartu, 2004. 175 p.
91. **Kadri Tali.** Species structure of *Neotinea ustulata*. Tartu, 2004. 109 p.
92. **Kristiina Tambets.** Towards the understanding of post-glacial spread of human mitochondrial DNA haplogroups in Europe and beyond: a phylogeographic approach. Tartu, 2004. 163 p.
93. **Arvi Jõers.** Regulation of p53-dependent transcription. Tartu, 2004. 103 p.
94. **Lilian Kadaja.** Studies on modulation of the activity of tumor suppressor protein p53. Tartu, 2004. 103 p.
95. **Jaak Truu.** Oil shale industry wastewater: impact on river microbial community and possibilities for bioremediation. Tartu, 2004. 128 p.
96. **Maire Peters.** Natural horizontal transfer of the *pheBA* operon. Tartu, 2004. 105 p.
97. **Ülo Maiväli.** Studies on the structure-function relationship of the bacterial ribosome. Tartu, 2004. 130 p.
98. **Merit Otsus.** Plant community regeneration and species diversity in dry calcareous grasslands. Tartu, 2004. 103 p.
99. **Mikk Heidemaa.** Systematic studies on sawflies of the genera *Dolerus*, *Empria*, and *Caliroa* (Hymenoptera: Tenthredinidae). Tartu, 2004. 167 p.
100. **Ilmar Tõnno.** The impact of nitrogen and phosphorus concentration and N/P ratio on cyanobacterial dominance and N₂ fixation in some Estonian lakes. Tartu, 2004. 111 p.
101. **Lauri Saks.** Immune function, parasites, and carotenoid-based ornaments in greenfinches. Tartu, 2004. 144 p.
102. **Siiri Rootsi.** Human Y-chromosomal variation in European populations. Tartu, 2004. 142 p.
103. **Eve Vedler.** Structure of the 2,4-dichloro-phenoxyacetic acid-degradative plasmid pEST4011. Tartu, 2005. 106 p.
104. **Andres Tover.** Regulation of transcription of the phenol degradation *pheBA* operon in *Pseudomonas putida*. Tartu, 2005. 126 p.
105. **Helen Udras.** Hexose kinases and glucose transport in the yeast *Hansenula polymorpha*. Tartu, 2005. 100 p.

106. **Ave Suija.** Lichens and lichenicolous fungi in Estonia: diversity, distribution patterns, taxonomy. Tartu, 2005. 162 p.
107. **Piret Lõhmus.** Forest lichens and their substrata in Estonia. Tartu, 2005. 162 p.
108. **Inga Lips.** Abiotic factors controlling the cyanobacterial bloom occurrence in the Gulf of Finland. Tartu, 2005. 156 p.
109. **Kaasik, Krista.** Circadian clock genes in mammalian clockwork, metabolism and behaviour. Tartu, 2005. 121 p.
110. **Juhan Javoiš.** The effects of experience on host acceptance in ovipositing moths. Tartu, 2005. 112 p.
111. **Tiina Sedman.** Characterization of the yeast *Saccharomyces cerevisiae* mitochondrial DNA helicase Hmi1. Tartu, 2005. 103 p.
112. **Ruth Aguraiuja.** Hawaiian endemic fern lineage *Diellia* (Aspleniaceae): distribution, population structure and ecology. Tartu, 2005. 112 p.
113. **Riho Teras.** Regulation of transcription from the fusion promoters generated by transposition of Tn4652 into the upstream region of *pheBA* operon in *Pseudomonas putida*. Tartu, 2005. 106 p.
114. **Mait Metspalu.** Through the course of prehistory in india: tracing the mtDNA trail. Tartu, 2005. 138 p.
115. **Elin Lõhmussaar.** The comparative patterns of linkage disequilibrium in European populations and its implication for genetic association studies. Tartu, 2006. 124 p.
116. **Priit Kopper.** Hydraulic and environmental limitations to leaf water relations in trees with respect to canopy position. Tartu, 2006. 126 p.
117. **Heili Ilves.** Stress-induced transposition of Tn4652 in *Pseudomonas Putida*. Tartu, 2006. 120 p.
118. **Silja Kuusk.** Biochemical properties of Hmi1p, a DNA helicase from *Saccharomyces cerevisiae* mitochondria. Tartu, 2006. 126 p.
119. **Kersti Püssa.** Forest edges on medium resolution landsat thematic mapper satellite images. Tartu, 2006. 90 p.
120. **Lea Tummeleht.** Physiological condition and immune function in great tits (*Parus major* L.): Sources of variation and trade-offs in relation to growth. Tartu, 2006. 94 p.
121. **Toomas Esperk.** Larval instar as a key element of insect growth schedules. Tartu, 2006. 186 p.
122. **Harri Valdmann.** Lynx (*Lynx lynx*) and wolf (*Canis lupus*) in the Baltic region: Diets, helminth parasites and genetic variation. Tartu, 2006. 102 p.
123. **Priit Jõers.** Studies of the mitochondrial helicase Hmi1p in *Candida albicans* and *Saccharomyces cerevisia*. Tartu, 2006. 113 p.
124. **Kersti Lilleväli.** Gata3 and Gata2 in inner ear development. Tartu, 2007. 123 p.
125. **Kai Rünk.** Comparative ecology of three fern species: *Dryopteris carthusiana* (Vill.) H.P. Fuchs, *D. expansa* (C. Presl) Fraser-Jenkins & Jermy and *D. dilatata* (Hoffm.) A. Gray (Dryopteridaceae). Tartu, 2007. 143 p.

126. **Aveliina Helm.** Formation and persistence of dry grassland diversity: role of human history and landscape structure. Tartu, 2007. 89 p.
127. **Leho Tedersoo.** Ectomycorrhizal fungi: diversity and community structure in Estonia, Seychelles and Australia. Tartu, 2007. 233 p.
128. **Marko Mägi.** The habitat-related variation of reproductive performance of great tits in a deciduous-coniferous forest mosaic: looking for causes and consequences. Tartu, 2007. 135 p.
129. **Valeria Lulla.** Replication strategies and applications of Semliki Forest virus. Tartu, 2007. 109 p.
130. **Ülle Reier.** Estonian threatened vascular plant species: causes of rarity and conservation. Tartu, 2007. 79 p.
131. **Inga Jüriado.** Diversity of lichen species in Estonia: influence of regional and local factors. Tartu, 2007. 171 p.
132. **Tatjana Krama.** Mobbing behaviour in birds: costs and reciprocity based cooperation. Tartu, 2007. 112 p.
133. **Signe Saumaa.** The role of DNA mismatch repair and oxidative DNA damage defense systems in avoidance of stationary phase mutations in *Pseudomonas putida*. Tartu, 2007. 172 p.
134. **Reedik Mägi.** The linkage disequilibrium and the selection of genetic markers for association studies in european populations. Tartu, 2007. 96 p.
135. **Priit Kilgas.** Blood parameters as indicators of physiological condition and skeletal development in great tits (*Parus major*): natural variation and application in the reproductive ecology of birds. Tartu, 2007. 129 p.
136. **Anu Albert.** The role of water salinity in structuring eastern Baltic coastal fish communities. Tartu, 2007. 95 p.
137. **Kärt Padari.** Protein transduction mechanisms of transportans. Tartu, 2008. 128 p.
138. **Siiri-Lii Sandre.** Selective forces on larval colouration in a moth. Tartu, 2008. 125 p.
139. **Ülle Jõgar.** Conservation and restoration of semi-natural floodplain meadows and their rare plant species. Tartu, 2008. 99 p.
140. **Lauri Laanisto.** Macroecological approach in vegetation science: generality of ecological relationships at the global scale. Tartu, 2008. 133 p.
141. **Reidar Andreson.** Methods and software for predicting PCR failure rate in large genomes. Tartu, 2008. 105 p.
142. **Birgot Paavel.** Bio-optical properties of turbid lakes. Tartu, 2008. 175 p.
143. **Kaire Torn.** Distribution and ecology of charophytes in the Baltic Sea. Tartu, 2008, 98 p.
144. **Vladimir Vimberg.** Peptide mediated macrolide resistance. Tartu, 2008, 190 p.
145. **Daima Örd.** Studies on the stress-inducible pseudokinase TRB3, a novel inhibitor of transcription factor ATF4. Tartu, 2008, 108 p.
146. **Lauri Saag.** Taxonomic and ecologic problems in the genus *Lepraria* (*Stereocaulaceae*, lichenised *Ascomycota*). Tartu, 2008, 175 p.

147. **Ulvi Karu.** Antioxidant protection, carotenoids and coccidians in green-finches – assessment of the costs of immune activation and mechanisms of parasite resistance in a passerine with carotenoid-based ornaments. Tartu, 2008, 124 p.
148. **Jaanus Remm.** Tree-cavities in forests: density, characteristics and occupancy by animals. Tartu, 2008, 128 p.
149. **Epp Moks.** Tapeworm parasites *Echinococcus multilocularis* and *E. granulosus* in Estonia: phylogenetic relationships and occurrence in wild carnivores and ungulates. Tartu, 2008, 82 p.
150. **Eve Eensalu.** Acclimation of stomatal structure and function in tree canopy: effect of light and CO₂ concentration. Tartu, 2008, 108 p.
151. **Janne Pullat.** Design, functionlization and application of an *in situ* synthesized oligonucleotide microarray. Tartu, 2008, 108 p.
152. **Marta Putrinš.** Responses of *Pseudomonas putida* to phenol-induced metabolic and stress signals. Tartu, 2008, 142 p.
153. **Marina Semtšenko.** Plant root behaviour: responses to neighbours and physical obstructions. Tartu, 2008, 106 p.
154. **Marge Starast.** Influence of cultivation techniques on productivity and fruit quality of some *Vaccinium* and *Rubus* taxa. Tartu, 2008, 154 p.
155. **Age Tats.** Sequence motifs influencing the efficiency of translation. Tartu, 2009, 104 p.
156. **Radi Tegova.** The role of specialized DNA polymerases in mutagenesis in *Pseudomonas putida*. Tartu, 2009, 124 p.
157. **Tsipe Aavik.** Plant species richness, composition and functional trait pattern in agricultural landscapes – the role of land use intensity and landscape structure. Tartu, 2009, 112 p.
158. **Kaja Kiiver.** Semliki forest virus based vectors and cell lines for studying the replication and interactions of alphaviruses and hepaciviruses. Tartu, 2009, 104 p.
159. **Meelis Kadaja.** Papillomavirus Replication Machinery Induces Genomic Instability in its Host Cell. Tartu, 2009, 126 p.
160. **Pille Hallast.** Human and chimpanzee Luteinizing hormone/Chorionic Gonadotropin beta (*LHB/CGB*) gene clusters: diversity and divergence of young duplicated genes. Tartu, 2009, 168 p.
161. **Ain Vellak.** Spatial and temporal aspects of plant species conservation. Tartu, 2009, 86 p.
162. **Triinu Rimmel.** Body size evolution in insects with different colouration strategies: the role of predation risk. Tartu, 2009, 168 p.
163. **Jaana Salujõe.** Zooplankton as the indicator of ecological quality and fish predation in lake ecosystems. Tartu, 2009, 129 p.
164. **Ele Vahtmäe.** Mapping benthic habitat with remote sensing in optically complex coastal environments. Tartu, 2009, 109 p.
165. **Liisa Metsamaa.** Model-based assessment to improve the use of remote sensing in recognition and quantitative mapping of cyanobacteria. Tartu, 2009, 114 p.

166. **Pille Säälük.** The role of endocytosis in the protein transduction by cell-penetrating peptides. Tartu, 2009, 155 p.
167. **Lauri Peil.** Ribosome assembly factors in *Escherichia coli*. Tartu, 2009, 147 p.
168. **Lea Hallik.** Generality and specificity in light harvesting, carbon gain capacity and shade tolerance among plant functional groups. Tartu, 2009, 99 p.
169. **Mariliis Tark.** Mutagenic potential of DNA damage repair and tolerance mechanisms under starvation stress. Tartu, 2009, 191 p.
170. **Riinu Rannap.** Impacts of habitat loss and restoration on amphibian populations. Tartu, 2009, 117 p.
171. **Maarja Adojaan.** Molecular variation of HIV-1 and the use of this knowledge in vaccine development. Tartu, 2009, 95 p.
172. **Signe Altmäe.** Genomics and transcriptomics of human induced ovarian folliculogenesis. Tartu, 2010, 179 p.
173. **Triin Suvi.** Mycorrhizal fungi of native and introduced trees in the Seychelles Islands. Tartu, 2010, 107 p.
174. **Velda Lauringson.** Role of suspension feeding in a brackish-water coastal sea. Tartu, 2010, 123 p.
175. **Eero Talts.** Photosynthetic cyclic electron transport – measurement and variably proton-coupled mechanism. Tartu, 2010, 121 p.
176. **Mari Nelis.** Genetic structure of the Estonian population and genetic distance from other populations of European descent. Tartu, 2010, 97 p.
177. **Kaarel Krjutškov.** Arrayed Primer Extension-2 as a multiplex PCR-based method for nucleic acid variation analysis: method and applications. Tartu, 2010, 129 p.
178. **Egle Köster.** Morphological and genetical variation within species complexes: *Anthyllis vulneraria* s. l. and *Alchemilla vulgaris* (coll.). Tartu, 2010, 101 p.
179. **Erki Õunap.** Systematic studies on the subfamily Sterrhinae (Lepidoptera: Geometridae). Tartu, 2010, 111 p.
180. **Merike Jõesaar.** Diversity of key catabolic genes at degradation of phenol and *p*-cresol in pseudomonads. Tartu, 2010, 125 p.
181. **Kristjan Herkül.** Effects of physical disturbance and habitat-modifying species on sediment properties and benthic communities in the northern Baltic Sea. Tartu, 2010, 123 p.
182. **Arto Pulk.** Studies on bacterial ribosomes by chemical modification approaches. Tartu, 2010, 161 p.
183. **Maria Põllupüü.** Ecological relations of cladocerans in a brackish-water ecosystem. Tartu, 2010, 126 p.
184. **Toomas Silla.** Study of the segregation mechanism of the Bovine Papillomavirus Type 1. Tartu, 2010, 188 p.
185. **Gyaneshwer Chaubey.** The demographic history of India: A perspective based on genetic evidence. Tartu, 2010, 184 p.
186. **Katrin Kepp.** Genes involved in cardiovascular traits: detection of genetic variation in Estonian and Czech populations. Tartu, 2010, 164 p.

187. **Virve Sõber.** The role of biotic interactions in plant reproductive performance. Tartu, 2010, 92 p.
188. **Kersti Kangro.** The response of phytoplankton community to the changes in nutrient loading. Tartu, 2010, 144 p.
189. **Joachim M. Gerhold.** Replication and Recombination of mitochondrial DNA in Yeast. Tartu, 2010, 120 p.
190. **Helen Tammert.** Ecological role of physiological and phylogenetic diversity in aquatic bacterial communities. Tartu, 2010, 140 p.
191. **Elle Rajandu.** Factors determining plant and lichen species diversity and composition in Estonian *Calamagrostis* and *Hepatica* site type forests. Tartu, 2010, 123 p.
192. **Paula Ann Kivistik.** ColR-ColS signalling system and transposition of Tn4652 in the adaptation of *Pseudomonas putida*. Tartu, 2010, 118 p.
193. **Siim Sõber.** Blood pressure genetics: from candidate genes to genome-wide association studies. Tartu, 2011, 120 p.
194. **Kalle Kipper.** Studies on the role of helix 69 of 23S rRNA in the factor-dependent stages of translation initiation, elongation, and termination. Tartu, 2011, 178 p.
195. **Triinu Siibak.** Effect of antibiotics on ribosome assembly is indirect. Tartu, 2011, 134 p.
196. **Tambet Tõnissoo.** Identification and molecular analysis of the role of guanine nucleotide exchange factor RIC-8 in mouse development and neural function. Tartu, 2011, 110 p.
197. **Helin Räägel.** Multiple faces of cell-penetrating peptides – their intracellular trafficking, stability and endosomal escape during protein transduction. Tartu, 2011, 161 p.
198. **Andres Jaanus.** Phytoplankton in Estonian coastal waters – variability, trends and response to environmental pressures. Tartu, 2011, 157 p.
199. **Tiit Nikopensius.** Genetic predisposition to nonsyndromic orofacial clefts. Tartu, 2011, 152 p.
200. **Signe Värvi.** Studies on the mechanisms of RNA polymerase II-dependent transcription elongation. Tartu, 2011, 108 p.
201. **Kristjan Välik.** Gene expression profiling and genome-wide association studies of non-small cell lung cancer. Tartu, 2011, 98 p.
202. **Arno Põllumäe.** Spatio-temporal patterns of native and invasive zooplankton species under changing climate and eutrophication conditions. Tartu, 2011, 153 p.
203. **Egle Tammeleht.** Brown bear (*Ursus arctos*) population structure, demographic processes and variations in diet in northern Eurasia. Tartu, 2011, 143 p.
205. **Teele Jairus.** Species composition and host preference among ectomycorrhizal fungi in Australian and African ecosystems. Tartu, 2011, 106 p.
206. **Kessy Abarenkov.** PlutoF – cloud database and computing services supporting biological research. Tartu, 2011, 125 p.

207. **Marina Grigorova.** Fine-scale genetic variation of follicle-stimulating hormone beta-subunit coding gene (*FSHB*) and its association with reproductive health. Tartu, 2011, 184 p.
208. **Anu Tiitsaar.** The effects of predation risk and habitat history on butterfly communities. Tartu, 2011, 97 p.
209. **Elin Sild.** Oxidative defences in immunoeological context: validation and application of assays for nitric oxide production and oxidative burst in a wild passerine. Tartu, 2011, 105 p.
210. **Irja Saar.** The taxonomy and phylogeny of the genera *Cystoderma* and *Cystodermella* (Agaricales, Fungi). Tartu, 2012, 167 p.
211. **Pauli Saag.** Natural variation in plumage bacterial assemblages in two wild breeding passerines. Tartu, 2012, 113 p.
212. **Aleksei Lulla.** Alphaviral nonstructural protease and its polyprotein substrate: arrangements for the perfect marriage. Tartu, 2012, 143 p.
213. **Mari Järve.** Different genetic perspectives on human history in Europe and the Caucasus: the stories told by uniparental and autosomal markers. Tartu, 2012, 119 p.
214. **Ott Scheler.** The application of tmRNA as a marker molecule in bacterial diagnostics using microarray and biosensor technology. Tartu, 2012, 93 p.
215. **Anna Balikova.** Studies on the functions of tumor-associated mucin-like leukosialin (CD43) in human cancer cells. Tartu, 2012, 129 p.
216. **Triinu Kõressaar.** Improvement of PCR primer design for detection of prokaryotic species. Tartu, 2012, 83 p.
217. **Tuul Sepp.** Hematological health state indices of greenfinches: sources of individual variation and responses to immune system manipulation. Tartu, 2012, 117 p.
218. **Rya Ero.** Modifier view of the bacterial ribosome. Tartu, 2012, 146 p.
219. **Mohammad Bahram.** Biogeography of ectomycorrhizal fungi across different spatial scales. Tartu, 2012, 165 p.
220. **Annely Lorents.** Overcoming the plasma membrane barrier: uptake of amphipathic cell-penetrating peptides induces influx of calcium ions and downstream responses. Tartu, 2012, 113 p.
221. **Katrin Männik.** Exploring the genomics of cognitive impairment: whole-genome SNP genotyping experience in Estonian patients and general population. Tartu, 2012, 171 p.
222. **Marko Prouš.** Taxonomy and phylogeny of the sawfly genus *Empria* (Hymenoptera, Tenthredinidae). Tartu, 2012, 192 p.
223. **Triinu Visnapuu.** Levansucrases encoded in the genome of *Pseudomonas syringae* pv. tomato DC3000: heterologous expression, biochemical characterization, mutational analysis and spectrum of polymerization products. Tartu, 2012, 160 p.
224. **Nele Tamberg.** Studies on Semliki Forest virus replication and pathogenesis. Tartu, 2012, 109 p.

225. **Tõnu Esko.** Novel applications of SNP array data in the analysis of the genetic structure of Europeans and in genetic association studies. Tartu, 2012, 149 p.
226. **Timo Arula.** Ecology of early life-history stages of herring *Clupea harengus membras* in the northeastern Baltic Sea. Tartu, 2012, 143 p.
227. **Inga Hiiesalu.** Belowground plant diversity and coexistence patterns in grassland ecosystems. Tartu, 2012, 130 p.
228. **Kadri Koorem.** The influence of abiotic and biotic factors on small-scale plant community patterns and regeneration in boreonemoral forest. Tartu, 2012, 114 p.
229. **Liis Andresen.** Regulation of virulence in plant-pathogenic pectobacteria. Tartu, 2012, 122 p.
230. **Kaupo Kohv.** The direct and indirect effects of management on boreal forest structure and field layer vegetation. Tartu, 2012, 124 p.
231. **Mart Jüssi.** Living on an edge: landlocked seals in changing climate. Tartu, 2012, 114 p.
232. **Riina Klais.** Phytoplankton trends in the Baltic Sea. Tartu, 2012, 136 p.
233. **Rauno Veeroja.** Effects of winter weather, population density and timing of reproduction on life-history traits and population dynamics of moose (*Alces alces*) in Estonia. Tartu, 2012, 92 p.
234. **Marju Keis.** Brown bear (*Ursus arctos*) phylogeography in northern Eurasia. Tartu, 2013, 142 p.
235. **Sergei Põlme.** Biogeography and ecology of *alnus*- associated ectomycorrhizal fungi – from regional to global scale. Tartu, 2013, 90 p.
236. **Liis Uusküla.** Tartu, 2013, 173 p.
237. **Marko Lõoke.** Studies on DNA replication initiation in *Saccharomyces cerevisiae*. Tartu, 2013, 112 p.
238. **Anne Aan.** Light- and nitrogen-use and biomass allocation along productivity gradients in multilayer plant communities. Tartu, 2013, 127 p.
239. **Heidi Tamm.** Comprehending phylogenetic diversity – case studies in three groups of ascomycetes. Tartu, 2013, 136 p.
240. **Liina Kangur.** High-Pressure Spectroscopy Study of Chromophore-Binding Hydrogen Bonds in Light-Harvesting Complexes of Photosynthetic Bacteria. Tartu, 2013, 150 p.
241. **Margus Leppik.** Substrate specificity of the multisite specific pseudouridine synthase RluD. Tartu, 2013, 111 p.
242. **Lauris Kaplinski.** The application of oligonucleotide hybridization model for PCR and microarray optimization. Tartu, 2013, 103 p.
243. **Merli Pärnoja.** Patterns of macrophyte distribution and productivity in coastal ecosystems: effect of abiotic and biotic forcing. Tartu, 2013, 155 p.
244. **Tõnu Margus.** Distribution and phylogeny of the bacterial translational GTPases and the Mqsr/YgiT regulatory system. Tartu, 2013, 126 p.