DISSERTATIONES BIOLOGICAE UNIVERSITATIS TARTUENSIS 205

DISSERTATIONES BIOLOGICAE UNIVERSITATIS TARTUENSIS 205

TEELE JAIRUS

Species composition and host preference among ectomycorrhizal fungi in Australian and African ecosystems



TARTU UNIVERSITY PRESS

Department of Botany, Institute of Ecology and Earth Sciences, Faculty of Science and Technology, University of Tartu, Estonia

Dissertation was accepted for the commencement of the degree of *Doctor philosophiae* in Botany and Mycology at the University of Tartu on 13th of June 2011 by the Scientific Council of the Institute of Ecology and Earth Sciences University of Tartu.

Supervisor: Dr. Leho Tedersoo, University of Tartu, Estonia

Opponent: Dr. Taina Pennanen, The Finnish Forest Research Institute, Finland

Commencement: Room 1019, 14A Ravila Street, Tartu, on 23 September 2011 at 12.15 p.m.

Publication of this thesis is granted by the Institute of Ecology and Earth Sciences, University of Tartu and by the Doctoral School of Earth Sciences and Ecology created under the auspices of European Social Fund.





European Union European Social Fund

Investing in your future

ISSN 1024–6479 ISBN 978–9949–19–843–6 (trükis) ISBN 978–9949–19–844–3 (PDF)

Autoriõigus Teele Jairus, 2011

Tartu Ülikooli Kirjastus www.tyk.ee Tellimus nr 512

CONTENTS

LIST OF ORIGINAL PUBLICATIONS	6
INTRODUCTION	7
MATERIALS AND METHODS Study sites Molecular analyses Phylogenetic analyses Statistical analyses	10 10 11 13 13
RESULTS	15
DISCUSSION	18
CONCLUSIONS	21
REFERENCES	22
SUMMARY IN ESTONIAN	27
ACKNOWLEDGEMENTS	29
PUBLICATIONS	31
CURRICULUM VITAE	92

LIST OF ORIGINAL PUBLICATIONS

This thesis is based on the following papers that are referred to in the text by Roman numerals:

- Tedersoo L, Jairus T, Horton BM, Abarenkov A, Suvi T, Saar I, Kõljalg U. 2008. Strong host preference of ectomycorrhizal fungi in a Tasmanian wet sclerophyll forest as revealed by DNA barcoding and taxon-specific primers. *New Phytologist* 180: 479–490.
- II. Tedersoo L, Gates G, Dunk CW, Lebel T, May TW, Kõljalg U, Jairus T. 2009. Establishment of ectomycorrhizal fungal community on isolated *Nothofagus cunninghamii* seedlings regenerating on dead wood in Australian wet temperate forests: does fruit-body type matter? *Mycorrhiza* 19: 403–416.
- III. Tedersoo L, Bahram M, Jairus T, Bechem E, Chinoya S, Mpumba R, Leal M, Randrianjohany E, Razafimandimbison S, Sadam A, Naadel T, Kõljalg U. 2011. Spatial structure and the effects of host and soil environments on communities of ectomycorrhizal fungi in wooded savannas and rainforests of Continental Africa and Madagascar. *Molecular Ecology*: in press. doi: 10.1111/j.1365-294X.2011.05145.x
- IV. Jairus T, Mpumba R, Chinoya S, Tedersoo L. 2011. Invasion potential and host shifts of Australian and African ectomycorrhizal fungi in mixed eucalypt plantations. *New Phytologist*: in press. doi: 10.1111/j.1469-8137.2011.03775.x

	Ι	II	III	IV
Idea and design				*
Sampling			*	*
Molecular analysis	*	*	*	*
Data analysis				*
Writing	*	*	*	*

Author's contribution to each paper

INTRODUCTION

Mycorrhiza is a mutualistic association between plant roots and fungi. Fungi receive carbon from plants and plants get mineral nutrients in return (Smith & Read 1997). The most common mycorrhiza types are arbuscular mycorrhiza and ectomycorrhiza (EcM). About 80% of studied plant species and 92% of plant families are mycorrhizal (Wang & Qiu 2006), of which 3% form EcM (Taylor & Alexander 2005). EcM is formed mainly between woody plants and fungi from the phyla Ascomycota or Basidiomycota. In temperate regions, the majority of trees are ectomycorrhizal, whereas arbuscular mycorrhiza dominates in most tropical plants. EcM trees tend to form monodominant patches in tropical forests (Connell & Lowman 1989, Alexander & Lee 2005). The symbiosis has specific anatomical appearance: root tips of plants are covered with a fungal sheath termed as 'mantle'. In addition, fungal hyphae surround epidermal or cortical cells of root tips of plants, forming a structure termed as a 'Hartig net'.

Ectomycorrhizal trees are usually able to form symbiotic associations with several different fungal species. For example, *Pseudotsuga menziesii* hosts about 2000 species of EcM fungi throughout its range (Trappe 1977), whereas a single *Populus tremula* tree may harbor more than a hundred fungal species (Bahram et al. 2011). Most EcM fungi are able to associate with multiple hosts simultaneously, but many subdominant species show high levels of host preference or even strict specificity (Molina et al. 1992; Horton & Bruns 1998). Fungi from the genera *Suillus* and *Rhizopogon* are strongly specific to tree species from the *Pinaceae* family (Molina et al. 1992, Collier & Bidartondo 2009, Nunez et al. 2009), whereas *Alnus* spp. associate with only a limited number of fungal species (Molina et al. 1992, Tedersoo et al. 2009).

Ectomycorrhizal symbiosis has independently evolved >60 times in fungi. These monophyletic groups of fungi are referred to as 'lineages' and they correspond to section- to order-level taxa in the fungal tree of life (Tedersoo et al. 2010a). At the global scale, the most species-rich EcM fungal lineages are /tomentella-thelephora and /russula-lactarius, followed by /cortinarius, /sebacina, /clavulina, /boletus and /inocybe (Tedersoo & Nara 2010). In Australia, fruit-body surveys have revealed that the EcM lineages /cortinarius, /descolea and /russula-lactarius are the most species-rich (Claridge et al. 1999, Lu et al. 1999, Bougher & Lebel 2001, Gates et al. 2005, Ratkowsky & Gates 2005), whereas soil mycelial communities are dominated by /russula-lactarius, /tomentella-thelephora, followed by /cortinarius and /inocybe (Bastias et al. 2006, Midgley et al. 2007). In tropical forests, the lineages of /russula-lactarius, /tomentella-thelephora and /boletus dominate on root systems of host trees (Diédhiou et al. 2010, Peay et al. 2010).

Ectomycorrhizal fungal species composition and diversity in a certain site is also influenced by the presence of different forest microsites (Tedersoo et al. 2003). Some EcM plants prefer to germinate in pits, root mounds and decaying logs (Hofgaard 1993). The EcM fungal community in dead wood differs from the soil EcM fungal community (Tedersoo et al. 2003, 2008). Due to the higher moisture content and other favorable microclimatic conditions, dead wood provides a suitable substrate for proliferation of roots and growth of EcM fungi (Harvey et al. 1978). In boreal forests, dead wood-inhabiting seedlings host a limited number of EcM fungi (Iwanski & Rudawska 2007) and their community structure depends on tree species and the presence of root connections with mature host trees (Tedersoo et al. 2008). Tedersoo et al. (2008) observed that in Estonian boreal forests, isolated seedlings harbor only a few EcM fungal species that belong to the orders Atheliales and Thelephorales that predominately form resupinate fruit-bodies.

Forest plantations have become an increasingly important supply for wood during the era of rapid deforestation of primary habitats (Richardson 1998). In 2010, plantations covered about 264 million ha globally and contributed about 7% to the world's forested areas (FAO 2010). Exotic trees are preferred over native ones due to their shorter rotation period, well-studied biology and the lack of pests in new habitats. Among hundreds of tree species used, Eucalyptus, Pinus and Acacia dominate in forest plantations (Richardson 1998, West 2006). When trees are transplanted to new localities, some of them grow well, whereas others thrive only after co-introduction of their native fungi. Eucalyptus trees grow better with their naturally symbiotic fungi compared to fungi from other regions (Chen et al. 2007). Eucalyptus spp. are, nevertheless, able to grow without their naturally symbiotic fungi (Tedersoo et al. 2007) and establish symbiosis with native fungi that are available in exotic habitats (Yazid 1994, Tedersoo et al. 2007, Buyck 2008). *Pinus* spp., by contrast, are not able to grow without their naturally symbiotic fungi (Mikola 1969), and they cannot form symbiosis with local fungi in exotic tropical or southern temperate habitats (Tedersoo et al. 2007, Nunez et al. 2009, Dickie et al. 2010, Walbert et al. 2010). When Pinus spp. are growing with co-introduced fungi in suitable habitats, they may naturalize and become invasive in suitable climatic conditions (Nunez et al. 2009).

It is relatively easy to follow the fate of introduced large organisms such as trees and mammals in novel habitats, but the establishment and spread of microbes may easily go unnoticed. The behavior of pathogenic fungi in novel habitats is usually detected through the damage they cause. Symbiotic and commensal fungi do not cause easily detectable ecological damage and their presence remains difficult to detect. It may take a lot of time before the introduced species become naturalized or detectable. For example, the fly-agaric *Amanita muscaria*, which was co-introduced with pines, broadened its host range from northern temperate plant families to *Nothofagus* species and gained the capacity to complete its life cycle with its novel host species in New Zealand and Australia a few hundred years later (Dunk 2002, Orlovich & Cairney 2004). A species of *Laccaria*, introduced along with *Eucalyptus* from Australia, has become naturalized on root systems of some native plant species

in the Iberian Peninsula (Diez 2005). In Madagascar, introduced fungi have provided a new food source for local people in *Eucalyptus* plantations (Buyck 2008). In several parts of the world, local fungal species have been found fruiting in exotic *Eucalyptus* plantations, indicating that host shifts are possible in both native and introduced fungi (Chen et al. 2007, Tedersoo et al. 2007, Buyck 2008).

In my thesis, the following main research hypotheses were tested: 1) Australian and African EcM fungal communities differ in species richness and phylogenetic community composition; 2) EcM fungi display differential host preference in African, Australian and temperate ecosystems; 3) EcM fungi are capable of shifting hosts especially when roots of native and introduced trees intermingle; and 4) EcM fungi with resupinate fruit-bodies preferentially inhabit tree roots in dead wood compared to forest floor soil.

MATERIALS AND METHODS

Study sites

In all study sites, soil cores (15x15 cm to 5 (I, II) or 10 (III, IV) cm depth) were collected from below potential host trees. In addition, fruit-bodies of all putatively EcM fungi were collected, vouchered and deposited in the herbarium of Tartu University (TU). Sampling in different studies is briefly described below (see methods of each study for detailed information).

Study I

Samples were collected from wet sclerophyll forest in Tall Trees Walk, Mt Field National Park, Tasmania (42°40.9'S; 146°42.2'E), where three 1-ha plots were established in sites, where at least three EcM host trees– *Eucalyptus regnans* F. Müll., *Nothofagus cunninghamii* (Hook.) Oerst and *Pomaderris apetala* Labill. - co-occured. Sampling was performed in August, 2006.

Study II

Samples were collected from two sites. The 1-ha Yarra site is located in the cool temperate rain forest dominated by *Nothofagus cunninghamii* at Acheron Gap in the Yarra Ranges National Park ($37^{\circ}41$ 'S $145^{\circ}44$ 'E, altitude 770 m), Victoria, Australia. The 600-m² Warra site is located in a wet sclerophyll vegetation dominated by *Eucalyptus obliqua* L'Hér a site at Warra LTER near Tahune ($43^{\circ}04$ 'S; $146^{\circ}40$ 'E, altitude 155 m), Geeveston, Tasmania, Australia. Soil samples were collected from below mature *N. cunninghamii*. In addition, seedlings (1–5 years old) of *N. cunninghamii* were sampled from decaying logs of medium decay classes (III and IV sensu Christy & Mack 1984). Up to two *Nothofagus* seedlings were sampled from the same log at least 5 m distance. The absence of mature tree roots in seedlings rooting zone was confirmed by digging the wood with a sharp knife.

Study III

Four study sites were established in African ecoregions that support many EcM hosts. The Kashima site is located in an ecotone of miombo (woodland) and dambo (seasonally flooded wetland) ecosystems in Kabompo district, NW Zambia (13°19'S; 24°30'E, altitude 1080 m). Sampling was performed in a 4-ha plot in January, 2008. The Tchimbele study site is in primary lowland rainforest in Mbé National Park in Littoral province of North-Western Gabon (0°37'N; 10°24'E, altitude 400 m). Sampling was performed in a 20-ha plot in May, 2009. The Mandena site lies in a littoral forest in Toliara province of SE Madagascar (24°58'S; 47°00'E, altitude 40 m). Sampling was performed in a 2-ha plot in March, 2010. The Korup site represents a primary lowland rainforest in Korup

National Park in Southwest Province of Cameroon (5°01'N; 8°48'E, altitude 120 m). Sampling was performed in a 12-ha plot in April, 2008.

Study IV

Samples were collected from two neighbouring plantations of *Eucalyptus grandis* W. Hill ex Maiden and *Eucalyptus camaldulensis* Dehnh, where native trees form an understorey. The plantations are located near Tumbama village, Kabompo district, Northwestern Province of Zambia (13°28'S; 24°25'E, altitude 1077 m). Two 2-ha plots were established in these plantations in January, 2008.

Molecular analyses

Fungal and plant species were identified based on molecular methods. In all studies, the aim was to maximize the number of identified species and to optimize the DNA extraction methodology. Genomic DNA was extracted from 1-6 root tips per sample using a Qiagen MagAttract 96 DNA Plant Kit or Qiagen Dneasy 96 Plant Kit for studies III and IV, and High Pure polymerase chain reaction (PCR) Template Preparation Kit for Isolation of Nucleic Acids from Mammalian Tissue for studies I and II. Nuclear rDNA ITS and Large Subunit (LSU) regions were amplified with PCR-methodology using respective primers (Table 1). To obtain the best success from amplification, new primers were developed to target specific fungal taxa. For amplification, PuReTaq Ready-To-Go PCR kit (GE Healthcare UK Limited, Little Chalfont, Buckinghamshire, UK) was used. For each reaction, 24 µl DNA extract and 0.5 µl of each primer were added. PCR products were checked on 1% agarose gels under UVlight and purified using Exo-Sap enzymes (Sigma, St Louis, MO, USA). DNA sequencing was performed using primers ITS4 (5'-tcctccgcttattgatatgc-3'), ITS5 (5'-ggaagtaaaagtcgtaacaagg-3') and/or LF340 (5'-tacttgtkcgctatcgg-3') for the and ctb6 (5'-gcatatcaataagcggagg-3') and/or LR5 ITS region. (5'tcctgagggaaacttcg-3') for the LSU. Sequencing was performed in Macrogen Inc. (Seoul, South Korea or Amsterdam, The Netherlands). Sequences were assembled into contigs and checked for quality with Sequencher 4.7 or 4.9 software (GeneCodes Corp., Ann Arbor, MI, USA).

DNA sequences were assigned to molecular species based on 97% similarity (excluding flanking rDNA sequences). For the lineages of /cortinarius and /laccaria, a 98% similarity was used instead (I). Species were identified by running bulk megablast queries against the International Sequence Database (INSD) and UNITE (Abarenkov et al. 2010a) as implemented in the PlutoF workbench (Abarenkov et al. 2010b). Host plant species were identified from the same DNA extract as fungi based on the plastid *trnL* region. DNA was amplified using primers trnC (5'-cgaaatcggtagacgtagacgactgaac-3') following the same protocol as for fungi. Plant DNA was sequenced using a primer trnD. Plant species were identified based on the size differences (I, II) or sequence similarity (III, IV) of the *trnL* region.

Primer	Sequence	DNA region	Target	Study	References
ITS1-F	5'-cttggtcatttagaggaagtaa-3'	ITS	Fungal	I, II	Gardes & Bruns 1993
ITS2	5'-gctgcgttcttcatcgatgc-3'	ITS	Universal	II, I<	White et al. 1990
ITS4	5'-tcctccgcttattgatatgc-3'	ITS	Universal	II, I<	White et al. 1990
ITS4B	5'-caggagacttgtacacggtccag-3'	ITS	Agaricales, Boletales	≡	Gardes & Bruns 1993
ITS4-Clav	5'-ggtagtcccacctgattc-3'	ITS	/clavulina	≡	Study III
ITS4–Russ	5'-agcgggtagtctcaccc-3'	ITS	/russula-lactarius	≡	Study III
ITS4-Seb	5'-tcagcgggtartcctactc-3'	ITS	Sebacinales	≡	Study III
ITS4-Sord	5'-cccgttccagggaactc-3'	ITS	Sordariomycetes	I, III	Study I
ITS4-Tom	5'-aactcggacgaccagaggca-3'	ITS	/tomentella-thelephora	≡	Study III
ITS5	5'-ggaagtaaaagtcgtaacaagg-3'	ITS	Universal	_	White et al. 1990
ITSOF-T	5'-acttggtcatttagaggaagt-3'	ITS	Fungal	I, III, I∕	Taylor & McCormick 2008
LA-W	5'cttttcatctttcgatcactc- 3'	ITS	Ascomycota	I, II, III	Study I
LB-Z	5' aaaaatggcccactagaaact 3'	nLSU	Basidiomycota	I, II, III	Study I
LB-W	5'-cttttcatctttccctcacgg-3'	ITS	Basidiomycota	I, II, III, IV	Study I
LB-Y	5'-ggtccgtgtttcaagacg-3'	nrsu	Basidiomycota	_	Study I
LROR	5'-acccgctgaacttaagc-3'	nLSU	Universal	I, II, III	Hopple & Vilgalys 1994
LR21–Cer	5'-cgactcgttgagagcacaa-3'	nrsu	Ceratobasidiaceae	≡	Study III
LR3-Asc	5'-cacytactcaaatccwagcg-3'	nrsu	Ascomycota	I, II	Study I
LR3–Pez	5'-cmtcrggatcggtcgatgg-3'	nrsu	Pezizales	≡	Study I
LR3–Tom	5'-ctaccgtagaaccgtctcc-3'	nrsu	/tomentella-thelephora	≡	Study I
LR3-Tul	5'-bactcgcatgcaaggtgca-3'	ITS	Tulasnellales	_	Study I
LR5	5'-tcctgagggaaacttcg-3'	nLSU	Universal	≡	Vilgalys & Hester 1990
LR5-F	5'-cgatcgatttgcacgtcaga-3'	nrsu	Fungal	_	Study I
LR5-Seb	5'-attcgctttaccgcacaagg-3'	ITS	Sebacinales	_	Study I
LR5-Tom	5'-ctaccgtagaaccgtctcc-3'	ITS	/tomentella-thelephora	I, III, I<	Study I
ML5	5'-ctcggaaattatcctcataag-3'	mtLSU	Fungal	=	Bruns et al. 1998
ML6	5'-cagtagaagctgcatagggtc-3'	mtLSU	Fungal	=	Bruns et al. 1998

Table 1 PCR primers used in different studies

Phylogenetic analyses

Phylogenetic analyses were prepared to identify the potential origin of EcM fungi (IV), assuming that closest relatives are likely to originate from the same continent (Hosaka et al. 2008, Matheny et al. 2009). Separate phylogenetic analyses were performed for different EcM fungal lineages (cf. Tedersoo et al. 2010a). Alignments were conducted using sequences from different study sites in Africa and all relevant INSD and UNITE sequences. The phylogenetic analyses of the /tomentella-thelephora and /russula-lactarius lineages were restricted to tropical and Southern Hemisphere isolates, because of the data set size and distinction of the northern boreal and tropical clades as seen from the other lineages. The alignments of ITS sequences were constructed using Mafft ver. 6 (Katoh & Toh 2008) and corrected manually. Unrooted Maximum Likelihood phylograms were constructed using RAxML with 200 bootstrap replicates and default options (Stamatakis et al. 2008). Rapid bootstrap values were used to estimate the robustness of branches and the potential origin of isolates. Isolates were considered to be native African if the same species were found in natural African habitats or they clustered with other African and/or SE Asian isolates with strong support (>70% BS). Conversely, taxa were regarded as Australian if they belonged to EcM lineages absent from the native vegetation of Sub-Saharan Africa, occurred in other eucalypt plantations, and/or belonged to the strongly supported Austral (incl. Australia, New Zealand, New Caledonia and Argentina) clades. Isolates that matched none of these criteria were conservatively assigned to uncertain origin.

Statistical analyses

Fisher's Exact tests were calculated to assess the occurrence of host preference (I, III, IV) and the effect of nearest host (I) on EcM fungi at the species and family levels of host trees. Species accumulation curves were calculated using a computer program EstimateS ver. 8 (Colwell 2006) to compare species richness. Moran's I was calculated to establish whether spatial factors cause biases in fungal distribution. Detrended Correspondence Analysis (DCA) using PC-ORD ver. 5.04 (McCune & Mefford 2006) was employed to reveal the effects of host species, host vicinity and plot on EcM fungal community structure (I). To address the relative importance of spatial effects on the EcM fungal communities, vectors of Principal Coordinates of Neighbor Matrices (PCNM) were constructed according to Borcard & Legendre (2002)(III). PCNM vectors were calculated based on the geographical distance matrix and ordered by coefficients of determination based on forward selection and 1,000 permutations as implemented in the Pakfor package of R (R Core Development Team 2007). All significant (P<0.05) PCNM vectors were included in the distance matrix for multivariate analysis. The relative effects of the PCNM vectors, host and soil horizons, were addressed based on Bray-Curtis distance measure as implemented in Adonis permutational multivariate analysis of variance function of the Vegan package of R (III). For the fourth study (IV), the relative effects of host were assessed by use of Adonis. Based on the same options, Non-metric Multidimensional Scaling (NMS) plots were constructed to visualize the relative effects of these variables on EcM fungal communities (III, IV). Vectors and centroids of each variables were fitted into the NMS plots using function envfit. Significance level of $\alpha < 0.05$ was used throughout the studies. In study II, two-way ANOVA was performed to study the effects of site and substrate on species density of EcM fungi. Mann- Whitney U tests and Fischer's exact tests were calculated to address the differences in relative abundance of EcM fungal lineages and frequency of species, respectively, between forest floor soil and dead wood. To control false discovery rate and reduce family wise error rate arising with multiple statistical testing, obtained P values were subjected to a sharpening procedure of Benjamin-Hochberg correction (Verhoeven et al. 2005). A canonical multivariate analysis was performed to evaluate differences in fungal communities between forest floor soil and dead wood and between sites using a computer program CAP (Anderson & Willis 2003). To test whether EcM fungal taxa with resupinate and non-resupinate fruit bodies occur at similar frequency in dead wood or not, a chi-square test was used. To further address potential phylogenetical biases among EcM fungal communities among substrates, G-tests using species frequency data were performed for nine most common EcM lineages as implemented in computer program Unifrac (Lozupone et al. 2006). For this analysis, available LSU sequences from all species and ITS sequences from /cortinarius and /laccaria were aligned using Mafft 5.861 (Katoh et al. 2005) and corrected manually. Eight species lacking >250 bp LSU sequence and *Tulasnella* sp3 were removed from the analysis because of the presence of long, unstable branches and unstable position. A parsimony analysis using 100 random start generations and TBR branch swapping was run in PAUP* 4.0d81 (Swofford 2002). One of the 21,834 most parsimony trees was randomly selected as an input to Unifrac. The P values of G tests were subjected to the Benjamini-Hochberg correction.

RESULTS

In Mt. Field site, Australia (I), 123 species of EcM fungi were recovered. The majority of species (55.3%) were found only once. Laccaria sp. 1 and Lactarius eucalypti were the most frequent. The /cortinarius (28 spp.), /tomentellathelephora (18 spp.) and /russula-lactarius (10 spp.) were the most species-rich lineages of EcM fungi. At Warra and Yarra, respectively, 86 and 25 species of EcM fungi were identified on the roots of Nothofagus cunninghamii. The lineage of /cortinarius dominated in both sites, followed by /clavulina and /russula-lactarius at Warra, and /descolea and /tomentella-thelephora at Yarra. In Africa, 94, 101, 46, 111 and 60 species of EcM fungi were recovered in Kashima, Tchimbele, Mandena, Korup and Tumbama, respectively. The lineages of /russula-lactarius, /tomentella-thelephora and /boletus dominated in all sites except Tumbama, where the third most species-rich lineage was /clavulina instead of /boletus. Because of differential sampling effort, the number of species is not directly comparable among sites. Species accumulation curves suggest that Australian sites comprise more species than African sites (Figure 1).



Figure 1 Rarefied species accumulation curves (solid lines) and their 95% confidence intervals (pointed lines) of ectomycorrhizal fungi at the seven study sites.

Australian EcM fungi displayed strong preference for host plants (I). Nineteen species (65.5%) were significantly affected by host plant roots in Mt Field site (Figure 2). Host species effect was stronger than host vicinity effect in 12 out of 15 cases (I). Plot had no significant effect on any species. However, host had no effect on the frequency of the EcM fungal species in Africa (III). Only in Tchimbele (III), five species showed significantly biased association towards host species, but in most cases (except one) the effect was explained by spatial autocorrelation. In the mixed eucalypt plantation in Tumbama (IV), three species of fungi with Australian origin showed host preference for their native eucalypt hosts. In all African and Australian sites, host trees did not differ in the richness of associated EcM fungi.



Figure 2 Ranked frequency of ectomycorrhizal fungi on three host species in study I. Differential shading demonstrates the weighted proportion of each host species. Asterisks denote statistically significant host-biased frequency according to Fisher's Exact test and sharpened Benjamini-Hochberg corrections.

In Tumbama, 28 species were associated with the introduced *Eucalyptus* spp., and 46 species with native Caesalpinioideae. Nine species of EcM fungi had potential Australian origin and eight species had uncertain origin. Fourteen species were shared between the native and exotic host trees. Majority of those species were of African origin and only three species (/terfezia-peziza depressa Z01, /tomentella-thelephora Z01, /tomentellopsis Z01) were of Australian

origin. Potentially Australian species /hysterangium Z04 was found only once on native tree. Species accumulation curves for different host trees showed no difference of species diversity for native and exotic tree species neither on species nor family level (Figure 3).



Figure 3 Rarefied species accumulation curves (solid lines) and their 95% confidence intervals (dotted lines) of ectomycorrhizal fungi on root systems as based on (a) host species; and (b) host families.

Dead wood harbored 32 species at Warra and six species in Yarra. At Warra, species density, species richness in soil and in dead wood was significantly higher than in Yarra. EcM fungal taxa with resupinate fruit-bodies were found in dead wood on 39.4% occasions, whereas taxa with no or other types of fruit bodies occurred in dead wood on 24.4% occasions. The differences in these ratios were statistically significantly different based on chi-square tests. However, lineage level tests provided no support for preferential association of resupinate fruitbody type with dead wood. Nevertheless, there were statistically significant differences in abundance of in several fungal lineages between forest floor and dead wood. In particular, /cortinarius, /russula-lactarius and /sordariales were more abundant in forest floor soil. At species level, 14 of the 21 statistically compared data displayed significant differences for substrate. In particular, /cortinarius, /descolea, /russula-lactarius and /tomentella-thelephora comprised species that were more frequent in forest floor soil than in dead wood. Species occurring most commonly in logs (Tomentella sp12, Laccaria sp7, Cenococcum sp., Helotiales sp4) were among the most frequent EcM symbionts in forest floor soil.

DISCUSSION

In a Tasmanian wet sclerophyll forest (I), EcM fungi formed a phylogenetically diverse community that comprised 123 species. Minimum species richness estimators suggested the potential species richness to range from 210 to 247 species. However, the EcM fungal diversity at Yarra (II) was relative low in the forest floor soil in Nothofagus cunninghamii forest patch. In different ecosystems of Africa, fungal species richness was lower compared to Australia and Northern Hemisphere in spite of generally higher sampling effort (III: Tedersoo & Nara 2010). In all Australian sites (I, II), the most species-rich and abundant EcM fungal lineages were /cortinarius, /tomentella-thelephora, /russulalactarius, /clavulina, /descolea and /laccaria (I). All these lineages, except /descolea and /laccaria, are species-rich and abundant also in Holarctic fungal communities (Gardes & Dahlberg 1996, Horton & Bruns 1998, Tedersoo et al. 2006). The /descolea lineage is endemic to the Southern Hemisphere with an exception of a single species (Tedersoo et al. 2010a). By contrast, the /laccaria lineage is relatively species-poor (Rinaldi et al. 2008) and the species are usually relatively infrequent in mature forests of the Northern Hemisphere (Gherbi et al. 1999, Genney et al. 2006). These two lineages have not been found in African and South American tropical ecosystems (Tedersoo et al. 2010a). The most common and abundant EcM fungal lineages across all African sites included /russula-lactarius, /tomentella-thelephora and /boletus. Altogether 18 EcM fungal lineages were found from these study sites combined (III). In comparison, a single site in temperate ecosystems commonly supports more than 20 lineages (Tedersoo & Nara 2010), which indicates the relative phylogenetic poverty of African EcM communities. The community composition of EcM fungi is highly similar across different African sites, independent from ecosystem type, and resembles other tropical forests (Diédhiou et al. 2010, Peav et al. 2010). Only species of the /inocybe lineage tended to be more common in miombo woodlands of Zambia (III:IV) and Benin (N.S. Yorou & L. Tedersoo unpublished) compared to the rainforest habitats. The /russula-lactarius and /tomentella-thelephora are the dominant lineages both in Australian, African and northern temperate sites, indicating their global dominance. In particular, the /russula-lactarius lineage seems to comprise the greatest proportion of species in African ecosystems that is consistent with implications from fruitbody surveys (Buyck et al. 1996). Tropical ecosystems are inferred as potentially ancestral habitats and evolutionary centres for the /russula-lactarius lineage (Buyck et al. 2008). However, the generally lower species richness of EcM fungi in tropical regions is attributable to the poorer soil environment and evolutionary ecology of the major host plants (Tedersoo & Nara 2010).

In boreal and temperate ecosystems, host is usually one of the most important factors that affect EcM fungal species composition (Ishida et al. 2007, Morris et al. 2008). In Tasmania, 66% of the common EcM fungal species showed statistically significant host preference, although exclusive specificity was less common among the most frequent species (I). In this forest, host preference is clearly stronger than in Northern temperate ecosystems, where only few species have typically host-biased distribution (Horton & Bruns 1998, Ishida et al. 2007). Conversely, the dominant fungal species showed no host preference, independent from the plant groups and ecosystem types, in the African study sites (III), However, in NW Amazon, most of the dominant EcM fungal species displayed strong genus-level host specifity, which may be related to the inherent specificity in the *Nvctaginaceae* family (Tedersoo et al. 2010b). Alnus spp. in the Northern Hemisphere (Molina et al. 1992) and Gnetum spp. in the tropical regions (Bechem 2004) are also highly selective for their EcM symbionts, suggesting that the relative importance of host specificity is a function of the plant taxa in question rather than ecosystem types. Thus, (taxonomic) sampling effect may play a substantial role in addressing host preference and specificity patterns. Of fungi, most members of the /suillusrhizopogon lineage are exclusively associated with certain members of the *Pinaceae* family (Molina et al. 1992). Niche partitioning via host specificity has been hypothesized to contribute to the high fungal species richness in temperate ecosystems (Ishida et al. 2007), but experimental evidence for this is lacking. High host specificity but low fungal richness in NW Amazon (Tedersoo et al. 2010b), however, contradicts these predictions on a global scale.

Host shifts rather than strict coevolution appears to be a common pattern in EcM fungi in sympatric conditions (den Bakker et al. 2004, Matheny et al. 2009). There is also limited evidence for long-distance dispersal and simultaneous host shifts in Basidiomycota (Hosaka et al. 2008, Matheny et al. 2009). We demonstrated that broadening of host range may occur spontaneously in field conditions during the establishment of exotic forestry plantations and cointroduction of mycorrhizal symbionts (IV). As expected, shifts to alien hosts were more frequent when the root systems of different host trees were overlapping. We estimated that essentially all African EcM fungal species found in Tumbama are able to colonize Eucalyptus spp. in vivo, because no host preference was recovered among the native fungal species (IV). Previously, Garbaye et al. (1988) demonstrated that Eucalyptus spp. may establish EcM symbiosis with native fungi in experimental conditions, although a few fungal species failed to form EcM in other trials (Bâ et al. 2010). In contrast to African fungi, the three most common Australian fungal species did not colonize the native trees despite their prevalence on *Eucalyptus* spp. Nevertheless, some of the alien fungi were able to complete their life cycle and form sporulating fruitbodies. This indicates that these introduced species are adapting to the native conditions. Eucalyptus spp. perform better with their co-introduced fungi than with local fungi (Malaiczuk et al. 1984, Chen et al. 2007). Taken together, these results suggest that adaptive coevolution between fungi and their hosts is influenced by competition among fungi in the soil habitat.

Naturalization and invasion of EcM fungi may be more common with phylogenetically closely related hosts (Selosse et al. 1998, Pringle et al. 2009).

Many invasive species need time for naturalization before they become invasive (Richardson et al. 2000). Whether or not invasive EcM fungal species may influence their new habitat is not yet predictable. In novel habitats, EcM fungi may transform soil carbon cycling (Gadgil & Gadgil 1975, Chapela et al. 2001), affect mineral nutrient dynamics (Philips & Fahey 2006), and alter surrounding vegetation (Richardson & Rejmanek 2004).

Seedlings in dead wood hosted less EcM fungal species compared to mature trees and the species composition differed from soil EcM fungal composition (II). While Tedersoo et al. (2008) found out that in northern hemisphere EcM fungi with resupinate fruit-bodies are dominating in dead wood, such relationship was also found, but not regarded as causal in the current study (II). Rather, the dominant EcM fungi in soil also dominate on root systems of seedlings in dead wood (II), except members of the /cortinarius and /russula-lactarius lineages that are considered as late successional colonizers (Last et al. 1987, Gibson & Deacon 1990). The EcM lineages of /laccaria, /descolea and /tomentella-thelephora dominated on isolated seedlings in dead wood in Australia. Of these taxa, only members of the /tomentella-thelephora lineage form resupinate fruit-bodies. In contrast to Australia, *Laccaria* spp. rarely colonize seedlings on dead wood in European boreal forests (Tedersoo et al. 2008) and they are also infrequent members of northern temperate fungal communities (Tedersoo & Nara 2010).

CONCLUSIONS

Species richness was highest in Tasmanian wet sclerophyll forests and was comparable with species richness in Holarctic region (I). In contrast, in Africa, species richness was lower, and fewer fungal lineages were present (III). Tropical EcM fungal communities are phylogenetically highly similar in different continents. There was a strong host preference among EcM fungi in a Tasmanian wet sclerophyll forest. Compared to other hosts, Pomaderris apetala hosted highly different EcM fungal communities (I). However, host preference was not detected among the most common fungal species in Africa (III). Based on these and previously published results, host specificity is probably related to the (taxonomic) sampling effect of hosts. Host shifts among ectomycorrhizal fungi may occur particularly commonly, when the roots of trees intermingle (IV). These host shifts may lead to naturalization and invasion of the introduced Eucalyptus species or spread of cointroduced EcM fungal species as detected in other ecosystems. Seedlings of Nothofagus associate with EcM fungi during their establishment on dead wood in Australian wet temperate forest. General commonness explains the spread of EcM fungal species in dead wood better than resupinate fruiting habit.

REFERENCES

- Abarenkov K, Nilsson RH, Larsson K-H, Alexander IJ, Eberhardt U, Erland S, Høiland K, Kjøller R, Larsson E, Pennanen T, Sen R, Taylor AFS, Tedersoo L, Ursing B, Vrålstad T, Liimatainen K, Peintner U, Kõljalg U. 2010a. The UNITE database for molecular identification of fungi recent updates and future perspectives. *New Phytologist* 186: 281–285.
- Abarenkov K, Tedersoo L, Nilsson RH, Vellak K, Saar I, Veldre V, Parmasto E, Prous M, Aan A, Ots M, Kurina O, Ostonen I, Jõgeva J, Halapuu S, Põldmaa K, Toots M, Truu J, Larsson K-H, Kõljalg U. 2010b. PlutoF a web based workbench for ecological and taxonomic research with an online implementation for fungal ITS sequences. *Evolutionary Bioinformatics* 6: 198–196.
- Alexander IJ, Lee SS. 2005. Mycorrhizas and ecosystem processes in tropical rainforest: implications for diversity. In: Biotic Interactions in the Tropics: Their Role in the Maintenance of Species Diversity. Ed. Burslem DFRP, Pinard MA, Hartley SE. Cambridge University Press: London. Pp. 165–203.
- Anderson MJ, Willis TJ. 2003. Canonical analysis of principal coordinates: a useful method of constrained ordination for ecology. *Ecology* 84: 511–525.
- Bâ A, Diédhiou AG, Prin Y, Galiana A, Duponnois R. 2010. Management of ectomycorrhizal symbionts associated to useful exotic tree species to improve reforestation performances in tropical Africa. *Annals of Forest Science* 67: 301.
- Bahram M, Põlme S, Kõljalg U, Tedersoo L. 2011. A single European aspen (*Populus tremula*) tree individual may potentially harbour dozens of *Cenococcum geophilum* ITS genotypes and hundreds of species of ectomycorrhizal fungi. *FEMS Microbiology Ecology* 75: 313–320.
- Bastias BA, Xu Z, Cairney JWG. 2006. Influence of long-term repeated prescribed burning on mycelial communities of ectomycorrhizal fungi. *New Phytologist* 172: 149–158.
- Bechem EET. 2004. Mycorrhizal Status of *Gnetum* spp. in Cameroon: Evaluating Diversity with a View to Ameliorating Domestication Efforts. PhD Thesis, University of Aberdeen, UK.
- Borcard D, Legendre P. 2002. All-scale spatial analysis of ecological data by means of principal 9 coordinates of neighbour matrices. *Ecological Modelling* 153: 51–68.
- Bougher NL, Lebel T. 2001. Sequestrate (truffle-like) fungi of Australia and New Zealand. *Australian Systematic Botany* 14: 439–484.
- Bruns TD, Szaro TM, Gardes M, Cullings KW, Pan JJ, Taylor DL, Horton TR, Kretzer A, Garelotto M, Li Y. 1998. A sequence database for the identification of ectomycorrhizal basidiomycetes by phylogenetic analysis. *Molecular Ecology* 7: 257–272.
- Buyck B, Hofstetter V, Eberhardt U, Verbeken A, Kauff F. 2008. Walking the thin line between *Russula* and *Lactarius*: the dilemma of *Russula* subsect *Ochricompactae*. *Fungal Diversity* 28: 15–40.
- Buyck B, Thoen D, Watling R.1996. Ectomycorrhizal fungi of the Guinea-Congo Region. *Proceedings of the Royal Society of Edinburgh Section B: Biological Sciences* 104: 313–333.
- Buyck B. 2008. The edible mushrooms of Madagascar: an evolving enigma. *Economic Botany* 62: 509–520.

- Chapela IH, Osher LJ, Horton TR, Henn MR. 2001. Ectomycorrhizal fungi introduced with exotic pine plantations induce soil carbon depletion. *Soil Biology and Biochemistry* 33: 1733–1740.
- Chen YL, Liu S, Dell B. 2007. Mycorrhizal status of *Eucalyptus* plantations in South China and implications for management. *Mycorrhiza* 17: 527–535.
- Christy EJ, Mack RN. 1984. Variation in demography of juvenile *Tsuga heterophylla* across the substratum mosaic. *Journal of ecology* 72: 75–91.
- Claridge AW, Cork SJ, Trappe JM. 1999. Diversity and habitat relationships of hypogeous fungi. I. Study design, sampling techniques and general survey results. *Biodiversity and Conservation* 9: 151–173.
- Collier FA, Bidartondo MI. 2009. Waiting for fungi: the ectomycorrhizal invasion of lowland heathlands. *Journal of Ecology* 97: 950–963.
- Colwell RK. 2006. EstimateS: statistical estimation of species richness and shared species from samples. Version 8. Persistent URL <purl.oclc.org/estimates>.
- Connel JH, Lowman MD. 1989. Low diversity tropical rainforests: Some possible mechanisms for their existence. *The American Naturalist* 134: 88–119.
- den Bakker HC, Zuccarello GC, Kuyper YW, Noordeloos ME. 2004. Evolution and host specificity in the ectomycorrhizal genus *Leccinum*. *New Phytologist* 163: 201–215.
- Dickie IA, Bolstridge N, Cooper JA, Peltzer DA. 2010. Co-invasion by *Pinus* and its mycorrhizal fungi. *New Phytologist* 187: 475–484.
- Diédhiou A, Selosse M-A, Galiana A, Diabate M, Dreyfus B, Bâ A, de Faria S, Bena G. 2010. Multi-host ectomycorrhizal fungi are predominant in a Guinean tropical rainforest and shared between canopy trees and seedlings. *Environmental Microbiology* 12: 2219–2232.
- Diez J. 2005. Invasion biology of Australian ectomycorrhizal fungi introduced with eucalypt plantations into the Iberian Peninsula. *Biological Invasions* 7: 3–15.
- Dunk CV. 2002. The ectomycorrhizae of Nothofagus cunninghamii (Hook.) Oerst. in Noojee State Forest, Victoria, and evidence of host shifting by the introduced fungus Amanita muscaria (L.: Fr.) Pers. – identification, host determination, and characterisation of the new association. B. Sc. Thesis. La Trobe University. Bundoora, Victoria.
- FAO. 2010. Global forest resources assessment 2010. Main report. FAO Forestry Paper 163. Food and Agricultural Organisation of the United Nations, Rome.
- Gadgil RL, Gadgil PD. 1975. Suppression of litter decomposition by mycorrhizal roots of *Pinus radiata*. *New Zealand Journal of Science* 5: 33–41.
- Garbaye J, Delwaulle JC, Diangana D. 1988. Growth response of eucalypts in the Congo to ectomycorrhizal inoculation. *Forest Ecology and Management* 24: 151–157.
- Gardes M, Bruns TD. 1993. ITS primers with enhanced specificity for basidiomycetes application to the identification of mycorrhizas and rusts. *Molecular Ecology* 2: 113–118.
- Gardes M, Dahlberg A. 1996. Mycorrhizal diversity in arctic and alpine tundra: an open question. *New Phytologist* 133: 147–157.
- Gates GM, Ratkowsky DA, Grove SJ. 2005. A comparison of macrofungi in young silvivultural regeneration and mature forest at the Warra LTER site in the southern forests of Tasmania. *Tasforests* 16: 127–152.
- Genney DR, Anderson IC, Alexander IJ. 2006. Fine-scale distribution of pine ectomycorrhizas and their extramatrical mycelium. *New Phytologist* 170: 381–390.

- Gherbi H, Delaruelle C, Selosse M-A, Martin F. 1999. High genetic diversity in the population of the ectomycorrhizal basidiomycete *Laccaria amethystina* in a 150 year-old beech forest. *Molecular Ecology* 8: 2003–2013.
- Gibson F, Deacon JW. 1990. Establishment of ectomycorrhizas in aseptic culture: effects of glucose, nitrogen and phosphorus in relation to successions. *Mycological Research* 94: 166–172.
- Harvey AE, Jurgensen MF, Larsen MJ. 1978. Seasonal distribution of ectomycorrhizae in a mature Douglas fir / larch forest soil in Western Montana. *Forest Science* 24: 203–208.
- Hofgaard A. 1993. Structure and regeneration patterns in a virgin *Picea abies* forest in northern Sweden. *Journal of Vegetation Science* 4: 601–608.
- Hopple JS, Vilgalys R. 1994. Phylogenetic relationships among coprinoid taxa and allies based on data from restriction site mapping of nuclear rDNA. *Mycologia* 86: 96–107.
- Horton TR, Bruns TD. 1998. Multiple-host fungi are the most frequent and abundant ectomycorrhizal types in a mixed stand of Douglas fir (*Pseudotsuga menziesii*) and bishop pine (*Pinus muricata*). *New Phytologist* 139: 331–339.
- Hosaka K, Castellano MA, Spatafora JW. 2008. Biogeography of Hysterangiales (Phallomycetidae, Basidiomycota). *Mycological Research* 112: 448–462.
- Ishida TA, Nara K, Hogetsu T. 2007. Host effects on ectomycorrhizal fungal communities: insight from eight host species in mixed conifer-broadleaf forests. *New Phytologist* 174: 430–440.
- Iwanski M, Rudawska M. 2007. Ectomycorrhizal colonization of naturally regenerating *Pinus sylvestris* L. seedlings growing in different microhabitats in boreal forest. *Mycorrhiza* 17: 461–467.
- Katoh K, Kuma K, Toh H, Miyata T. 2005. MAFFT version 5: improvement in accuracy of multiple sequence alignment. *Nucleic Acids Research* 33: 511–518.
- Katoh K, Toh H. 2008. Recent developments in the MAFFT multiple sequence alignment program. *Briefings in Bioinformatics* 9: 286–298.
- Last FT, Dighton J, Mason PA. 1987. Successions of sheathing mycorrhizal fungi. *Trends in Ecology & Evolution* 2: 157–161.
- Lozupone CA, Hamady M, Knight R. 2006. UniFrac-An online tool for comparing microbial community diversity in a phylogenetic context. *BMC Bioinformatics* 7: 1–14.
- Lu X, Malajczuk N, Brundrett M, Dell B. 1999. Fruiting of putative ectomycorrhizal fungi under blue gum (*Eucalyptus globulus*) plantations of different ages in Western Australia. *Mycorrhiza* 8: 255–261.
- Malajczuk N, Molina R, Trappe JM. 1984. Ectomycorrhiza formation in *Eucalyptus*. II. The ultrastructure of compatible and incompatible mycorrhizal fungi and associated roots. *New Phytologist* 96: 43–53.
- Matheny PB, Aime MC, Bougher NL, Buyck B, Desjardin D, Horak E, Kropp BR, Lodge DJ, Soytong K, Trappe JM, Hibbett DS. 2009. Out of the palaeotropics? Historical biogeography and diversification in the cosmopolitan ectomycorrhizal mushroom family *Inocybaceae*. *Journal of Biogeography* 36: 577–592.
- McCune B, Mefford MJ. 2006. PC-ORD. Multivariate Analysis of Ecological Data. Version 5.04. MjM Software, Gleneden Beach, Oregon, USA.
- Midgley DJ, Saleeba JA, Stewart MI, Simpson AE, McGee PA. 2007. Molecular diversity of soil basidiomycete communities in northern-central New South Wales, Australia. *Mycological Research* 111: 370–378.

Mikola P. 1969. Afforestation of treeless areas. Unasylva 23: S1-S20.

- Molina R, Massicotte H, Trappe JM. 1992. Specificity phenomena in mycorrhizal symbiosis: community-ecological consequences and practical implications. In: Allen M (ed) Mycorrhizal functioning. An integrative plant-fungal process. New York, USA: Chapman and Hall. Pp. 357–423.
- Morris MH, Smith ME, Rizzo DM, Rejmanek M, Bledsoe CS. 2008. Contrasting ectomycorrhizal fungal communites on the roots of co-occuring oaks (*Quercus* spp.) in a California woodland. *New Phytologist* 178: 167–176.
- Nunez MA, Horton TR, Simberloff D. 2009. Lack of belowground mutualisms hinders *Pinaceae* invasions. *Ecology* 90: 2352–2359.
- Orlovich DA, Cairney JWG. 2004. Ectomycorrhizal fungi in New Zealand: current perspectives and future directions. *New Zealand Journal of Botany* 42: 721–738.
- Peay KG, Kennedy PG, Davies SJ, Tan S, Bruns TD. 2010. Potential link between plant and fungal distributions in a dipterocarp rainforest: community and phylogenetic structure of tropical ectomycorrhizal fungi across a plant and soil ecotone. *New Phytologist* 185: 529–542.
- Phillips RP, Fahey TJ. 2006. Tree species and mycorrhizal associations influence the magnitude of rhizosphere effects. *Ecology* 87: 1302–1313.
- Pringle A, Bever JD, Gardes M, Parrent JL, Rillig MC, Klironomos JN. 2009. Mycorrhizal symbioses and plant invasions. *Annual Reviews in Ecology, Evolution* and Systematics 40: 699–715.
- R Core Development Team (2007) R: A Language and Environment for Statistical Computing. R Foundation for Statistical Computing. Vienna, Austria.
- Ratkowsky DA, Gates GM. 2005. An inventory of macrofungi observed in Tasmanian forests over a six-year period. *Tasforests* 16: 153–168.
- Richardson DM, Allsopp N, D'Antonio C, Milton SJ, Rejmanek M. 2000. Plant invasions the role of mutualisms. *Biological Reviews* 75: 65–93.
- Richardson DM, Rejmanek M. 2004. Conifers as invasive aliens: a global survey and predictive framework. *Diversity and Distributions* 10: 321–331.
- Richardson DM. 1998. Forestry trees as invasive aliens. *Conservation Biology* 12: 18–26.
- Rinaldi AC, Comadini O, Kuyper TW. 2008. Ectomycorrhizal fungal diversity: separating the wheat from the chaff. *Fungal Diversity* 33: 1–45.
- Selosse M-A, Jacquot D, Bouchard D, Martin F, Le Tacon F. 1998. Temporal persistence and spatial distribution of an American inoculant strain of the ectomycorrhizal basidiomycete *Laccaria bicolor* in a French forest plantation. *Molecular Ecology* 7: 561–573.
- Smith SE, Read DJ, 1997. Mycorrhizal Symbiosis, 2nd edn. Academic Press, London.
- Stamatakis A, Hoover P, Rougemont J. 2008. A rapid bootstrap algorithm for the RAxML web-servers. *Systematic Biology* 75: 758–771.
- Swofford DL. 2002. PAUP*. Phylogenetic Analysis Using Parsimony (*and Other Methods). Version 4. Sinauer Associates, Sunderland, MA.
- Taylor AFS, Alexander IJ. 2005. The ectomycorrhizal symbiosis: life in the real world. *Mycologist* 19: 102–112.
- Taylor DL, McCormick MK. 2008. Internal transcribed spacer primers and sequences for improved characterization of basidomycetous orchid mycorrhizas. *New Phytologist* 177: 1020–1033.

- Tedersoo L, Kõljalg U, Hallenberg N, Larsson K-H. 2003. Fine scale distribution of ectomycorrhizal fungi and roots across substrate layers including coarse woody debris in a mixed forest. *New Phytologist* 159: 153–165.
- Tedersoo L, May TW, Smith ME. 2010a. Ectomycorrhizal lifestyle in fungi: global diversity, distribution, and evolution of phylogenetic lineages. *Mycorrhiza* 20: 217– 263.
- Tedersoo L, Nara K. 2010. General latitudinal gradient of biodiversity is reversed in ectomycorrhizal fungi. *New Phytologist* 185: 351–354.
- Tedersoo L, Sadam A, Zambrano M, Valencia R, Bahram M. 2010b. Low diversity and high host preference of ectomycorrhizal fungi in Western Amazonia, a neotropical biodiversity hotspot. *The ISME Journal* 4: 465–471.
- Tedersoo L, Suvi T, Beaver K, Kõljalg U. 2007. Ectomycorrhizal fungi of the Seychelles: diversity patterns and host shifts from the native *Vateriopsis* seychellarum (Dipterocarpaceae) and Intsia bijuga (Caesalpiniaceae) to the introduced Eucalyptus robusta (Myrtaceae), but not Pinus caribea (Pinaceae). New Phytologist 175: 321–333.
- Tedersoo L, Suvi T, Jairus T, Kõljalg U. 2008. Forest microsite effects on community composition of ectomycorrhizal fungi on seedlings of Picea abies and Betula pendula. *Environmental Microbiology* 10: 1189–1201.
- Tedersoo L, Suvi T, Jairus T, Ostonen I, Põlme S. 2009. Revisiting ectomycorrhizal fungi of Alnus: differential host specificity, diversity and determinants of the fungal community. *New Phytologist* 182: 727–735.
- Tedersoo L, Suvi T, Larsson E, Kõljalg U. 2006. Diversity and community structure of ectomycorrhizal fungi in a wooded meadow. *Mycological Research* 110: 734–748.
- Trappe JM. 1977. Selection of fungi for ectomycorrhizal inoculation in nurseries. *Annual Review of Phytophatology* 15: 203–222.
- Walbert K, Ramsfield TD, Ridgway HJ, Jones EE. 2010. Ectomycorrhizal species associated with *Pinus radiata* in New Zealand including novel association determined by molecular analysis. *Mycorrhiza* 20: 209–215.
- Wang B, Qiu Y-L. 2006. Phylogenetic distribution and evolution of mycorrhizas in land plants. *Mycorrhiza* 16: 299–363.
- Verhoeven KJF, Simonsen KL, McIntyre LM. 2005. Implementing false discovery rate control: increasing your power. *Oikos* 108: 643–647.
- West P. 2006. Growing plantation forests. Springer-Verlag, Berlin Heidelberg
- White TJ, Bruns TD, Lee S, Taylor J. 1990. Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics. In: Innis MA, Gelfand DH. (eds). PCR Protocols: A Guide to Methods and Applications. Academic Press: London. Pp. 315–322.
- Vilgalys R, Hester M. 1990. Rapid genetic identification and mapping of enzymatically amplified ribosomal DNA from several *Cryptococcus* species. *Journal of Bacteriology* 172: 4238–4246.
- Yazid SM, Lee SS, Lapeyrie F. 1994. Growth stimulation of *Hopea* spp. (*Dipterocarpaceae*) seedlings following ectomycorrhizal inoculation with an exotic strain of *Pisolithus tinctorius*. *Forest Ecology and Management* 67: 339–343.

SUMMARY IN ESTONIAN

Ektomükoriissete seente liigiline koosseis ja peremehe-eelistused Austraalia ja Aafrika ökosüsteemides

Mükoriisa on taimede ja seente vaheline vastastikku kasulik kooselu, mille käigus taim varustab seent suhkruga ning saab vastu mineraalaineid. Levinuimad mükoriisa tüübid on arbuskulaarne mükoriisa ja ektomükoriisa. Parasvöötme metsades on ülekaalus ektomükoriisat moodustavad puud, samas kui valdav enamus troopiliste alade puid moodustab arbuskulaarset mükoriisat. Ektomükoriissed puud moodustavad troopilistes ökosüsteemides monodominantseid kooslusi. Enamikul ektomükoriissetest puuliikidest ei ole kindlat seensümbiondi eelistust. Näiteks üks hariliku haava (*Populus tremula*) isend võib korraga moodustada mükoriisat rohkem kui saja erineva seeneliigiga. Samas lepp (*Alnus* spp.) kasvab koos vaid väheste temale omaste seentega. Lisaks peremeespuudele mõjutab ektomükoriisaseente liigilist koosseisu ka kasvukoha heterogeensus. Näiteks lagupuidu mükoriisaseente kooslus erineb mullas olevast ektomükoriisaseente kooslustest.

Seoses kasvava nõudlusega puidu järele suureneb kogu maailmas metsaistanduste pindala. Istandustes kasutatakse peamiselt kiirekasvulisi sissetoodud puuliike, millel on uues keskkonnas vähem kahjureid. Enamik istandustes kasvavaid puid on ektomükoriissed ja mitmed neist sõltuvad seensümbiontide olemasolust. Minu töö eesmärgid olid: 1) kirjeldada ektomükoriisaseente kooslusi Austraalia niisket tüüpi sklerofüllimetsas ja Aafrika valitud ökosüsteemides; 2) tuvastada ektomükoriissete seente peremehe-eelistused nendes kooslustes; 3) tuvastada, kas kohalikud ja sissetoodud seened on võimelised oma looduslikult peremeestaimelt võõrale peremeestaimele üle kanduma ja 4) võrrelda ektomükoriisaseente kooslusi lagupuidus ja mullas Austraalia lõunapöögi metsades.

Uurimustöö viidi läbi ühes Tasmaania niisket tüüpi sklerofüllimetsas, ühes Austraalia ja ühes Tasmaania lõunapöögi metsas, neljas erinevas Aafrika looduslikus ökosüsteemis ning kahes eukalüpti-istanduses Kesk-Aafrika lõunaosas, kus teise rinde moodustasid kohalikud puud. Igalt proovialalt koguti proovid sama metoodikat kasutades. Nii seene- kui ka taimeliigid määrati molekulaarsete meetodite abil. Liikide geograafilise päritolu kindlaks tegemisel kasutati võrdluseks teistest Aafrika looduslikest kasvukohtadest kogutud materjale.

Selgus, et Austraalias on ektomükoriisaseente liigiline ja fülogeneetiline mitmekesisus suurem kui Aafrikas. Kui Aafrika erinevates ökosüsteemides olid arvukamalt esindatud seened rühmadest /russula-lactarius, /tomentella-thelephora ja /boletus, siis Austraalias oli arvukaim /cortinarius, millele järgnesid /tomentella-thelephora, /russula-lactarius, /clavulina, /descolea ja /laccaria. Austraalias oli 66% ektomükoriissetest seeneliikidest peremehe spetsiifilised, samas kui Aafrika seentel peremehe-eelistust ei esinenud. Aafrikast ja

Austraaliast pärit seened olid võimelised eukalüptiistanduses mükoriisat moodustama mitteomaste peremeestaimedega.

Liigirikkus oli suurim Tasmaania niisket tüüpi sklerofüllimetsas, mis on võrreldav parasvöötme metsadega (I). Aafrikas oli nii ektomükoriissete seente liigirikkus kui ka olemasolevate seenerühmade arv väiksem (III) ning sarnanes teiste troopikas tehtud uurimuste tulemustele. Tasmaania niisket tüüpi sklerofüllimetsas eelistasid seeneliigid teatud peremeestaimi. Liigi *Pomaderris apetala* seenekooslus erines selgelt teiste puude omast (I). Aafrikas ei leitud ühelgi domineerival seeneliigil peremehe-eelistusi (III). Peremehe vahetused on ektomükoriissetel seentel suhteliselt tavalised, kui kohalike ja sissetoodud puude juured puutuvad kokku (IV). Introdutseeritud seente peremehe vahetused võivad põhjustada sissetoodud puude ja seente naturaliseerumist või bioloogilist invasiooni (IV). Lagupuidul kasvavad lõunapöögi seemikud olid sümbioosis ektomükoriisaseentega (II) ning nende seente liigilist koosseisu mõjutas antud seeneliikide ohtrus ümbritsevas mullas, mitte seente viljakeha tüüp.

ACKNOWLEDGEMENTS

First I thank my supervisor Leho for all the help, advice and guidance. I am grateful for the people who taught me fungi – Kuulo Kalamees, Anu Kollom, Urmas Kõljalg, Vello Liiv, Kadri Põldmaa, Kadri Pärtel, Irja Saar and the late Mall Vaasma who brought me to the world of mushroom trips and exhibitions.

I thank my roommates Heidi, Jane and Sergei and will always remember our "scientific" discussions. I also thank Triin and Kessy, who made our lunches and teatimes memorable. I am thankful to all the other mycologists from Tartu University, Estonian University of Life Sciences and from Tallinn. I also thank all the co-authors and the people who helped us during fieldwork.

I thank my family for all the support and understanding. I am always thankful that you let me choose my own path. I thank all the friends who have been there for me. Many thanks also to people who commented on the manuscript of this thesis.

PUBLICATIONS

CURRICULUM VITAE

I. General

Name: Teele Jairus

Date and place of birth: 31.10.1984, Hanila

Citizenship and nationality: Estonian

Address, phone, e-mail: Institute of Ecology and Earth Sciences, University of Tartu. 14A Ravila Street 50411 Tartu, Estonia. +3727376672, teele.jairus@ut.ee

Current position: Department of Botany, Institute of Ecology and Earth Sciences, University of Tartu, PhD student

Education:

2007 University of Tartu, Master of Science in Natural Sciences 2005 University of Tartu, diploma in botany

2002 Gymnasium of Lihula

1999 Elementary School of Vatla

Language skills: Estonian, English, German

Institution and position held:

2006–2008 University of Tartu, Museums of Tartu University; specialist (0.40) 2006–2006 University of Tartu, Faculty of Science and Technology, Institute of Ecology and Earth Sciences, Tartu University, Chair of Mycology; laboratory technician

II. Scientific and research activity

Research interests

Ecology and biogeography of ectomycorrhizal fungi Host preference of ectomycorrhizal fungi Host shifts among native and introduced ectomycorrhizal fungi

Publications:

- Tedersoo L, Bahram M, Jairus T, Bechem E, Chinoya S, Mpumba R, Leal M, Randrianjohany E, Razafimandimbison S, Sadam A, Naadel T, Kõljalg U. 2011. Spatial structure and the effects of host and soil environments on communities of ectomycorrhizal fungi in wooded savannas and rain forests of Continental Africa and Madagascar. Molecular Ecology doi: 10.1111/j.1365-294X.2011.05145.x [in print]
- Jairus T, Mpumba R, Chinoya S, Tedersoo L. 2011. Invasion potential and host shifts of Australian and African ectomycorrhizal fungi in mixed eucalypt plantations. New Phytologist doi: 10.1111/j.1469-8137.2011.03775.x [in print]
- Tedersoo L, Nilsson RH, Abarenkov K, Jairus T, Sadam A, Saar I, Bahram M, Bechem E, Chuyong G, Kõljalg U. 2010. 454 Pyrosequencing and Sanger sequencing of tropical mycorrhizal fungi provide similar results but reveal substantial methodological biases. New Phytologist 188: 291–301.

- Tedersoo L, Pärtel K, Jairus T, Gates G, Põldmaa K, Tamm H. 2009. Ascomycetes associated with ectomycorrhizas: molecular diversity and ecology with particular reference to the Helotiales. Environmental Microbiology 3166– 3179.
- Tedersoo L, Suvi T, Jairus T, Ostonen I, Põlme S. 2009. Revisiting ectomycorrhizal fungi of Alnus: differential host specificity, diversity and determinants of the fungal community. New Phytologist 182: 727–735.
- Tedersoo L, Gates G, Dunk C, Lebel T, May TW, Kõljalg U, Jairus T. 2009. Establishment of ectomycorrhizal fungal community on isolated Nothofagus cunninghamii seedlings regenerating on dead wood in Australian wet temperate forests: does fruit-body type matter. Mycorrhiza 19: 403–416.
- Tedersoo L, Jairus T, Horton BM, Abarenkov A, Suvi T, Saar I, Kõljalg U. 2008. Strong host preference of ectomycorrhizal fungi in a Tasmanian wet sclerophyll forest as revealed by DNA barcoding and taxon-specific primers. New Phytologist 180: 479–490.
- Tedersoo L, Suvi T, Jairus T, Kõljalg U. 2008. Forest microsite effects on community composition of ectomycorrhizal fungi on seedlings of Picea abies and Betula pendula. Environmental Microbiology 10: 1189–1201.

Conference presentations:

Jairus T, Chinoya S, Mpumba R, Tedersoo L. Host shifts between Australian and African ectomycorrhizal fungi. In conference "Next generation insights into geosciences and ecology", may 2011, Tartu, Estonia.

Awards and scholarships

Doctoral school of Ecology and Earth Sciences travelling grant, 2011 DoRa travelling grant, 2008 Kaleva Travel's travelling grant, 2007 FEMS Research Fellowship, 2007 Doctoral School of Ecology and Environmental Sciences PhD student scholarship, 2007

Other activities and membership:

Since 2007 member of Estonian Microbiologist society

III. Improvement of skills

Courses attended:

- PhD course: Identifying ectomycorrhizal fungi from environmental samples to DNA sequences, 12–24 August, 2007
- PhD course: Analyzing high throughput sequence data in microbial community ecology, 23–26 March, 2010

PhD course: Dynamics of Organic Matter in Soil, 22-28 May, 2011

CURRICULUM VITAE

I. Üldandmed

Ees- ja perekonnanimi: Teele Jairus

Sünniaeg ja koht: 31.10.1984, Hanila vald

Kodakondsus: Eesti

Aadress, telefon, e-post: Tartu Ülikool, Ökoloogia ja Maateaduste Instituut. Ravila 14 A, 50411. +3727376672, teele.jairus@ut.ee

Praegune töökoht, amet: Botaanika osakond, Ökoloogia ja Maateaduste Instituut, Tartu Ülikool, doktorant

Haridus:

2007 Tartu Ülikool Loodusteaduse magister (MSc), taime- ja seeneteadus 2005 Tartu Ülikool Loodusteaduse bakalaureus, Bioloogia 2002 Lihula Gümnaasium 1999 Vatla Põhikool

Keelteoskus: eesti, inglise, saksa

Töökogemus:

2006–2007 Tartu Ülikool, Tartu Ülikooli Muuseumid; spetsialist (0.40) 2006–2006 Tartu Ülikool, Loodus- ja tehnoloogiateaduskond, Tartu Ülikooli Ökoloogia- ja Maateaduste Instituut, Mükoloogia õppetool; laborant

II. Teaduslik ja arendustegevus

Peamised uurimisvaldkonnad:

Aafrika ektomükoriisat moodustavate seente ökoloogia ja biogeograafia Ektomükoriissete seente peremehe-eelistused

Peremehevahetused kohalikel ja sisstoodud ektomükoriisa seentel

Publikatsioonid:

- Tedersoo L, Bahram M, Jairus T, Bechem E, Chinoya S, Mpumba R, Leal M, Randrianjohany E, Razafimandimbison S, Sadam A, Naadel T, Kõljalg U. 2011. Spatial structure and the effects of host and soil environments on communities of ectomycorrhizal fungi in wooded savannas and rain forests of Continental Africa and Madagascar. Molecular Ecology doi: 10.1111/j.1365-294X.2011.05145.x [ilmumas]
- Jairus T, Mpumba R, Chinoya S, Tedersoo L. 2011. Invasion potential and host shifts of Australian and African ectomycorrhizal fungi in mixed eucalypt plantations. New Phytologist doi: 10.1111/j.1469-8137.2011.03775.x [ilmumas]
- Tedersoo L, Nilsson RH, Abarenkov K, Jairus T, Sadam A, Saar I, Bahram M, Bechem E, Chuyong G, Kõljalg U. 2010. 454 Pyrosequencing and Sanger sequencing of tropical mycorrhizal fungi provide similar results but reveal substantial methodological biases. New Phytologist 188: 291–301.

- Tedersoo L, Pärtel K, Jairus T, Gates G, Põldmaa K, Tamm H. 2009. Ascomycetes associated with ectomycorrhizas: molecular diversity and ecology with particular reference to the Helotiales. Environmental Microbiology 3166–3179.
- Tedersoo L, Suvi T, Jairus T, Ostonen I, Põlme S. 2009. Revisiting ectomycorrhizal fungi of Alnus: differential host specificity, diversity and determinants of the fungal community. New Phytologist 182: 727–735.
- Tedersoo L, Gates G, Dunk C, Lebel T, May TW, Kõljalg U, Jairus T. 2009. Establishment of ectomycorrhizal fungal community on isolated Nothofagus cunninghamii seedlings regenerating on dead wood in Australian wet temperate forests: does fruit-body type matter. Mycorrhiza 19: 403–416.
- Tedersoo L, Jairus T, Horton BM, Abarenkov A, Suvi T, Saar I, Kõljalg U. 2008. Strong host preference of ectomycorrhizal fungi in a Tasmanian wet sclerophyll forest as revealed by DNA barcoding and taxon-specific primers. New Phytologist 180: 479–490.
- Tedersoo L, Suvi T, Jairus T, Kõljalg U. 2008. Forest microsite effects on community composition of ectomycorrhizal fungi on seedlings of Picea abies and Betula pendula. Environmental Microbiology 10: 1189–1201.

Konverentsiettekanded:

Jairus T, Chinoya S, Mpumba R, Tedersoo L. Host shifts between Australian and African ectomycorrhizal fungi. Maateaduste ja ökoloogia erialade doktorantide konverents "Next generation insights into geosciences and ecology", mai 2011, Tartu, Eesti

Saadud uurimistoetused ja stipendiumid

Maateaduste ja Ökoloogia doktorikooli välissõidutoetus 2011 DoRa doktorantide semester välismaal 2008 Kaleva Traveli reisistipendium 2007 FEMS Research Fellowship 2007 Ökoloogia ja keskkonnateaduste doktorikool, toetus doktorandi teadustöö finantseerimiseks 2007

Muu teaduslik organisatsiooniline ja erialane tegevus

2007 – Eesti Mikrobioloogide Ühenduse liige

III. Erialane enesetäiendus

- Kursus: Identifying ectomycorrhizal fungi from environmental samples to DNA sequences, 12–24 august 2007
- Kursus: Analyzing high throughput sequence data in microbial community ecology, 23–26 märts 2010

Kursus: Dynamics of Organic Matter in Soil, 22–28 mai 2011

DISSERTATIONES BIOLOGICAE UNIVERSITATIS TARTUENSIS

- 1. Toivo Maimets. Studies of human oncoprotein p53. Tartu, 1991, 96 p.
- 2. Enn K. Seppet. Thyroid state control over energy metabolism, ion transport and contractile functions in rat heart. Tartu, 1991, 135 p.
- 3. Kristjan Zobel. Epifüütsete makrosamblike väärtus õhu saastuse indikaatoritena Hamar-Dobani boreaalsetes mägimetsades. Tartu, 1992, 131 lk.
- 4. Andres Mäe. Conjugal mobilization of catabolic plasmids by transposable elements in helper plasmids. Tartu, 1992, 91 p.
- 5. Maia Kivisaar. Studies on phenol degradation genes of *Pseudomonas* sp. strain EST 1001. Tartu, 1992, 61 p.
- 6. Allan Nurk. Nucleotide sequences of phenol degradative genes from *Pseudomonas sp.* strain EST 1001 and their transcriptional activation in *Pseudomonas putida*. Tartu, 1992, 72 p.
- 7. Ülo Tamm. The genus *Populus* L. in Estonia: variation of the species biology and introduction. Tartu, 1993, 91 p.
- 8. Jaanus Remme. Studies on the peptidyltransferase centre of the *E.coli* ribosome. Tartu, 1993, 68 p.
- 9. Ülo Langel. Galanin and galanin antagonists. Tartu, 1993, 97 p.
- 10. Arvo Käärd. The development of an automatic online dynamic fluorescense-based pH-dependent fiber optic penicillin flowthrought biosensor for the control of the benzylpenicillin hydrolysis. Tartu, 1993, 117 p.
- 11. Lilian Järvekülg. Antigenic analysis and development of sensitive immunoassay for potato viruses. Tartu, 1993, 147 p.
- 12. Jaak Palumets. Analysis of phytomass partition in Norway spruce. Tartu, 1993, 47 p.
- 13. Arne Sellin. Variation in hydraulic architecture of *Picea abies* (L.) Karst. trees grown under different environmental conditions. Tartu, 1994, 119 p.
- 13. Mati Reeben. Regulation of light neurofilament gene expression. Tartu, 1994, 108 p.
- 14. Urmas Tartes. Respiration rhytms in insects. Tartu, 1995, 109 p.
- 15. **Ülo Puurand.** The complete nucleotide sequence and infections *in vitro* transcripts from cloned cDNA of a potato A potyvirus. Tartu, 1995, 96 p.
- 16. **Peeter Hõrak**. Pathways of selection in avian reproduction: a functional framework and its application in the population study of the great tit (*Parus major*). Tartu, 1995, 118 p.
- 17. Erkki Truve. Studies on specific and broad spectrum virus resistance in transgenic plants. Tartu, 1996, 158 p.
- 18. Illar Pata. Cloning and characterization of human and mouse ribosomal protein S6-encoding genes. Tartu, 1996, 60 p.
- Ülo Niinemets. Importance of structural features of leaves and canopy in determining species shade-tolerance in temperature deciduous woody taxa. Tartu, 1996, 150 p.

- 20. Ants Kurg. Bovine leukemia virus: molecular studies on the packaging region and DNA diagnostics in cattle. Tartu, 1996, 104 p.
- 21. Ene Ustav. E2 as the modulator of the BPV1 DNA replication. Tartu, 1996, 100 p.
- 22. Aksel Soosaar. Role of helix-loop-helix and nuclear hormone receptor transcription factors in neurogenesis. Tartu, 1996, 109 p.
- 23. Maido Remm. Human papillomavirus type 18: replication, transformation and gene expression. Tartu, 1997, 117 p.
- 24. **Tiiu Kull**. Population dynamics in *Cypripedium calceolus* L. Tartu, 1997, 124 p.
- 25. Kalle Olli. Evolutionary life-strategies of autotrophic planktonic microorganisms in the Baltic Sea. Tartu, 1997, 180 p.
- 26. **Meelis Pärtel**. Species diversity and community dynamics in calcareous grassland communities in Western Estonia. Tartu, 1997, 124 p.
- 27. Malle Leht. The Genus *Potentilla* L. in Estonia, Latvia and Lithuania: distribution, morphology and taxonomy. Tartu, 1997, 186 p.
- 28. **Tanel Tenson**. Ribosomes, peptides and antibiotic resistance. Tartu, 1997, 80 p.
- 29. Arvo Tuvikene. Assessment of inland water pollution using biomarker responses in fish *in vivo* and *in vitro*. Tartu, 1997, 160 p.
- 30. Urmas Saarma. Tuning ribosomal elongation cycle by mutagenesis of 23S rRNA. Tartu, 1997, 134 p.
- 31. **Henn Ojaveer**. Composition and dynamics of fish stocks in the gulf of Riga ecosystem. Tartu, 1997, 138 p.
- 32. Lembi Lõugas. Post-glacial development of vertebrate fauna in Estonian water bodies. Tartu, 1997, 138 p.
- 33. Margus Pooga. Cell penetrating peptide, transportan, and its predecessors, galanin-based chimeric peptides. Tartu, 1998, 110 p.
- 34. Andres Saag. Evolutionary relationships in some cetrarioid genera (Lichenized Ascomycota). Tartu, 1998, 196 p.
- 35. Aivar Liiv. Ribosomal large subunit assembly in vivo. Tartu, 1998, 158 p.
- 36. **Tatjana Oja**. Isoenzyme diversity and phylogenetic affinities among the eurasian annual bromes (*Bromus* L., Poaceae). Tartu, 1998, 92 p.
- 37. **Mari Moora**. The influence of arbuscular mycorrhizal (AM) symbiosis on the competition and coexistence of calcareous crassland plant species. Tartu, 1998, 78 p.
- 38. **Olavi Kurina**. Fungus gnats in Estonia (*Diptera: Bolitophilidae, Keroplatidae, Macroceridae, Ditomyiidae, Diadocidiidae, Mycetophilidae*). Tartu, 1998, 200 p.
- 39. Andrus Tasa. Biological leaching of shales: black shale and oil shale. Tartu, 1998, 98 p.
- 40. Arnold Kristjuhan. Studies on transcriptional activator properties of tumor suppressor protein p53. Tartu, 1998, 86 p.

- 41. **Sulev Ingerpuu.** Characterization of some human myeloid cell surface and nuclear differentiation antigens. Tartu, 1998, 163 p.
- 42. Veljo Kisand. Responses of planktonic bacteria to the abiotic and biotic factors in the shallow lake Võrtsjärv. Tartu, 1998, 118 p.
- 43. Kadri Põldmaa. Studies in the systematics of hypomyces and allied genera (Hypocreales, Ascomycota). Tartu, 1998, 178 p.
- 44. Markus Vetemaa. Reproduction parameters of fish as indicators in environmental monitoring. Tartu, 1998, 117 p.
- 45. **Heli Talvik.** Prepatent periods and species composition of different *Oesophagostomum* spp. populations in Estonia and Denmark. Tartu, 1998, 104 p.
- 46. Katrin Heinsoo. Cuticular and stomatal antechamber conductance to water vapour diffusion in *Picea abies* (L.) karst. Tartu, 1999, 133 p.
- 47. **Tarmo Annilo.** Studies on mammalian ribosomal protein S7. Tartu, 1998, 77 p.
- 48. **Indrek Ots.** Health state indicies of reproducing great tits (*Parus major*): sources of variation and connections with life-history traits. Tartu, 1999, 117 p.
- 49. Juan Jose Cantero. Plant community diversity and habitat relationships in central Argentina grasslands. Tartu, 1999, 161 p.
- 50. **Rein Kalamees.** Seed bank, seed rain and community regeneration in Estonian calcareous grasslands. Tartu, 1999, 107 p.
- 51. **Sulev Kõks.** Cholecystokinin (CCK) induced anxiety in rats: influence of environmental stimuli and involvement of endopioid mechanisms and erotonin. Tartu, 1999, 123 p.
- 52. Ebe Sild. Impact of increasing concentrations of O₃ and CO₂ on wheat, clover and pasture. Tartu, 1999, 123 p.
- 53. Ljudmilla Timofejeva. Electron microscopical analysis of the synaptonemal complex formation in cereals. Tartu, 1999, 99 p.
- 54. Andres Valkna. Interactions of galanin receptor with ligands and G-proteins: studies with synthetic peptides. Tartu, 1999, 103 p.
- 55. **Taavi Virro.** Life cycles of planktonic rotifers in lake Peipsi. Tartu, 1999, 101 p.
- 56. Ana Rebane. Mammalian ribosomal protein S3a genes and intron-encoded small nucleolar RNAs U73 and U82. Tartu, 1999, 85 p.
- 57. **Tiina Tamm.** Cocksfoot mottle virus: the genome organisation and translational strategies. Tartu, 2000, 101 p.
- 58. **Reet Kurg.** Structure-function relationship of the bovine papilloma virus E2 protein. Tartu, 2000, 89 p.
- 59. **Toomas Kivisild.** The origins of Southern and Western Eurasian populations: an mtDNA study. Tartu, 2000, 121 p.
- 60. **Niilo Kaldalu.** Studies of the TOL plasmid transcription factor XylS. Tartu 2000. 88 p.

- 61. **Dina Lepik.** Modulation of viral DNA replication by tumor suppressor protein p53. Tartu 2000. 106 p.
- 62. Kai Vellak. Influence of different factors on the diversity of the bryophyte vegetation in forest and wooded meadow communities. Tartu 2000. 122 p.
- 63. Jonne Kotta. Impact of eutrophication and biological invasionas on the structure and functions of benthic macrofauna. Tartu 2000. 160 p.
- 64. **Georg Martin.** Phytobenthic communities of the Gulf of Riga and the inner sea the West-Estonian archipelago. Tartu, 2000. 139 p.
- 65. Silvia Sepp. Morphological and genetical variation of *Alchemilla L*. in Estonia. Tartu, 2000. 124 p.
- 66. Jaan Liira. On the determinants of structure and diversity in herbaceous plant communities. Tartu, 2000. 96 p.
- 67. **Priit Zingel.** The role of planktonic ciliates in lake ecosystems. Tartu 2001. 111 p.
- 68. **Tiit Teder.** Direct and indirect effects in Host-parasitoid interactions: ecological and evolutionary consequences. Tartu 2001. 122 p.
- 69. **Hannes Kollist.** Leaf apoplastic ascorbate as ozone scavenger and its transport across the plasma membrane. Tartu 2001. 80 p.
- 70. **Reet Marits.** Role of two-component regulator system PehR-PehS and extracellular protease PrtW in virulence of *Erwinia Carotovora* subsp. *Carotovora*. Tartu 2001. 112 p.
- 71. **Vallo Tilgar.** Effect of calcium supplementation on reproductive performance of the pied flycatcher *Ficedula hypoleuca* and the great tit *Parus major*, breeding in Nothern temperate forests. Tartu, 2002. 126 p.
- 72. **Rita Hõrak.** Regulation of transposition of transposon Tn4652 in *Pseudomonas putida*. Tartu, 2002. 108 p.
- 73. Liina Eek-Piirsoo. The effect of fertilization, mowing and additional illumination on the structure of a species-rich grassland community. Tartu, 2002. 74 p.
- 74. **Krõõt Aasamaa.** Shoot hydraulic conductance and stomatal conductance of six temperate deciduous tree species. Tartu, 2002. 110 p.
- 75. **Nele Ingerpuu.** Bryophyte diversity and vascular plants. Tartu, 2002. 112 p.
- 76. Neeme Tõnisson. Mutation detection by primer extension on oligonucleotide microarrays. Tartu, 2002. 124 p.
- 77. **Margus Pensa.** Variation in needle retention of Scots pine in relation to leaf morphology, nitrogen conservation and tree age. Tartu, 2003. 110 p.
- 78. **Asko Lõhmus.** Habitat preferences and quality for birds of prey: from principles to applications. Tartu, 2003. 168 p.
- 79. Viljar Jaks. p53 a switch in cellular circuit. Tartu, 2003. 160 p.
- 80. Jaana Männik. Characterization and genetic studies of four ATP-binding cassette (ABC) transporters. Tartu, 2003. 140 p.
- 81. Marek Sammul. Competition and coexistence of clonal plants in relation to productivity. Tartu, 2003. 159 p

- 82. **Ivar Ilves.** Virus-cell interactions in the replication cycle of bovine papillomavirus type 1. Tartu, 2003. 89 p.
- 83. Andres Männik. Design and characterization of a novel vector system based on the stable replicator of bovine papillomavirus type 1. Tartu, 2003. 109 p.
- 84. **Ivika Ostonen.** Fine root structure, dynamics and proportion in net primary production of Norway spruce forest ecosystem in relation to site conditions. Tartu, 2003. 158 p.
- 85. **Gudrun Veldre.** Somatic status of 12–15-year-old Tartu schoolchildren. Tartu, 2003. 199 p.
- 86. Ülo Väli. The greater spotted eagle *Aquila clanga* and the lesser spotted eagle *A. pomarina*: taxonomy, phylogeography and ecology. Tartu, 2004. 159 p.
- 87. **Aare Abroi.** The determinants for the native activities of the bovine papillomavirus type 1 E2 protein are separable. Tartu, 2004. 135 p.
- 88. Tiina Kahre. Cystic fibrosis in Estonia. Tartu, 2004. 116 p.
- 89. **Helen Orav-Kotta.** Habitat choice and feeding activity of benthic suspension feeders and mesograzers in the northern Baltic Sea. Tartu, 2004. 117 p.
- 90. **Maarja Öpik.** Diversity of arbuscular mycorrhizal fungi in the roots of perennial plants and their effect on plant performance. Tartu, 2004. 175 p.
- 91. Kadri Tali. Species structure of *Neotinea ustulata*. Tartu, 2004. 109 p.
- 92. Kristiina Tambets. Towards the understanding of post-glacial spread of human mitochondrial DNA haplogroups in Europe and beyond: a phylogeographic approach. Tartu, 2004. 163 p.
- 93. Arvi Jõers. Regulation of p53-dependent transcription. Tartu, 2004. 103 p.
- 94. Lilian Kadaja. Studies on modulation of the activity of tumor suppressor protein p53. Tartu, 2004. 103 p.
- 95. Jaak Truu. Oil shale industry wastewater: impact on river microbial community and possibilities for bioremediation. Tartu, 2004. 128 p.
- 96. **Maire Peters.** Natural horizontal transfer of the *pheBA* operon. Tartu, 2004. 105 p.
- 97. Ülo Maiväli. Studies on the structure-function relationship of the bacterial ribosome. Tartu, 2004. 130 p.
- 98. Merit Otsus. Plant community regeneration and species diversity in dry calcareous grasslands. Tartu, 2004. 103 p.
- 99. Mikk Heidemaa. Systematic studies on sawflies of the genera *Dolerus*, *Empria*, and *Caliroa* (Hymenoptera: Tenthredinidae). Tartu, 2004. 167 p.
- 100. **Ilmar Tõnno.** The impact of nitrogen and phosphorus concentration and N/P ratio on cyanobacterial dominance and N_2 fixation in some Estonian lakes. Tartu, 2004. 111 p.
- 101. Lauri Saks. Immune function, parasites, and carotenoid-based ornaments in greenfinches. Tartu, 2004. 144 p.

- 102. **Siiri Rootsi.** Human Y-chromosomal variation in European populations. Tartu, 2004. 142 p.
- 103. Eve Vedler. Structure of the 2,4-dichloro-phenoxyacetic acid-degradative plasmid pEST4011. Tartu, 2005. 106 p.
- 104. Andres Tover. Regulation of transcription of the phenol degradation *pheBA* operon in *Pseudomonas putida*. Tartu, 2005. 126 p.
- 105. Helen Udras. Hexose kinases and glucose transport in the yeast *Hansenula polymorpha*. Tartu, 2005. 100 p.
- 106. Ave Suija. Lichens and lichenicolous fungi in Estonia: diversity, distribution patterns, taxonomy. Tartu, 2005. 162 p.
- 107. **Piret Lõhmus.** Forest lichens and their substrata in Estonia. Tartu, 2005. 162 p.
- 108. **Inga Lips.** Abiotic factors controlling the cyanobacterial bloom occurrence in the Gulf of Finland. Tartu, 2005. 156 p.
- 109. Kaasik, Krista. Circadian clock genes in mammalian clockwork, metabolism and behaviour. Tartu, 2005. 121 p.
- 110. Juhan Javoiš. The effects of experience on host acceptance in ovipositing moths. Tartu, 2005. 112 p.
- 111. **Tiina Sedman.** Characterization of the yeast *Saccharomyces cerevisiae* mitochondrial DNA helicase Hmi1. Tartu, 2005. 103 p.
- 112. **Ruth Aguraiuja.** Hawaiian endemic fern lineage *Diellia* (Aspleniaceae): distribution, population structure and ecology. Tartu, 2005. 112 p.
- 113. **Riho Teras.** Regulation of transcription from the fusion promoters generated by transposition of Tn4652 into the upstream region of *pheBA* operon in *Pseudomonas putida*. Tartu, 2005. 106 p.
- 114. **Mait Metspalu.** Through the course of prehistory in india: tracing the mtDNA trail. Tartu, 2005. 138 p.
- 115. Elin Lõhmussaar. The comparative patterns of linkage disequilibrium in European populations and its implication for genetic association studies. Tartu, 2006. 124 p.
- 116. **Priit Kupper.** Hydraulic and environmental limitations to leaf water relations in trees with respect to canopy position. Tartu, 2006. 126 p.
- 117. **Heili Ilves.** Stress-induced transposition of Tn4652 in *Pseudomonas Putida*. Tartu, 2006. 120 p.
- 118. **Silja Kuusk.** Biochemical properties of Hmi1p, a DNA helicase from *Saccharomyces cerevisiae* mitochondria. Tartu, 2006. 126 p.
- 119. Kersti Püssa. Forest edges on medium resolution landsat thematic mapper satellite images. Tartu, 2006. 90 p.
- 120. Lea Tummeleht. Physiological condition and immune function in great tits (*Parus major* 1.): Sources of variation and trade-offs in relation to growth. Tartu, 2006. 94 p.
- 121. **Toomas Esperk.** Larval instar as a key element of insect growth schedules. Tartu, 2006. 186 p.

- 122. Harri Valdmann. Lynx (*Lynx lynx*) and wolf (*Canis lupus*) in the Baltic region: Diets, helminth parasites and genetic variation. Tartu, 2006. 102 p.
- 123. **Priit Jõers.** Studies of the mitochondrial helicase Hmi1p in *Candida albicans* and *Saccharomyces cerevisia*. Tartu, 2006. 113 p.
- 124. Kersti Lilleväli. Gata3 and Gata2 in inner ear development. Tartu, 2007. 123 p.
- 125. Kai Rünk. Comparative ecology of three fern species: Dryopteris carthusiana (Vill.) H.P. Fuchs, D. expansa (C. Presl) Fraser-Jenkins & Jermy and D. dilatata (Hoffm.) A. Gray (Dryopteridaceae). Tartu, 2007. 143 p.
- 126. **Aveliina Helm.** Formation and persistence of dry grassland diversity: role of human history and landscape structure. Tartu, 2007. 89 p.
- 127. **Leho Tedersoo.** Ectomycorrhizal fungi: diversity and community structure in Estonia, Seychelles and Australia. Tartu, 2007. 233 p.
- 128. **Marko Mägi.** The habitat-related variation of reproductive performance of great tits in a deciduous-coniferous forest mosaic: looking for causes and consequences. Tartu, 2007. 135 p.
- 129. Valeria Lulla. Replication strategies and applications of Semliki Forest virus. Tartu, 2007. 109 p.
- 130. Ülle Reier. Estonian threatened vascular plant species: causes of rarity and conservation. Tartu, 2007. 79 p.
- 131. **Inga Jüriado**. Diversity of lichen species in Estonia: influence of regional and local factors. Tartu, 2007. 171 p.
- 132. **Tatjana Krama.** Mobbing behaviour in birds: costs and reciprocity based cooperation. Tartu, 2007. 112 p.
- 133. **Signe Saumaa.** The role of DNA mismatch repair and oxidative DNA damage defense systems in avoidance of stationary phase mutations in *Pseudomonas putida*. Tartu, 2007. 172 p.
- 134. **Reedik Mägi**. The linkage disequilibrium and the selection of genetic markers for association studies in european populations. Tartu, 2007. 96 p.
- 135. **Priit Kilgas.** Blood parameters as indicators of physiological condition and skeletal development in great tits (*Parus major*): natural variation and application in the reproductive ecology of birds. Tartu, 2007. 129 p.
- 136. **Anu Albert**. The role of water salinity in structuring eastern Baltic coastal fish communities. Tartu, 2007. 95 p.
- 137. **Kärt Padari.** Protein transduction mechanisms of transportans. Tartu, 2008. 128 p.
- 138. **Siiri-Lii Sandre.** Selective forces on larval colouration in a moth. Tartu, 2008. 125 p.
- 139. Ülle Jõgar. Conservation and restoration of semi-natural floodplain meadows and their rare plant species. Tartu, 2008. 99 p.
- 140. Lauri Laanisto. Macroecological approach in vegetation science: generality of ecological relationships at the global scale. Tartu, 2008. 133 p.
- 141. **Reidar Andreson**. Methods and software for predicting PCR failure rate in large genomes. Tartu, 2008. 105 p.

- 142. Birgot Paavel. Bio-optical properties of turbid lakes. Tartu, 2008. 175 p.
- 143. **Kaire Torn.** Distribution and ecology of charophytes in the Baltic Sea. Tartu, 2008, 98 p.
- 144. **Vladimir Vimberg.** Peptide mediated macrolide resistance. Tartu, 2008, 190 p.
- 145. **Daima Örd.** Studies on the stress-inducible pseudokinase TRB3, a novel inhibitor of transcription factor ATF4. Tartu, 2008, 108 p.
- 146. Lauri Saag. Taxonomic and ecologic problems in the genus *Lepraria* (*Stereocaulaceae*, lichenised *Ascomycota*). Tartu, 2008, 175 p.
- 147. Ulvi Karu. Antioxidant protection, carotenoids and coccidians in greenfinches – assessment of the costs of immune activation and mechanisms of parasite resistance in a passerine with carotenoid-based ornaments. Tartu, 2008, 124 p.
- 148. Jaanus Remm. Tree-cavities in forests: density, characteristics and occupancy by animals. Tartu, 2008, 128 p.
- 149. **Epp Moks.** Tapeworm parasites *Echinococcus multilocularis* and *E. granulosus* in Estonia: phylogenetic relationships and occurrence in wild carnivores and ungulates. Tartu, 2008, 82 p.
- 150. Eve Eensalu. Acclimation of stomatal structure and function in tree canopy: effect of light and CO₂ concentration. Tartu, 2008, 108 p.
- 151. **Janne Pullat**. Design, functionlization and application of an *in situ* synthesized oligonucleotide microarray. Tartu, 2008, 108 p.
- 152. Marta Putrinš. Responses of *Pseudomonas putida* to phenol-induced metabolic and stress signals. Tartu, 2008, 142 p.
- 153. Marina Semtšenko. Plant root behaviour: responses to neighbours and physical obstructions. Tartu, 2008, 106 p.
- 154. **Marge Starast.** Influence of cultivation techniques on productivity and fruit quality of some *Vaccinium* and *Rubus* taxa. Tartu, 2008, 154 p.
- 155. Age Tats. Sequence motifs influencing the efficiency of translation. Tartu, 2009, 104 p.
- 156. **Radi Tegova.** The role of specialized DNA polymerases in mutagenesis in *Pseudomonas putida*. Tartu, 2009, 124 p.
- 157. **Tsipe Aavik.** Plant species richness, composition and functional trait pattern in agricultural landscapes the role of land use intensity and landscape structure. Tartu, 2008, 112 p.
- 158. **Kaja Kiiver.** Semliki forest virus based vectors and cell lines for studying the replication and interactions of alphaviruses and hepaciviruses. Tartu, 2009, 104 p.
- 159. **Meelis Kadaja.** Papillomavirus Replication Machinery Induces Genomic Instability in its Host Cell. Tartu, 2009, 126 p.
- 160. **Pille Hallast.** Human and chimpanzee Luteinizing hormone/Chorionic Gonadotropin beta (*LHB/CGB*) gene clusters: diversity and divergence of young duplicated genes. Tartu, 2009, 168 p.

- 161. Ain Vellak. Spatial and temporal aspects of plant species conservation. Tartu, 2009, 86 p.
- 162. **Triinu Remmel.** Body size evolution in insects with different colouration strategies: the role of predation risk. Tartu, 2009, 168 p.
- 163. **Jaana Salujõe.** Zooplankton as the indicator of ecological quality and fish predation in lake ecosystems. Tartu, 2009, 129 p.
- 164. Ele Vahtmäe. Mapping benthic habitat with remote sensing in optically complex coastal environments. Tartu, 2009, 109 p.
- 165. Liisa Metsamaa. Model-based assessment to improve the use of remote sensing in recognition and quantitative mapping of cyanobacteria. Tartu, 2009, 114 p.
- 166. **Pille Säälik.** The role of endocytosis in the protein transduction by cellpenetrating peptides. Tartu, 2009, 155 p.
- 167. Lauri Peil. Ribosome assembly factors in *Escherichia coli*. Tartu, 2009, 147 p.
- Lea Hallik. Generality and specificity in light harvesting, carbon gain capacity and shade tolerance among plant functional groups. Tartu, 2009, 99 p.
- 169. **Mariliis Tark.** Mutagenic potential of DNA damage repair and tolerance mechanisms under starvation stress. Tartu, 2009, 191 p.
- 170. **Riinu Rannap.** Impacts of habitat loss and restoration on amphibian populations. Tartu, 2009, 117 p.
- 171. **Maarja Adojaan.** Molecular variation of HIV-1 and the use of this knowledge in vaccine development. Tartu, 2009, 95 p.
- 172. **Signe Altmäe.** Genomics and transcriptomics of human induced ovarian folliculogenesis. Tartu, 2010, 179 p.
- 173. **Triin Suvi.** Mycorrhizal fungi of native and introduced trees in the Seychelles Islands. Tartu, 2010, 107 p.
- 174. **Velda Lauringson.** Role of suspension feeding in a brackish-water coastal sea. Tartu, 2010, 123 p.
- 175. **Eero Talts.** Photosynthetic cyclic electron transport measurement and variably proton-coupled mechanism. Tartu, 2010, 121 p.
- 176. Mari Nelis. Genetic structure of the Estonian population and genetic distance from other populations of European descent. Tartu, 2010, 97 p.
- 177. **Kaarel Krjutškov.** Arrayed Primer Extension-2 as a multiplex PCR-based method for nucleic acid variation analysis: method and applications. Tartu, 2010, 129 p.
- 178. Egle Köster. Morphological and genetical variation within species complexes: *Anthyllis vulneraria* s. l. and *Alchemilla vulgaris* (coll.). Tartu, 2010, 101 p.
- 179. Erki Õunap. Systematic studies on the subfamily Sterrhinae (Lepidoptera: Geometridae). Tartu, 2010, 111 p.
- 180. **Merike Jõesaar.** Diversity of key catabolic genes at degradation of phenol and *p*-cresol in pseudomonads. Tartu, 2010, 125 p.

- 181. **Kristjan Herkül.** Effects of physical disturbance and habitat-modifying species on sediment properties and benthic communities in the northern Baltic Sea. Tartu, 2010, 123 p.
- 182. Arto Pulk. Studies on bacterial ribosomes by chemical modification approaches. Tartu, 2010, 161 p.
- 183. **Maria Põllupüü.** Ecological relations of cladocerans in a brackish-water ecosystem. Tartu, 2010, 126 p.
- 184. **Toomas Silla.** Study of the segregation mechanism of the Bovine Papillomavirus Type 1. Tartu, 2010, 188 p.
- 185. **Gyaneshwer Chaubey.** The demographic history of India: A perspective based on genetic evidence. Tartu, 2010, 184 p.
- 186. **Katrin Kepp.** Genes involved in cardiovascular traits: detection of genetic variation in Estonian and Czech populations. Tartu, 2010, 164 p.
- 187. Virve Sõber. The role of biotic interactions in plant reproductive performance. Tartu, 2010, 92 p.
- 188. Kersti Kangro. The response of phytoplankton community to the changes in nutrient loading. Tartu, 2010, 144 p.
- 189. Joachim M. Gerhold. Replication and Recombination of mitochondrial DNA in Yeast. Tartu, 2010, 120 p.
- 190. Helen Tammert. Ecological role of physiological and phylogenetic diversity in aquatic bacterial communities. Tartu, 2010, 140 p.
- 191. **Elle Rajandu.** Factors determining plant and lichen species diversity and composition in Estonian *Calamagrostis* and *Hepatica* site type forests. Tartu, 2010, 123 p.
- 192. **Paula Ann Kivistik.** ColR-ColS signalling system and transposition of Tn4652 in the adaptation of *Pseudomonas putida*. Tartu, 2010, 118 p.
- 193. Siim Sõber. Blood pressure genetics: from candidate genes to genomewide association studies. Tartu, 2011, 120 p.
- 194. **Kalle Kipper.** Studies on the role of helix 69 of 23S rRNA in the factordependent stages of translation initiation, elongation, and termination. Tartu, 2011, 178 p.
- 195. **Triinu Siibak.** Effect of antibiotics on ribosome assembly is indirect. Tartu, 2011, 134 p.
- 196. **Tambet Tõnissoo.** Identification and molecular analysis of the role of guanine nucleotide exchange factor RIC-8 in mouse development and neural function. Tartu, 2011, 110 p.
- 197. **Helin Räägel.** Multiple faces of cell-penetrating peptides their intracellular trafficking, stability and endosomal escape during protein transduction. Tartu, 2011, 161 p.
- 198. Andres Jaanus. Phytoplankton in Estonian coastal waters variability, trends and response to environmental pressures. Tartu, 2011, 157 p.
- 199. **Tiit Nikopensius.** Genetic predisposition to nonsyndromic orofacial clefts. Tartu, 2011, 152 p.

- 200. **Signe Värv.** Studies on the mechanisms of RNA polymerase II-dependent transcription elongation. Tartu, 2011, 108 p.
- 201. Kristjan Välk. Gene expression profiling and genome-wide association studies of non-small cell lung cancer. Tartu, 2011, 98 p.
- 202. Arno Põllumäe. Spatio-temporal patterns of native and invasive zooplankton species under changing climate and eutrophication conditions. Tartu, 2011, 153 p.
- 203. Egle Tammeleht. Brown bear (*Ursus arctos*) population structure, demographic processes and variations in diet in northern Eurasia. Tartu, 2011, 143 p.