

PETR KOHOUT

Ecology of ericoid mycorrhizal fungi



DISSERTATIONES BIOLOGICAE UNIVERSITATIS TARTUENSIS

349

PETR KOHOUT

Ecology of ericoid mycorrhizal fungi



UNIVERSITY OF TARTU
Press

Department of Botany, Institute of Ecology and Earth Sciences, Faculty of Science and Technology, University of Tartu, Estonia

Dissertation was accepted for the commencement of the degree of Doctor philosophiae in Botany and Mycology at the University of Tartu on 29 August 2018 by the Scientific Council of the Institute of Ecology and Earth Sciences, University of Tartu.

Supervisor: prof. Leho Tedersoo, University of Tartu, Estonia,
 prof. Urmas Kõljalg, University of Tartu, Estonia

Opponent: Prof. Björn Lindahl,
 Swedish University of Agricultural Sciences

Commencement: A. Vaga auditorium, Lai 40, Tartu; 9:15, 10 December 2018

Publication of this thesis is granted by the Institute of Ecology and Earth Sciences, University of Tartu.

ISSN 1024-6479

ISBN 978-9949-77-857-7 (print)

ISBN 978-9949-77-858-4 (pdf)

Copyright: Petr Kohout, 2018
Tartu University

University of Tartu Press
www.tyk.ee

CONTENTS

| | |
|--|-----|
| LIST OF ORIGINAL PUBLICATIONS | 6 |
| INTRODUCTION..... | 7 |
| Aims and hypothesis | 9 |
| MATERIALS AND METHODS | 10 |
| Sampling sites and study design..... | 10 |
| Data sources | 11 |
| Isolation of ericoid mycobionts and resynthesis experiments..... | 12 |
| Molecular analyses..... | 13 |
| Bioinformatics and phylogenetic analyses..... | 14 |
| Statistical analyses..... | 15 |
| RESULTS AND DISCUSSION | 17 |
| Identification of ericoid mycorrhizal fungi | 17 |
| Community composition of ericoid mycorrhizal fungi..... | 19 |
| CONCLUSIONS | 22 |
| SUMMARY | 23 |
| SUMMARY IN ESTONIAN | 25 |
| ACKNOWLEDGEMENTS | 26 |
| REFERENCES..... | 27 |
| PUBLICATIONS | 33 |
| CURRICULUM VITAE | 159 |
| ELULOOKIRJELDUS..... | 163 |

LIST OF ORIGINAL PUBLICATIONS

This thesis is based on the following publications that are referred in the text by their Roman numerals:

- I. Tedersoo L, Abarenkov K, Nilsson RH, Schussler A, Grelet GA, **Kohout P**, Oja J, Bonito GM, Veldre V, Jairus T, Ryberg M, Larsson KH, Kõljalg U. 2011. Tidying Up International Nucleotide Sequence Databases: Ecological, Geographical and Sequence Quality Annotation of ITS Sequences of Mycorrhizal Fungi. *Plos One* 6.
- II. Vohník M, Sadowsky JJ, **Kohout P**, Lhotáková Z, Nestby R, Kolařík M. 2012. Novel Root-Fungus Symbiosis in Ericaceae: Sheathed Ericoid Mycorrhiza Formed by a Hitherto Undescribed Basidiomycete with Affinities to Trechisporales. *Plos One* 7.
- III. Vohník M, Mrnka L, Lukešová T, Bruzone MC, **Kohout P**, Fehrer J. 2013. The cultivable endophytic community of Norway spruce ectomycorrhizas from microhabitats lacking ericaceous hosts is dominated by ericoid mycorrhizal *Meliniomyces variabilis*. *Fungal Ecology* 6: 281–292.
- IV. Lukešová T, **Kohout P**, Větrovský T, Vohník M. 2015. The Potential of Dark Septate Endophytes to Form Root Symbioses with Ectomycorrhizal and Ericoid Mycorrhizal Middle European Forest Plants. *Plos One* 10.
- V. **Kohout P**. 2017. Biogeography of Ericoid Mycorrhiza, in: Tedersoo, L. (Ed.), *Biogeography of Mycorrhizal Symbiosis*. Springer International Publishing, pp. 179–193.
- VI. **Kohout P**, Tedersoo L. 2017. Effect of soil moisture on root-associated fungal communities of *Erica dominans* in Drakensberg mountains in South Africa. *Mycorrhiza* 27: 397–406.
- VII. **Kohout P**, Bahram M, Põlme S, Tedersoo L. 2017. Elevation, space and host plant species structure Ericaceae root-associated fungal communities in Papua New Guinea. *Fungal Ecology* 30: 112–121.
- VIII. Põlme S, Bahram M, Jacquemyn H, Kennedy P, **Kohout P**, Moora M, Oja J, Öpik M, Pecoraro L, Tedersoo L. 2018. Host preference and network properties in biotrophic plant–fungal associations. *New Phytologist* 217: 1230–1239.

Author's contribution to each publication:

| | I | II | III | IV | V | VI | VII | VIII |
|-----------------------------|------|----|-----|----|------|-----|-----|------|
| Idea and design | – | – | – | ** | *** | *** | * | – |
| Sampling or data collection | * | – | * | * | *** | *** | – | * |
| Molecular analysis | n.a. | * | ** | * | n.a. | *** | *** | n.a. |
| Data analysis | – | * | * | ** | *** | *** | *** | – |
| Writing | * | ** | * | ** | *** | *** | *** | * |

* moderate contribution, ** high contribution, *** leading role

INTRODUCTION

Mycorrhizal symbiosis is a mutualistic partnership between plants and fungi that represents one of the oldest and the most widespread symbioses on the Earth (Redecker et al. 2000). It has been estimated that approximately 80% of vascular plant species form symbiosis with mycorrhizal fungi (Brundrett 2009). Mycorrhizal fungi play a crucial role in water and nutrient uptake to the host plant. They also enhance host plant defense mechanisms against pathogens and facilitate their growth in environments with high levels of heavy metals. In return, mycorrhizal plants provide carbohydrates, such as glucose and sucrose, to their symbiotic partners (Smith and Read 2008).

Several mycorrhizal types exist that have evolved independently multiple times during evolution for the last 500 million years. Arbuscular mycorrhizas (AM) evolved concurrently with the first colonization of land by plants some 450–500 million years ago, while ectomycorrhizas (EcM) evolved about 200 million years ago (Cairney 2000). One of the youngest mycorrhizal type is ericoid mycorrhizal (ErM) symbiosis, a mutualistic relationship formed between species belonging to several lineages of the Ericaceae family and diverse group of soil fungi. The first appearance of Ericaceae-like plants dates back to 90–75 million years (Nixon and Crepet 1993; Carpenter et al. 2015). It has been hypothesized that ErM symbiosis may have evolved in the same time frame (Cairney 2000). Ericoid mycorrhiza (one of the so-called endomycorrhizal types) is characterized by the intensive fungal colonization of the outermost root cell layer. Mycorrhizal fungi form a coiled intracellular hyphal complex. The fungal hyphae within the plant cell are usually hyaline with a thin cell wall. The plant plasma membrane of the root cells invaginates to envelope the fungal structures, but it is separated from the fungal cell by an interfacial matrix. This represents the interface between the two symbionts, where nutrient exchange takes place (Smith and Read 2008).

Ericoid mycorrhizal plants often occur on extremely poor soils, where most of the nutrients are locked up in complex forms of soil organic matter, with restricted biological availability. The ErM symbiosis represents a key evolutionary adaptation of ErM plants to mobilize the nutrients from such recalcitrant substrates (Kerley and Read 1998). However, ericoid mycorrhiza remains largely overlooked compared to the more common mycorrhizal types, such as AM and EcM, and a broader general understanding of the ErM symbiosis is lacking.

The Ericaceae family comprises 9 subfamilies, 124 genera, and approximately 4,250 species (Kron et al. 2002). Only the basal evolutionary lineages of the Ericaceae, namely, Enkianthoideae, Arbutoideae, Pyroloideae, and Monotropeoideae, lack the capability to form ErM. The earliest diverging lineage Enkianthoideae, represented by the sole genus *Enkianthus*, forms arbuscular mycorrhizal symbiosis (Gorman and Starrett 2003; Abe 2005). Instead, species of the Monotropeoideae subfamily form the so-called monotropoid mycorrhizal symbiosis (characterized by ectendomycorrhizal anatomical structures) with

specific groups of ectomycorrhizal fungi (EcMF) from the Basidiomycota phylum (Hynson and Bruns 2009), while members of the Arbutioideae and Pyroloideae subfamilies host a wide spectrum of EcM mycobionts in their roots (Krpata et al. 2007). Okuda et al. (2011) described a symbiosis resembling ErM in *Schizocodon soldanelloides* (Diapensiaceae) roots, but this requires independent confirmation. So far, the only sufficiently confirmed ErM plant species belong to the Cassiopoideae, Ericoideae, Harrimanelloideae, Styphelioideae (formally known as Epacridaceae), and Vaccinioideae subfamilies.

Compared to more common mycorrhiza types, such as AM and EcM, our knowledge about the diversity of ericoid mycorrhizal fungi (ErMF) is very superficial. While arbuscular mycorrhizal fungi (AMF) have a monophyletic origin, the ability to form ErM as well as EcM evolved independently multiple times in several fungal lineages (Smith and Read 2008). Earlier attempts to determine fungal diversity were based on direct observations of macroscopic (fungal fruit bodies) as well as microscopic (e.g., spores) structures. These methods allowed researchers to classify AMF to morphospecies based on their chlamydospore anatomy. Similarly, EcMF were classified based on the morphology and anatomy of the EcM colonization structure formed by each individual unique plant-fungal species combination (Agerer 1987–2006). Although these early methods suffered from many drawbacks, their implementation enabled us to classify uncultured fungal species, which would have been completely overlooked and uncommunicated otherwise. Subsequent implementation of molecular methods for fungal species determination boosted up our knowledge of AM as well as EcMF diversity (Õpik et al. 2014; Tedersoo et al. 2010; Tedersoo et al. 2014a). On the other hand, research focused on the ErMF diversity suffered from much more serious drawbacks. Determination of the ErMF lifestyle can neither be based on the phylogenetic affinity to any known lineage as it is the case of AMF and EcMF to some extent nor can ErM lifestyle be defined based on the occurrence of fungal species in Ericaceae root segments, because Ericaceae roots can also harbor non-mycorrhizal fungi (Bougoure and Cairney 2005a, b). Therefore, mycorrhizal resynthesis experiments are needed to describe the character of the association between the host plant and mycobiont and to sufficiently prove the ericoid mycorrhizal lifestyle of Ericaceae-associated mycobionts (Leake and Read 1991). Anatomical features of ErM symbiosis were described above as well as more specifically in Smith and Read (2008). Alternatively, methods applying transmission electron microscopy associated with molecular methods of fungal detection can be used in specific cases (Selosse et al. 2007).

The ErMF belong to several fungal lineages of Ascomycota as well as Basidiomycota. Nowadays, there are few sufficiently proven ErMF, which belong to four taxonomic groups, Helotiales, Chaetothyriales (Ascomycota) and Sebaciniales (Basidiomycota) (Allen et al. 2003; Selosse et al. 2007; Tedersoo et al. 2011; Vohnik et al. 2016). The most comprehensive knowledge about ericoid mycorrhizal lifestyle in fungi is derived from the *Pezoloma ericae* aggregate (PEA; previously known as *Hymenoscyphus ericae* or *Rhizoscyphus*

ericae aggregate), which contains several species (*Pezoloma ericae* and *Meliniomyces variabilis*) known as ErM symbionts of Ericaceae (Hambleton and Sigler, 2005). Besides the PEA, *Oidiodendron maius* is another well studied ErMF. Its sequenced genome offered the first insights into the evolution of ErM (Kohler et al. 2015).

Our current knowledge about the ErMF diversity and ecology largely relates to the Northern Hemisphere, where PEA (e.g. Vralstad et al. 2002; Usuki et al. 2003; Bougoure et al. 2007; Walker et al. 2011; Gorzelak et al. 2012) or Sebaciniales (Allen et al. 2003) often dominate ErMF communities associated with Ericaceae. Compared to the Northern Hemisphere, there are only a few studies addressing ErMF diversity and ecology in the Southern Hemisphere, almost exclusively from Australia (e.g. Williams et al. 2004; Bougoure and Cairney, 2005a,b). These studies showed that Australian Ericaceae interact with distinct species of Helotiales, although some of them are closely related to PEA but distinct such as the recently described *Cairneyella variabilis* (Midgley et al. 2016). On the contrary, *C. variabilis* has never been recorded outside Australia. Much more data is however needed before comprehensive insights into ErMF biogeography can be obtained.

Besides the global distribution of ErMF, composition of Ericaceae root associated fungal communities can be also affected by local environmental factors such as soil chemistry (Hazard et al. 2014), elevation (Gorzelak et al. 2012) and/or vegetation type (Bougoure et al. 2007). Besides that, host identity might also play an important role in structuring mycorrhizal fungal communities (e.g. Vandenkoornhuyse et al., 2002; Sýkorová et al., 2007). Previous studies on host preference in ErM systems provide inconsistent patterns. Although Bougoure et al. (2007) showed differences in ErMF community composition between *Calluna vulgaris* and *Vaccinium myrtillus*, subsequent more inclusive studies recovered no host effect on ErMF communities (Kjoller et al., 2010; Walker et al., 2011). All these studies were conducted at high latitudes, where the diversity of Ericaceae is relatively low. As recently shown, addressing host effect on mycorrhizal fungal communities might have higher significance in regions with higher plant biodiversity, where mycorrhizal fungi might contribute to speciation or species co-occurrence of their host plants (Waterman et al., 2011; Nurfadilah et al., 2013).

Aims and hypothesis

In this thesis, I focus on Ericaceae associated fungi from various aspects of the partnership, because the ericoid mycorrhizal symbiosis is probably the most overlooked mycorrhizal symbiotic type. Firstly, this thesis aims to determine the occurrence of ericoid mycorrhizal life-style among fungi (**I**, **II**, **III**, **IV**). Furthermore, I also focused on determination of environmental factors which significantly affect composition of fungal communities associated with roots of ericoid mycorrhizal plants (**VI**, **VII**, **VIII**). Last but not least, I addressed global distribution and biogeography of ericoid mycorrhizal fungi (**V**).

MATERIALS AND METHODS

Sampling sites and study design

To determine the occurrence of ericoid mycorrhizal life-style among fungi, we chose one sampling site in Norway (II) and two sampling sites in Czechia (III, IV). We repeatedly collected root samples of *Vaccinium myrtillus* in a forest plantation and a nearby natural forest in mid-Norway (II). The mats (plants and adhering soil, approx. 40×30×15 cm) of blueberry with some co-occurring *Vaccinium vitis-idaea* were taken from a regenerating *Picea abies* stand. Altogether 20 mats samples were collected between October 2010 and May 2011. Upon receipt in the laboratory, roots were washed free of the adhering substrate and stored at 5 °C until processed. Half of the ericoid roots were used for assessment of fungal colonization (using light microscopy and scanning electron microscopy) and the second half was used for isolation of root associated fungi.

In Czechia, one site was chosen within the Bohemian Forest NP in *P. abies* dominated forest with ericoid understory (*V. myrtillus* and *V. vitis-idaea*). In total, nineteen 3–5 year old spruce seedlings inhabiting different niches within the two sites were sampled on two occasions during the vegetation season (III). Seedlings were carefully dug up not to destroy their fine ectomycorrhizal roots, packed in plastic bags and stored at 5 °C until the isolation of the ectomycorrhiza-associated fungi on the following day. The second sampling site in Czechia was situated in Bohemian Switzerland NP (IV). Fifteen soil samples were collected on the site dominated by *Pinus sylvestris* forest with understory of ericaceous plants (*V. myrtillus*, *V. vitis-idaea* and *Calluna vulgaris*) on podzolic soil. Ericaceae roots were washed under tap water and separated into three fractions: the first part was used for measurement of fungal colonization (using light microscopy), the second part for mycobionts isolation and the third part was used for direct isolation of DNA and subsequent molecular determination of root associated fungi.

To determine the environmental factors, which affect composition of fungal communities associated with roots of ericoid mycorrhizal plants, we sampled roots of Ericaceae plants on Mount Wilhelm in Papua New Guinea (VI) and in Drakensberg mountain range in the Republic of South Africa (VII). Mount Wilhelm (4509 m.a.s.l.) is located within the Bismarck Range. An elevation transect with six sampling sites (70×70 m) was established in November 2011. The transect was situated on the eastern slope of Mt. Wilhelm at the following elevations: i) 4483 m.a.s.l., ii) 4266 m.a.s.l., iii) 4044 m.a.s.l., iv) 3830 m.a.s.l. v) 3600 m.a.s.l., and vi) 3387 m.a.s.l.. Five separated root samples (each at least 10 m distance from another one) of *Acrothamnus* sp. were taken at each site. For studying the host and spatial effects on putative ericoid mycorrhizal fungi and endophytic fungal diversity, the 3600 m site was chosen, because it had the highest number of Ericaceae species. All collected root samples were cleaned

from the attached soil and dried on silica gel. Each root sample contained 20–25 mg of fresh hair root mass. Roots were transported to a lab where they were refreshed by rinsing in sterile tap water for several minutes. Ericaceae hair roots were surface sterilized in 10% commercial house bleach (3% chlorine) for one minute. The fine roots were subsequently rinsed in sterile tap water, dried and stored at -20°C . All samples were analyzed separately, without any subsequent pooling.

In Drakensberg mountain range, we selected a study site on a slope of Champagne Castle (VII). Dominant vegetation represented open grassland with small patches of primeval forest. We selected nine sites (across an area of 10 km^2). On each site, we established two plots with contrasting soil moisture level. The dry microhabitat with low water content was dominated by sparse vegetation on shallow soils, typically on large erratic boulders with limited access to ground water. We selected a common Ericaceae species (*Erica dominans*), which occurred on all sites in both microhabitats to study differences in root-associated fungal communities. On each, plot one healthy-looking *E. dominans* plant was sampled. All collected root samples were cleaned from the attached soil. Ericaceae hair roots were surface sterilized in 10% commercial house bleach (3% of chlorine) for 1 min and subsequently rinsed in sterile tap water, dried and kept in silica gel. Each root sample contained 20 to 25 mg of fresh fine hair roots.

Data sources

Two studies included in the thesis were based on data collected from publicly available sources, such as the International Nucleotide Sequence Databases consortium (INSDc) (I) or publications (VIII). Study I aimed to extend the trait annotation of fungal INSDc sequences to provide insights into the biodiversity and ecology of mycorrhizal fungi. All fungal sequences (annotated as such in INSDc) of internal transcribed spacer (ITS) of ribosomal DNA (rDNA) were downloaded from INSDc to UNITE (Abarenkov et al. 2010a). Very short sequences (<200 bp) and sequences derived from Next Generation Sequencing techniques – that are normally not allowed in INSD – were excluded. Sequences were annotated by experts on particular mycorrhizal types and/or taxonomic groups. For EcM, AM, ErM and orchid mycorrhizal (OrM) fungi, all representatives of the major mycorrhizal taxonomic groups were retrieved through the use of names of the inclusive taxa as search strings in the organism field in the PlutoF workbench (Abarenkov et al. 2010b). All sequences that were poorly aligned to other species were subjected to bulk megablast searches against INSD and UNITE as implemented in the PlutoF workbench. This enabled us to identify potentially chimeric and reverse complementary sequences as well as sequences belonging to non-targeted taxa. Most of the chimeric and low-quality sequences were discovered by carefully inspecting the alignment. Potentially low-quality sequences were primarily recognized as sequences with unique gaps

and indels in the conserved regions, especially in the 5.8S gene, as compared to their closest sequences. Sequences were also considered of low quality when the beginning or end of the ITS spacers contained >2 obvious substitution errors or indels resulting from inadequate end trimming. Sequences passing the quality control steps were re-aligned with MAFFT; the alignments were corrected manually and subjected to Maximum Likelihood analyses using RAxML (<http://phylobench.vital-it.ch/raxml-bb/>) or PhyML (<http://www.biportal.uio.no/appinfo/show.php?app=phym1>) with default options. Sequences with disproportionately long branches were, once again, checked for potential chimeric insertions and low quality.

The second study (VIII) aimed to assess the generality of organizational patterns in biotrophic plant–fungal symbioses and builds on individual case studies that were compiled from the Web of Science by combining the search terms ‘host specificity’, ‘host preference’ and ‘host effect’ with ‘mycorrhiza’ and ‘endophytes’. The analysis includes studies in which at least two host plant species were sampled in multiple replicates per study area and fungi were identified using either molecular or morphological methods. In total, we were able to compile 67% datasets out of 73 studies that were regarded as suitable. In most datasets, taxa were delimited using molecular methods and termed as operational taxonomic units (OTUs).

The datasets of plant–fungal associations were categorized into the following guilds: AM, OrM, EcM, ErM, root endophytes and leaf endophytes. For each site, metadata on various geographic, floristic and sampling variables were retrieved from the original publications. Approximate mean annual temperature (MAT) and precipitation (MAP) were retrieved from a high resolution database of the Earth’s surface climate (Hijmans et al., 2005) using the software ARCMAP 10.3 (ESRI, Redlands, CA, USA). For each dataset, we calculated the average phylogenetic distance (APD) among hosts using the online phylogenetic query tool Phylomatic (<http://phylodiversity.net/phyloomatic/>).

Isolation of ericoid mycobionts and resynthesis experiments

To determine the occurrence of ericoid mycorrhizal lifestyle in fungi and ecology of ericoid mycorrhizal fungi, we isolated root associated fungi from sampled Ericaceae plants (II, IV) or seedlings of *P. abies* (III). Isolation of Ericaceae was performed from approx. 5-mm pieces of surface-sterilized (30 s in 10% house bleach containing 4.5% of chlorine and rinsed twice in autoclaved water) roots and cultivation on modified Melin Norkrans agar (MMN). All roots were incubated in the dark at 20 °C for 21 days.

Ecological lifestyle of isolated fungi (and other roots associated fungi previously isolated in other studies) was assessed in number of resynthesis experiments with *V. myrtilus* (II, III, IV), *P. abies* (III, IV) and *Betula pendula*

(IV). Seeds of the selected plant species were surface-sterilized with 30% hydrogen peroxide and placed on MMN agar. Non-contaminated seeds were left to germinate for 3 weeks. Lower compartments of split Petri dishes (9 cm in diam.) with perforated central septa were filled with MMN (10x diluted sugars), overlaid with a sterile cellophane membrane to prevent growth of roots into the medium and inoculated with agar plugs cut from margins of colonies of selected fungal isolates actively growing on MMN. The petri dishes with the plugs were left for 3 weeks at room temperature in the dark to produce vegetative mycelium. Then, 3-week-old seedlings were transferred to the dishes, the empty upper compartments accommodating their shoots and the lower compartments their roots. The roots of the seedlings were covered with a piece of sterile moistened filter paper. The dishes were sealed with air-permeable film, lower compartments covered with aluminum foil, and placed in a vertical position in a growth chamber (16/8 hr and 21/15 °C day/night cycle, irradiation 250 $\mu\text{mol m}^{-2} \text{s}^{-1}$). After 5 months, the plants were extracted and their roots were separated and processed as follows: *V. myrtillus* roots were cleared with 10% KOH (20 min at 121 °C), washed with tap water, acidified (1 min in 3% HCl), washed with tap water, stained with trypan blue (60 min at 121 °C) and de-stained overnight in lactoglycerol; *P. abies* and *B. pendula* roots were hand-sectioned and thin sections were stained with aniline or trypan blue. The stained roots and thin sections were observed using an Olympus BX60 microscope equipped with DIC at high magnification (400x and 1000x).

Molecular analyses

Taxonomic determination of root associated fungal isolates was mostly done by sequencing of ITS region (II, III). In some cases, small (SSU) or large (LSU) subunits of rDNA were used for more precise phylogenetical placement of the obtained isolates. Fungal DNA was extracted from fresh mycelia using the sorbitol method (Štorchová et al. 2000). DNA amplifications of the rDNA regions were performed using variety of eukaryotic primers. PCR products were checked for length and quality/quantity by gel electrophoresis (1.5% agarose) and purified using the High Pure PCR product purification kit (Roche Holding Ltd, Switzerland). Sanger sequencing was done by GATC Biotech AG (Germany) using the PCR primers, sequence electropherograms were edited manually. The obtained sequences were subjected to BLAST searches and submitted to the GenBank database.

To determine fungal communities associated with Ericaceae roots, we used the 454 pyrosequencing method (IV, VI, VII). Development of high throughput sequencing methods (such as 454 pyrosequencing) enables metagenomic analyses in a manner that exceeds the capacity of traditional Sanger sequencing-based approaches by several orders of magnitude. The DNA of all samples were extracted using a PowerSoil DNA Isolation Kit (MoBio, Carlsbad, CA, USA), following the manufacturer's instructions. For identification of fungi, we

selected the full ITS region (Schoch et al. 2012), amplified with ITS1Fngs, ITS1ngs, and ITS4ngs primers. Each of these primers was supplemented with a 10–12 base multiplex identifier (MID) tag in the 5' end (at least four differences to each other). PCR was performed in four replicates for each primer pair using 25 cycles. The amplicons were checked for the presence of a product on 1% agarose gel. In case of no visible band or a strong band, we repeated the amplification program by adjusting the number of cycles between 25 and 35. The PCR products were purified using Exo-Sap enzymes (Sigma, St. Louis, MO, USA), and 20 μ l of the purified PCR product was normalized using a Sequalprep™ Normalization Plate (96) Kit (Invitrogen Inc., Carlsbad, CA, USA) according to the manufacturer's instructions. The PCR products were pyrosequenced using the Roche GS FLX+ platform and Titanium chemistry in LGC Genomics (Berlin, Germany). Raw sequence data and associated metadata are available in UNITE repository.

Bioinformatics and phylogenetic analyses

Pyrosequencing reads (**IV**, **VI**, **VII**) were cleaned based on the quality information using Mothur 1.30.2 denoising algorithm (Schloss et al. 2009). Short sequences (<300 bp in length) and sequences possessing any mismatch to the primers were removed using the SEED 1.2.3 platform (Větrovský and Baldrian 2013). Sequences were demultiplexed based on the primer tags. Only reverse sequences from ITS4 were used for the subsequent data processing. Putative chimeras were identified and removed, and the remaining sequences were pooled and clustered into OTUs with USEARCH (Edgar et al. 2011) using 97% similarity level. We removed all global singletons, because nearly half of these are suggested to be artificial (Tedersoo et al. 2010). The remaining OTUs were taxonomically identified based on representative sequences using Biopython scripts for running BLASTn queries against the INSDc and UNITE. All sequences belonging to plants or other non-fungal groups were excluded from the dataset. We typically relied on 90, 85, 80, and 75% sequence identity as a criterion for assigning OTUs with names of a genus, family, order, or class, respectively. Representative sequencing of fungal OTUs (those with e-value $\geq e^{-50}$) were assigned to species hypothesis (SH) using UNITE (Kõljalg et al. 2013).

We assigned putative fungal ecology to obtained OTUs based on their taxonomic placement (**VI**, **VII**). The rough division of fungal OTUs into two ecological guilds, endophytic fungi (EndF) and putative ErMF was chosen, because the ability to form ericoid mycorrhizal symbiosis is not phylogenetically conserved among fungal species. In Helotiales, well known ErMF species are closely related to EcMF as well as non-mycorrhizal symbionts. On the contrary, many fungal taxa belonging to Helotiales, but distinct from the known ErMF species from the PEA, have been described as putative ericoid mycorrhizal symbionts (e.g. Grelet et al. 2009; Zhang et al. 2009). The ability to

form ericoid mycorrhizal symbiosis shows a similarly low phylogenetic signal in Sebaciales as well (Weiss et al. 2013). The third sufficiently confirmed group of ericoid mycorrhizal fungi belongs to Hymenochaetales (Kolařík and Vohník 2018). Therefore, fungal OTUs belonging to these three taxonomic groups were considered as putative ErMF.

Phylogenetic methods were used for more reliable identification of fungal isolates (**II**, **III**) or OTUs (**IV**, **V**, **VI**). Further sequences, derived preferentially from cultured isolates deposited in international culture collections, as well as suitable outgroup taxa were retrieved from INSDc to represent the genetic variability of the isolates or OTUs as comprehensively as possible. Sequences were aligned using the iterative refinement method of MAFFT (L-INS-i; <http://mafft.cbrc.jp/alignment/server/>). The alignment was checked and corrected manually using BioEdit V7.0.0 (Hall 1999). For phylogenetic analyses, maximum likelihood (ML) and Bayesian methods of inference were applied using MEGA 7 (Kumar et al. 2016) and MrBayes v3.2.6 (Ronquist and Huelsenbeck 2003). The ML analysis was done using GTR model with uniform rates among sites and NNI tree inference option. The Bayesian analysis was performed using the GTR substitution model with gamma distributed rate across the sites.

Statistical analyses

To test differences in colonization levels and root or shoot biomass production among experimental treatments (**II**, **III**, **IV**) we used the STATISTICA 12 software (StatSoft Inc., USA). Preferentially, we used parametric methods such as ANOVA. However, if the data did not meet the criteria of normal distribution and homogeneity of variances, the non-parametric Kruskal-Wallis test followed by a multiple-comparison z-value test were used.

To determine the drivers of fungal richness in Ericaceae-associated fungal communities (**IV**, **VI**, **VII**), we randomly subsampled the number of sequences per sample into the lowest number of sequences per sample, a procedure termed rarefaction. The general least squares (GLS) model was built to identify the main predictors of OTU richness, based on tested environmental variables using a routine implemented in the nlme package of R (Pinheiro et al., 2016). The best model was selected according to the corrected Akaike information criterion (AICc). Robustness of the best model was further evaluated by averaging models that fell into the 95% AICc confidence set.

Hellinger transformation was used to standardize fungal OTU abundance data across samples, whereas Bray-Curtis distance was used for computing dissimilarity matrices. The spatial distances between samples were included into subsequent analysis by reducing the Euclidean distance matrix into spatial Principal Coordinates of Neighbourhood Matrix (PCNM) vectors (Borcard and Legendre 2002). To address the relative importance of environmental factors and spatial distances between the samples on the fungal community structure,

we used PERMANOVA (**VI**, **VII**, **VIII**) as implemented in the `adonis` routine of the `vegan` package of R (Oksanen et al. 2012). Adonis tests the significance of discrete and continuous factors based on permutations. Adjusted R^2 was calculated based on the `adonis` results. Using the Bray-Curtis distance, we also constructed non-metric multidimensional scaling (NMDS) plots in the `Ecodist` package of R (Goslee and Urban 2007) for visualizing trends in fungal communities. Confidence ellipses for NMDS plots were calculated by the function `ordiellipse` in `Vegan` package. Indicator OTUs for different environments were determined using “`indVal`” function of the `labdsv` package of R (Roberts 2014), where the significance of indicator values is determined based on a permutation test. Only OTUs found in more than four samples were included in the analysis.

To assess network properties (**VIII**), we calculated nestedness and modularity metrics based on plant–fungi co-occurrence matrices. The modularity index of each dataset was calculated using a simulated annealing algorithm as implemented in `NetCarto` software (Guimerá & Amaral, 2005). We calculated the nestedness metric based on overlap and decreasing fill using the `NODF` function in the `bipartite` package of R. To test for differences among fungal guilds in host effect, nestedness and modularity, we applied a nonparametric multiple comparison procedure with an unbalanced one-way factorial design as implemented in the `gao_cs` function in the `nparcomp` package of R (Gao et al., 2008).

RESULTS AND DISCUSSION

Identification of ericoid mycorrhizal fungi

The study **I**, focused on occurrence of plant-fungal symbiotic life-styles among fungi, comprised 183,208 fungal ITS sequences. Of these, 28,791 (15.7%) sequences belonged to EcMF and 3,176 (1.7%) to AMF. In total, 1,457 (0.8%) and 2,267 (1.2%) sequences were recovered from roots of ErM plants and orchids, respectively. Metadata on interacting taxon were available for 6,272 (21.8%), 835 (26.3%), 1,093 (75.0%) and 1,608 (70.9%) entries of EcM, AM, ErM and OrM fungi, respectively. In AMF, plant roots, spores and soil contributed 41.2%, 33.9% and 19.0% to the source of isolation, respectively. In EcMF, fruit bodies, ectomycorrhizas and soil DNA accounted for 43.3%, 32.4% and 14.6% of the identification sources, respectively. In contrast to EcM and AM mycobionts, the fungi inhabiting roots of ericoid plants were identified directly from roots with or without a culturing step. In putatively ErMF, 690 (47.4%) sequences were obtained directly from ErM roots and 767 (52.6%) sequences were obtained from living cultures. In the cultured isolates, we could trace the symbiotic performance of 226 isolates in various experiments. Taken together, 60.2% of the isolates were capable of forming coils and/or stimulating growth of ericoid plants *in vitro*. More than 95% of the functional ErM mycobionts belonged to the Helotiales. Cultures identified as Hypocreales and Coniochaetales probably represent fast-growing contaminants, because these taxa have never been suggested as functional partners in ErM. While taxa from all fungal phyla have been identified from roots of ericoid plants, experimental evidence for functional association covers only a few, albeit large, groups of fungi. The remaining DNA-based identified taxa may belong either to unculturable mycorrhizal fungi or to non-mycorrhizal guilds of opportunistic pathogens, endophytes or saprobes (Allen et al. 2003; Walker et al. 2011). As an alternative to direct synthesis experiments, electron microscopy may provide *in situ* evidence for functional associations between plants and fungi at higher taxonomic levels (Moore et al. 1978).

A yet undescribed type of ericoid mycorrhizal association in *Vaccinium* spp. was observed in field-collected roots from semi-natural blueberry plantation in Norway (**II**). Its most prominent characteristic is a dense layer of clamp-bearing hyphae over the surface of terminal parts of young hair roots. The hyphae comprising the sheath were of variable diameter and penetrated epidermal cells, forming dense hyphal coils typical for ericoid mycorrhizae. Three isolates with distinct morphology and/or growth rate were obtained from 30 sheathed ErM roots. Both clamped basidiomycetes JPK 87 and JPK 90 grew well on MMN producing dense whitish colonies. The basidiomycete JPK 90 formed a long branch residing as sister to Trechisporales (PP = 0.98) in MB analyses, but being inconsistently clustered with Trechisporales, Hymenochaetales or Russulales in ML trees. Recent study, based on protein coding sequences and

morphology, placed the isolates within the Hymenochaetales order (Kolařík and Vohník 2018). On soil agar, sheathed ErM formed by the basidiomycetes JPK 87 and JPK 90 occurred in, but were not limited to, root apices and did not appear to follow a predictable pattern among root orders. The extent of intracellular colonization among inoculated plants was highly variable (coefficient of variation = 1.7). Significant correlations between ErM intracellular colonization and shoot length ($r = 0.74$, $p = 0.002$) and shoot weight ($r = 0.54$, $p = 0.04$), along with negligible shoot growth of non-(ErM)-colonized and non-inoculated control plants, indicate that colonization by the basidiomycetes JPK 87 and JPK 90 positively contributed to shoot growth. It is generally observed that ascomycetous ErM fungi do not produce developed extraradical hyphal mantles around colonized roots; so far the only described symbiosis of Ericaceae characteristically possessing hyphal mantles is cavendishoid ectendomycorrhiza. However, this symbiosis is formed by nonclamped hyphae of Sebaciales and its mantles are accompanied by intercellular fungal tissue resembling a Hartig net (Binder et al. 2005).

To determine the ecological interactions of ErMF (III), we focused on ascomycetes associating with basidiomycetous ectomycorrhizas of *Picea abies* in a temperate montane forest in central Europe, where ericoid shrubs dominate in forest understory. The 360 surface-sterilized basidiomycetous ectomycorrhizas yielded 128 relatively slow-growing non-sporulating isolates. Most of the isolates belonged to the PEA encompassing *Cadophora finlandica*, *Melinomyces bicolor*, *M. variabilis* and two unidentified ascomycetous isolates. In the resynthesis experiments, none of the *M. bicolor* nor *M. variabilis* isolates formed ectomycorrhiza or ectendomycorrhiza with spruce seedlings, but rather produced intracellular colonization in the root cortex cells which was usually accompanied with apparent darkening and thickening of their cell walls. In the same sampling sites, *M. variabilis* did not belong among the major soil fungal OTUs (Baldrian et al. 2012). These observations suggest that *M. variabilis* is an obligate biotroph rather than a free-living soil dweller. Additionally, *M. variabilis* seems to have the broadest ecological niche among the PEA members – it has been detected as an endophyte in the roots of a taxonomically diverse spectrum of plants including Fagaceae, Orchidaceae and Salicaceae (Hambleton and Sigler 2005), Betulaceae and Pinaceae (Kernaghan and Patriquin, 2011) and Poaceae and Primulaceae (Tejesvi et al. 2013), but seems to prefer Ericaceae, the only group which forms true mycorrhizas with *M. variabilis*. Except for one *M. variabilis* isolate, all the remaining PEA isolates colonized roots of *V. myrtillus* intracellularly. Most of the PEA isolates formed dense intracellular hyphal coils resembling ericoid mycorrhiza, including *C. finlandica* and *M. bicolor*. *C. finlandica* and *M. bicolor* have been previously detected as EcM co-associated fungi but they also form characteristic ecto- and ectendomycorrhizas (Wang & Wilcox 1985; Vralstad et al. 2000, 2002; Villarreal-Ruiz et al. 2004). Additionally, both species have been shown to form structures resembling ericoid mycorrhizas in vitro (Vralstad et al. 2002; Villarreal-Ruiz et al. 2004), but they are only rarely detected in

Ericaceae roots under natural conditions. Therefore, the potential of PEA fungi to potentially form common mycorrhizal mycelial network between ErM and EcM plants, seems to be rather ecologically insignificant.

The dual mycorrhizal ability of other Helotiales species (with affinity to *Phialocephala fortinii* s. l.—*Acephala applanata* species complex, PAC), that may form mycorrhizal links between Ericaceae and Pinaceae, was tested in resyntheses experiments in the subsequent study (IV). Roots of all inoculated plants (*P. abies* and *V. myrtillus*) possessed intraradical fungal colonization. All species belonging to PAC formed intracellular microsclerotia consisting of melanised or hyaline hyphae in both *P. abies* and *V. myrtillus*. In *P. abies*, microsclerotia were often found within the central stele. *Acephala macrosclerotiorum* colonized spruce intercellulary and formed a Hartig net and a parenchymatous hyphal net on the root surface resembling a loose hyphal mantle. This species formed darkly pigmented sclerotia on the surface of some roots, as previously showed by Münzenberger et al. (2009). Intracellular hyphal coils resembling ErM together with intracellular microsclerotia typical for endophytic colonization were observed in blueberry roots colonized by *A. macrosclerotiorum*. Therefore, *A. macrosclerotiorum* may potentially form a mycorrhizal link between *P. abies* and *V. myrtillus*. To test the occurrence of *A. macrosclerotiorum* in Ericaceae plants in situ, we chose a forest ecosystem where *A. macrosclerotiorum* dominates on *Pinus sylvestris* roots. Using direct isolation of DNA from Ericaceae roots with subsequent 454-sequencing of fungal ITS rDNA region, we did not obtain any sequences similar to *A. macrosclerotiorum*. It remains questionable, if mycorrhizal links between Ericaceae and Pinaceae occur in temperate or boreal forests.

Community composition of ericoid mycorrhizal fungi

The communities of Ericaceae root-associated fungi were significantly affected by environmental factors such as elevation gradient (VI, VII), host plant species (VII, VIII) as well as soil moisture content (VI). On the larger spatial scales, ericoid mycorrhizal fungi showed biogeographical patterns (V).

Although the ericoid mycorrhiza represents ecologically important symbiotic partnership, its global distribution has never been assessed previously (IV). In spite of the very broad distribution of the ErM plants (all continents except Antarctica), there are also many areas where ErM plants are missing, such as large parts of South America, SW Asia, and much of Africa and Australia. Compared to ErM host plants, much less is known about the distribution and global biogeography of their root-associated mycorrhizal symbionts. Traditionally, most of the studies focused on the diversity and community ecology of ErMF were performed on the Northern Hemisphere, particularly in Europe and North America. Preliminary insights indicate that some ErMF species (*Pezoloma ericae*) have a very broad distribution range. On the contrary, some

species have much narrower distribution range restricted to a single hemisphere (*Meliniomyces variabilis*) or continent (*Cairneyella variabilis*).

In the study **VII**, which focused on community ecology of Ericaceae roots associated fungi in Papua New Guinea, 670 OTUs were recovered from 24,008 sequences that passed through the denoising steps and removal of singletons and non-fungal sequences. The PERMANOVA test revealed significant effects of elevation and spatial vectors on putative ErMF and EndF community composition. Both variables (elevation and space) explained more variation in putative ErMF than EndF communities. The observed pattern in our study was mostly caused by the highest elevation site, which showed the greatest relative abundance of Helotiales. Broad ecological niche of the Helotiales species is well documented by numerous studies, describing them as abundant members of fungal communities from many different environments such as neotropical forests (Haug et al., 2004), glacier forefronts (Brown and Jumpponen, 2014), sub-Antarctic islands (Upson et al., 2007), or roots of submerged aquatic plants (Kohout et al., 2012). On the other hand, the second most widespread and well known ErMF order, Sebaciniales (Weiss et al., 2004), showed higher occurrence in the lowest-elevation sites. High relative abundance of Sebaciniales associated with Ericaceae roots was also found in other tropical mountain regions (Setaro and Kron, 2011), although our study is the first one focusing on sites above 2500 m.a.s.l. These findings support the ubiquity of Sebaciniales and their worldwide distribution (Oberwinkler et al., 2013; Tedersoo et al., 2014b). On the main study site, which was used to study host plant effect on Ericaceae associated fungal communities in Papua New Guinea, PERMANOVA analysis revealed a substantially different effect of Ericaceae host plant taxonomy on EndF and putative ErMF communities. While putative ErMF communities were significantly affected by host plant genus, EndF communities differed between host plant subfamilies. These results show that host plant taxonomic levels above the species level (Kjøller et al., 2010; Walker et al., 2011) play an important role in structuring EndF communities rather than putative ErMF communities. Differentiation of ErMF communities at lower taxonomic levels may promote coexistence of Ericaceae genera, if ErMF for example differ in exploitation of different nutrient sources (Cairney et al., 2000).

In study **VI**, which focused on community ecology of root-associated fungi of *Erica dominans* in Drakensberg Mountains in South Africa, 353 OTUs were recovered from 11,813 sequences which passed through the denoising steps, and removal of singletons and non-fungal sequences. Ascomycota dominated in the roots of *E. dominans* in most of the samples. We identified two OTUs corresponding to PEA, which represented approx. 2% of sequences in the whole dataset and occurred in 10 out of 15 samples. The rest of OTUs with high similarity to PEA clustered with *Meliniomyces* sp. 2, which belongs to the Clade 4 of PEA according to Hambleton and Sigler (2005). For the first time, we demonstrated the well-known ErMF *P. ericae* in Ericaceae roots in Africa. Although *P. ericae* has been frequently documented in Ericaceae roots from the Northern Hemisphere (e.g., Bougoure et al. 2007; Kjøller et al. 2010; Gorzelak

et al. 2012), it was so far mostly detected in non-Ericaceae plants and rhizoids of leafy liverworts in Antarctica or Chile (Chambers et al. 1999; Upson et al. 2007; Upson et al. 2009). The only record of *P. ericae* in Ericaceae roots from the Southern Hemisphere comes from a recent study in Patagonia (Bruzone et al. 2016). Taken together, *P. ericae* represents the most widespread fungal species (so far not detected only in Australia) with ericoid mycorrhizal lifestyle. *Erica dominans* root fungal communities were structured by geographical distance, altitude as well as microhabitat (dry vs. wet sites). We detected several fungal taxa which distribution was significantly affected by microhabitat type. Only the OTU0005, matching with *Meliniomyces* sp. 2 from *P. ericae* aggregate, was significantly more common in dry sites.

Based on synthesis on 111 independent sampling units from 44 published studies and five unpublished datasets, we did not find any significant difference in response between ErMF and other fungal guilds to host plant identity (VIII). The only exception was represented by OrM fungal communities, which responded most strongly to the identity of their host plants. Orchid and ericoid mycorrhizal fungal communities were also more modular than EcMF and EndF communities, with AMF in an intermediate position. Compared to EcMF symbionts with multiple examples of high unilateral or reciprocal specialization in EcM symbiosis, such as *Alnus* and *Alnus* associated fungi or *Pinus* and Suillaceae, (Bruns et al., 2002; Kennedy et al., 2015), the average host effect on ErMF community composition was comparable, although such examples of specificity are not known for ErMF symbiosis. Interestingly, the relatively low level of host effect in ErM associations contrasts with the high modularity among this group. Such contrasts between modularity and host effect have been reported previously, possibly deriving from the sensitivity of the modularity measure to the total links in the dataset (Bahram et al., 2014).

CONCLUSIONS

The following main conclusions can be inferred from my thesis:

- In the cultured isolates, more than 95% of the functional ErM mycobionts belonged to the Helotiales.
- Novel lineage of basidiomycete formed characteristic sheathed ErM symbiosis with mantle resembling a Hartig net and enhanced growth of *Vaccinium* spp. in vitro, and showed ability to degrade a recalcitrant aromatic substrate.
- Fungi belonging to the *Pezoloma ericae* aggregate co-associate as endophytes with ectomycorrhizas of *Picea abies* in a temperate montane forest in central Europe, where ericoid shrubs dominate in forest understory. Most of the PEA isolates formed dense intracellular hyphal coils resembling ericoid mycorrhiza in roots of *V. myrtillus*. Therefore EcM roots may represent an important refuge for ErMF.
- We found the ability of *Acephala macrosclerotiorum* to form EcM symbiosis with *P. abies* as well as ErM symbiosis with *V. myrtillus* based on resynthesis experiments in laboratory conditions. However, we did not detect any sequences similar to *A. macrosclerotiorum* in Ericaceae roots, although *Acephala macrosclerotiorum* dominated on surrounding EcM roots in natural forests. It remains questionable, if *Acephala macrosclerotiorum* may serve as a common mycorrhizal network between Ericaceae and Pinaceae in temperate or boreal forests.
- The communities of Ericaceae root-associated fungi were affected by environmental factors such as elevation, host plant species as well as soil moisture-related microhabitat, indicating the presence of multi-scale environmental filtering. On the global scale, ericoid mycorrhizal fungi showed strong biogeographical patterns. While some ErM fungal species (*Pezoloma ericae*) have a very broad distribution range, other species have much narrower distribution range restricted to a single hemisphere (*Meliniomyces variabilis*) or continent (*Cairneyella variabilis*). This suggests that also conidial microfungi can be limited by long-distance dispersal capacity and sometimes more than the host plants.

SUMMARY

Mycorrhizal symbiosis is a mutualistic partnership between plants and fungi that represents one of the oldest and the most widespread symbioses on the Earth. One of the youngest mycorrhizal type is ericoid mycorrhizal (ErM) symbiosis, a mutualistic relationship formed between species belonging to several lineages of the Ericaceae family and diverse group of soil fungi. Ericoid mycorrhiza is characterized by the intensive fungal colonization of the outermost root cell layer. Mycorrhizal fungi form a coiled intracellular hyphal complex. Ericoid mycorrhizal plants often occur on extremely poor soils, where most of the nutrients are locked up in complex forms of soil organic matter, with restricted biological availability. The ErM symbiosis represents a key evolutionary adaptation of ErM plants to mobilize the nutrients from such recalcitrant substrates. However, ErM remains largely overlooked compared to the more common mycorrhizal types, such as arbuscular mycorrhiza and ectomycorrhiza, and a broader general understanding of the ErM symbiosis is lacking. While arbuscular mycorrhizal fungi have a monophyletic origin, the ability to form ErM as well as ectomycorrhiza evolved independently multiple times in several fungal lineages. The ERM fungi belong to several fungal lineages of Ascomycota as well as Basidiomycota. Nowadays, there are few sufficiently proven ErM fungi, which belong to four taxonomic groups, Helotiales, Chaetothyriales (Ascomycota) and Sebaciniales and Hymenochaetales (Basidiomycota). In this thesis, I focus on Ericaceae associated fungi from various aspects of the partnership, because the ericoid mycorrhizal symbiosis is probably the most overlooked mycorrhizal symbiotic type. Firstly, this thesis aims to determine the occurrence of ericoid mycorrhizal life-style among fungi. Furthermore, I also focused on determination of environmental factors which significantly affect composition of fungal communities associated with roots of ericoid mycorrhizal plants. To determine the occurrence of ericoid mycorrhizal life-style among fungi, we isolated root associated fungi from sampled Ericaceae plants or seedlings of *Picea abies* from sampling sites in Norway or Czechia. Ecological lifestyle of isolated fungi was assessed in number of resynthesize experiments. To determine the environmental factors, which affect composition of fungal communities associated with roots of ericoid mycorrhizal plants, we sampled roots of Ericaceae plants on Mount Wilhelm in Papua New Guinea and in Drakensberg mountain range in the Republic of South Africa. I used the 454 pyrosequencing method to determine fungal communities associated with the Ericaceae roots.

The main results and conclusions are the following: 1) Novel lineage of Basidiomycota formed characteristic sheathed ErM symbiosis with mantle resembling a Hartig net and enhanced growth of *Vaccinium* spp. in vitro, and showed ability to degrade a recalcitrant aromatic substrate. 2) Most of the *Pezoloma ericae* aggregate isolates associated with *Picea abies* formed dense

intracellular hyphal coils resembling ericoid mycorrhiza in roots of *V. myrtillus*. 3) We found the ability of *Acephala macrosclerotiorum* to form EcM symbiosis with *P. abies* as well as ErM symbiosis with *V. myrtillus* based on resynthesis experiments in laboratory conditions. 4) The communities of Ericaceae root-associated fungi were affected by environmental factors such as elevation, host plant species as well as soil moisture-related microhabitat, indicating the presence of multi-scale environmental filtering. 5) On the global scale, ericoid mycorrhizal fungi showed strong biogeographical patterns. While some ErM fungal species (*Pezoloma ericae*) have a very broad distribution range, other species have much narrower distribution range restricted to a single hemisphere (*Meliniomyces variabilis*) or continent (*Cairneyella variabilis*).

SUMMARY IN ESTONIAN

Erikoidse mükoriisa ökoloogia

Sümbioos mükoriisa ehk seenjuure vahendusel on üks vanimaid ja enim levinud mutualismivorme maailmas. Mükoriissetes suhetes seenpartner varustab taime vees lahustunud mineraalainetega ning saab taimelt vastu suhkruid. Erikoidne mükoriisa (ErM) on arbuskulaarse mükoriisa ja ektomükoriisa kõrval evolutsiooniliselt noorim mükoriisatüüp. Taimed sugukonnast kanarbikulised (Ericaceae), kuhu kuuluvad ka mustikas, pohl ja kanarbik, moodustavad ErM sümbioosi mitmete rühmade mullas ja juurtes elavate endofüütsete seentega. Seenehüüfid kasvavad taime juurerakkudesse sisse ja moodustavad paunakesi – aju-kujulisi struktuure, kus taime rakumembraani ja seeneraku piirpind on oluliselt suurenenud toitainete vahetuse soodustamiseks. Kanarbikulised esinevad sageli vaestel happelistel muldadel, kus toitained on peamiselt orgaanilisel kujul ja raskesti taimedele omastatavad. Kanarbikuliste peamine ökoloogiline kohastumus ongi mükoriisaseente abil makro- ja mikroelementide kättesaamine. Kanarbikulised on metsanduse ja põllumajanduse aspektist väheolulised, mistõttu ErM ökofüsioloogia ja sellega seotud seente elurikkust on vähe uuritud võrreldes teiste mükoriisatüüpidega. Senised elurikkuse uuringud ja mükoriisa sünteesi katse näitavad, et ErM moodustavad seened kuuluvad mitmesse kottseente (*Helotiales*, *Chaetothyriales*) ja kandseente (*Sebacinales*, *Hymenochaetales*) seltsidesse. Oma väitekirjas keskendun erinevatele aspektidele ErM seente elurikkuses ja ökoloogias, eelkõige keskkonnaparametritele, mis mõjutavad seenekooslusi. ErM seente tuvastamiseks isoleerisin kanarbikuliste ja hariliku kuuse juurtes kasvavad seened puhaskultuuri Tsehhiima ja Norra metsadest ning viisin läbi mükoriisa sünteesi katsed laboris. ErM seente kooslusi mõjutavate keskkonnaparametrite tuvastamiseks kogusime kanarbikuliste juuri Mount Wilhelmi mäelt Paapua Uus-Guineas ja Drakensbergi mägedest Lõuna-Aafrika Vabariigis. Kasutasin Roche 454 pürosekvenerimise tehnoloogiat sealt kogutud juurtelt seente määramiseks.

Minu väitekirja peamised tulemused ja järeldused on järgmised: 1) seni määramata kandseente haru moodustab erikoidset mükoriisat, kus taimerakkude pinnal areneb õhuke seeneniidistiku kiht ja seen soodustab mustika kasvu ning lagundab aromaatsaid polümeere; 2) kuusejuurtelt isoleeritud seened *Pezoloma ericae* liigikompleksist on võimelised moodustama erikoidsele mükoriisale iseloomulikke struktuure mustika juurerakkudes; 3) seeneliik *Acephala macrosclerotiorum* moodustab ektomükoriisat hariliku kuusega ja erikoidset mükoriisat mustikaga laboritingimustes; 4) kanarbikuliste seenekooslusi mõjutavad paljud keskkonnaparametrid, eelkõige kõrgus üle merepinna, peremeestaimeliik ning mikroelupaik niiskuse gradiendil; 5) geograafiline kaugus globaalsel skaalal: mõnel seenerühmal on väga lai levik (*Pezoloma ericae*), ent teised on piiratuma levilaga (*Meliniomyces variabilis* ja *Cairneyella variabilis*).

ACKNOWLEDGEMENTS

First of all, I am very grateful to my supervisor, Prof. Leho Tedersoo, who has guided me throughout my first scientific steps and inspired me to learn more about fascinating world of fungal ecology. I am also thankful to Prof. Urmas Kõljalg for his valuable time and advice through these years. I acknowledge all of the co-authors of paper included in this thesis. Particularly, I would like to thank Mohammad Bahram who was always helpful with all bioinformatics and statistics. Besides that I owe gratitude to Sergei, Jane, Heidi, Sten, Rasmus, Kessy, Martin, Tereza and Jesse.

Special thanks to my family for the possibility to make my own life choices and all their support.

Financial support was provided by Estonian Science Foundation grants (9286, PUT0171 and EMP265) and FIBIR. Additional support was supplied by ESF Doctoral Studies and Internationalization Program DoRa.

REFERENCES

- Abarenkov K, Nilsson RH, Larsson K-H, Alexander IJ, Eberhardt U, Erland S, Høiland K, Kjølner R, Larsson E, Pennanen T, et al. (2010a) The UNITE database for molecular identification of fungi—recent updates and future perspectives. *New Phytol* 186: 281–285.
- Abarenkov K, Tedersoo L, Nilsson RH, Vellak K, Saar I, Veldre V, Parmasto E, Proulx M, Aan A, Ots M, et al. (2010b) PlutoF – a web based workbench for ecological and taxonomic research with an online implementation for fungal ITS sequences. *Evol Bioinform* 6: 189–196.
- Abe JP (2005) An arbuscular mycorrhizal genus in the Ericaceae. *Inoculum* 56: 6.
- Agerer R (ed) (1987–2006) *Colour atlas of Ectomycorrhizae*. Einhorn Verlag, Schwäbisch Gmünd.
- Allen TR, Millar T, Berch SM, Berbee ML (2003) Culturing and direct DNA extraction find different fungi from the same ericoid mycorrhizal roots. *New Phytol* 160: 255–272.
- Bahram M, Harend H, Tedersoo L (2014) Network perspectives of ectomycorrhizal associations. *Fung Ecol* 7: 70–77.
- Baldrian P, Kolařík M, Štursová M, Kopecký J, Valášková V, Větrovský T, Žifčáková L, Šnajdr J, Ridl J, et al. (2012) Active and total microbial communities in forest soil are largely different and highly stratified during decomposition. *ISME Journal* 6: 248–258.
- Berch SM, Allen TR, Berbee ML (2002) Molecular detection, community structure and phylogeny of ericoid mycorrhizal fungi. *Plant Soil* 244: 55–66.
- Binder M, Hibbett DS, Larsson KH, Larsson E, Langer E, Langer G (2005) The phylogenetic distribution of resupinate forms across the major clades of mushroom-forming fungi (Homobasidiomycetes). *Syst Biodivers* 3: 113–157.
- Borcard D, Legendre P (2002) All-scale spatial analysis of ecological data by means of principal coordinates of neighbour matrices. *Ecol Model* 153: 51–68.
- Bougoure DS, Cairney JWG (2005a) Fungi associated with hair roots of *Rhododendron lochiaie* (Ericaceae) in an Australian tropical cloud forest revealed by culturing and culture-independent molecular methods. *Environ Microbiol* 7: 1743–1754.
- Bougoure DS, Cairney JWG (2005b) Assemblages of ericoid mycorrhizal and other root-associated fungi from *Epacris pulchella* (Ericaceae) as determined by culturing and direct DNA extraction from roots. *Environ Microbiol* 7: 819–827.
- Bougoure DS, Parkin PI, Cairney JWG, Alexander IJ, Anderson IC (2007) Diversity of fungi in hair roots of Ericaceae varies along a vegetation gradient. *Mol Ecol* 16: 4624–4636.
- Brown SP, Jumpponen A (2014) Contrasting primary successional trajectories of fungi and bacteria in retreating glacier soils. *Mol Ecol* 23: 481–497.
- Brundrett MC (2009) Mycorrhizal associations and other means of nutrition of vascular plants: understanding the global diversity of host plants by resolving conflicting information and developing reliable means of diagnosis. *Plant Soil* 320: 37–77.
- Bruns TD, Bidartondo MI, Taylor DL (2002) Host specificity in ectomycorrhizal communities: what do the exceptions tell us? *Integr Compar Biol* 42: 352–359.
- Bruzzone MC, Fehrer J, Fontenla SB, Vohník M (2017) First record of *Rhizoscyphus ericae* in Southern Hemisphere's Ericaceae. *Mycorrhiza* 27: 147–163.
- Cairney JWG (2000) Evolution of mycorrhiza systems. *Naturwissenschaften* 87: 467–475.

- Carpenter RJ, Macphail MK, Jordan GJ, Hill RS (2015) Fossil evidence for open, Proteaceae dominated heathlands and fire in the Late Cretaceous of Australia. *Am J Bot* 12: 2092–2107.
- Chambers SM, Williams PG, Seppelt RD, Cairney JWG (1999) Molecular identification of *Hymenoscyphus* sp. from rhizoids of the leafy liverwort *Cephaloziella exiliflora* in Australia and Antarctica. *Mycol Res* 103: 286–288.
- Edgar RC, Haas BJ, Clemente JC, Quince C, Knight R (2011) UCHIME improves sensitivity and speed of chimera detection. *Bioinformatics* 27: 2194–2200.
- Gao X, Alvo M, Chen J, Li G (2008) Nonparametric multiple comparison procedures for unbalanced one-way factorial designs. *J Stat Plan Infer* 138: 2574–2591.
- Gorman NR, Starrett MC (2003) Host range of a select isolate of the ericoid mycorrhizal fungus *Hymenoscyphus ericae*. *Hortic Sci* 38: 1163–1166.
- Gorzalak MA, Hambleton S, Massicotte HB (2012) Community structure of ericoid mycorrhizas and root-associated fungi of *Vaccinium membranaceum* across an elevation gradient in the Canadian Rocky Mountains. *Fungal Ecol* 5: 36–45.
- Goslee SC, Urban DL (2007) The ecodist package for dissimilarity-based analysis of ecological data. *J Stat Soft* 22: 1–19.
- Grelet GA, Johnson D, Paterson E, Anderson IC, Alexander IJ (2009) Reciprocal carbon and nitrogen transfer between an ericaceous dwarf shrub and fungi isolated from *Piceirhiza bicolorata* ectomycorrhizas. *New Phytol* 182: 359–366.
- Guimera R, Amaral LAN (2005) Functional cartography of complex metabolic networks. *Nature* 433: 895–900.
- Hall TA (1999) BioEdit: a user-friendly biological sequence alignment editor and analysis program for Windows 95/98/NT. *Nucleic Acids Symposium Series* 41: 95–98.
- Hambleton S, Sigler L (2005) *Meliniomyces*, a new anamorph genus for root-associated fungi with phylogenetic affinities to *Rhizoscyphus ericae* (= *Hymenoscyphus ericae*), Leotiomycetes. *Styd Mycol* 53: 1–27.
- Haug I, Lempe J, Homeier J, Weiss M, Setaro S, Oberwinkler F, Kottke I (2004) *Graffenrieda emarginata* (Melastomataceae) forms mycorrhizas with Glomeromycota and with a member of the *Hymenoscyphus ericae* aggregate in the organic soil of a neotropical mountain rain forest. *Can. J. Botany-Rev. Can. de Botanique* 82: 340–356.
- Hazard C, Gosling P, Mitchell DT, Doohan FM, Bending GD (2014) Diversity of fungi associated with hair roots of ericaceous plants is affected by land use. *FEMS Microbiol Ecol* 87: 586–600.
- Hijmans RJ, Cameron SE, Parra JL, Jones PG, Jarvis A (2005) Very high resolution interpolated climate surfaces for global land areas. *Inter J Climat* 25: 1965–1978.
- Hynson NA, Bruns TD (2009) Evidence of a myco-heterotroph in the plant family Ericaceae that lacks mycorrhizal specificity. *Proc R Soc B* 276: 4053–4059.
- Kennedy PG, Walker JK, Bogar LM (2015) Interspecific mycorrhizal networks and non-networking hosts: exploring the ecology of the host genus *Alnus*. In: Horton TR, ed. *Mycorrhizal networks*. Dordrecht, the Netherlands: Springer 227–254.
- Kerley SJ, Read DJ (1998) The biology of mycorrhiza in the Ericaceae 20. Plant and mycorrhizal necromass as nitrogenous substrates for the ericoid mycorrhizal fungus *Hymenoscyphus ericae* and its host. *New Phytol* 139: 353–360.
- Kernaghan G, Patriquin G (2011) Host associations between fungal root endophytes and boreal trees. *Microbial Ecology* 62: 460–473.

- Kjøller R, Olsrud M, Michelsen A (2010) Co-existing ericaceous plant species in a subarctic mire community share fungal root endophytes. *Fungal Ecol* 3: 205–214.
- Kohler A, Kuo A, Nagy LG, Morin E, Barry KW, Buscot F, Canbäck B, Choi C, Cichocki N, Clum A, et al (2015) Convergent losses of decay mechanisms and rapid turnover of symbiosis genes in mycorrhizal mutualists. *Nat Genet* 47: 410–415.
- Kohout P, Sýkorová Z, Čtvrtlíková M, Rydlová J, Suda J, Vohník M, Sudová R (2012) Surprising spectra of root associated fungi in submerged aquatic plants. *FEMS Microbiol Ecol* 80: 216–235.
- Kolařík M, Vohník M (2018) When the ribosomal DNA does not tell the truth: The case of the taxonomic position of *Kurtia argillacea*, an ericoid mycorrhizal fungus residing among Hymenochaetales. *Fung Biol* 122: 1–18.
- Köljalg U, Nilsson RH, Abarenkov K, Tedersoo L, Taylor AFS, Bahram M, Bates ST, Bruns TD, Bengtsson-Palme J, Callaghan TM, et al. (2013) Towards a unified paradigm for sequence-based identification of fungi. *Mol Ecol* 22: 5271–5277.
- Kron KA, Judd WS, Stevens PF, Crayn D, Anderberg AA, Gadek P, Quinn CJ, Luteyn JL (2002) Phylogenetic classification of Ericaceae: molecular and morphological evidence. *Bot Rev* 68: 335–423.
- Krpata D, Mühlmann O, Kuhnert R, Ladurner H, Göbl F, Peintner U (2007) High diversity of ectomycorrhizal fungi associated with *Arctostaphylos uva-ursi* in subalpine and alpine zones: potential inoculum for afforestation. *For Ecol Manag* 250: 167–175.
- Kumar S, Stecher G, Tamura K (2016) MEGA7: molecular evolutionary genetics analysis version 7.0 for bigger datasets. *Mol Biol Evol* 33: 1870–1874.
- Leake JR, Read DJ (1991) Experiments with ericoid mycorrhiza. *Methods Microbiol* 23: 435–457.
- Midgley DJ, Rosewarne CP, Greenfield P, Li D, Vockler CJ, Hitchcock CJ, Sawyer NA, Brett R, Edwards J, Pitt JI, Tran-Dinh N (2016) Genomic insights into the carbohydrate catabolism of *Cairneyella variabilis* gen. nov. sp. nov., the first reports from a genome of an ericoid mycorrhizal fungus from the Southern Hemisphere. *Mycorrhiza* 23: 345–352.
- Moore RT (1978) Taxonomic significance of septal ultrastructure with particular reference to the jelly fungi. *Mycologia* 70: 1007–1024.
- Münzenberger B, Bubner B, Wöllecke J (2009) The ectomycorrhizal morphotype *Pinirhiza sclerotia* is formed by *Acephala macrosclerotiorum* sp. nov., a close relative of *Phialocephala fortinii*. *Mycorrhiza* 19: 481–492.
- Nixon KC, Crepet WL (1993) Late Cretaceous fossil flowers of ericalean affinity. *Am J Bot* 80: 616–623.
- Nurfadilah S, Swarts ND, Dixon KW, Lambers H, Merritt DJ (2013) Variation in nutrient-acquisition patterns by mycorrhizal fungi of rare and common orchids explains diversification in a global biodiversity hotspot. *Ann Bot* 6: 1233–1241.
- Oberwinkler F, Riess K, Bauer R, Selosse MA, Weiss M, Garnica S, Zuccaro A (2013) Enigmatic Sebaciniales. *Mycol Prog* 12: 1–27.
- Oksanen J, Blanchet FG, Kindt R (2012) Vegan: community ecology package. R package version 2.4–1, <http://veganr-forgerproject.org/>
- Okuda A, Yamato M, Iwase K (2011) The mycorrhiza of *Schizocodon soldanelloides* var. *magnus* (Diapensiaceae) is regarded as ericoid mycorrhiza from its structure and fungal identities. *Mycoscience* 52: 425–430.

- Öpik M, Davison J, Moora M, Zobel M (2014) DNA-based detection and identification of Glomeromycota: the virtual taxonomy of environmental sequences. *Botany* 92: 135–147.
- Pinheiro J, Bates D, DebRoy S, Sarkar D, R Core Team (2016) nlme: linear and non-linear mixed effects models. R package version 3.1–128, <http://CRAN.R-project.org/package=nlme>
- Redecker D, Kodner R, Graham LE (2000) Glomalean fungi from the Ordovician. *Science* 289: 1920–1921.
- Roberts DW (2014) labdsv: ordination and multivariate analysis for ecology. R package version 1.8–0, <http://ecology.msu.montana.edu/labdsv/R/labdsv>
- Ronquist F, Huelsenbeck J (2003) MrBayes 3: Bayesian phylogenetic inference under mixed models. *Bioinformatics* 19: 1572–1574.
- Schloss PD, Westcott SL, Ryabin T, Hall JR, Hartmann M, Hollister EB, Lesniewski RA, Oakley BB, Parks DH, Robinson C (2009) Introducing mothur: open-source, platform-independent, community-supported software for describing and comparing microbial communities. *Appl Environ Microbiol* 75: 7537–7541.
- Schoch CL, Seifert KA, Huhndorf S, Robert V, Spouge JL, Levesque CA, Chen W, Bolchacova E, Voigt K, Crous PW et al. (2012) Nuclear ribosomal internal transcribed spacer (ITS) region as a universal DNA barcode marker for fungi. *Proc Natl Acad Sci USA* 109: 6241–6246.
- Selosse MA, Setaro S, Glatard F, Richard F, Urcelay C, Weiss M (2007) Sebacinaleae are common mycorrhizal associates of Ericaceae. *New Phytol* 174: 864–878.
- Setaro SD, Kron K (2011) Neotropical and North American Vaccinioideae (Ericaceae) share their mycorrhizal Sebacinaleae—an indication for concerted migration? *PLoS Curr* 3: 1227.
- Smith SE, Read DJ (2008) *Mycorrhizal symbiosis*, 3rd edn. Academic, London.
- Štorchová H, Hrdlicková R, Chrtek J, Tetera M, Fitze D, Fehrer J (2000) An improved method of DNA isolation from plants collected in the field and conserved in saturated NaCl/CTAB solution. *Taxon* 49: 79–84.
- Sýkorová Z, Wiemken A, Redecker D (2007) Cooccurring *Gentiana verna* and *Gentiana acaulis* and their neighboring plants in two swiss upper montane meadows harbor distinct arbuscular mycorrhizal fungal communities. *Appl Environ Microbiol* 73: 5426–5434.
- Tedersoo L, May TW, Smith ME (2010) Ectomycorrhizal lifestyle in fungi: global diversity, distribution, and evolution of phylogenetic lineages. *Mycorrhiza* 20: 217–263.
- Tedersoo L, Bahram M, Põlme S, Kõljalg U, Yorou NS, Wijesundera R, Villarreal-Ruiz L, Vasco-Palacios AM, Thu PQ, Suija A et al. (2014a) Global diversity and geography of soil fungi. *Science* 346: 1256688.
- Tedersoo L, Bahram M, Ryberg M, Otsing E, Kõljalg U, Abarenkov K (2014b) Global biogeography of the ectomycorrhizal/sebacina lineage (Fungi, Sebacinaleae) as revealed from comparative phylogenetics analyses. *Mol Ecol* 23: 4168–4183.
- Tejesvi MV, Sauvola T, Pirttilä AM, Ruotsalainen AL (2013) Neighboring *Deschampsia flexuosa* and *Trientalis europaea* harbor contrasting root fungal endophytic communities. *Mycorrhiza* 23: 1–10.
- Upton R, Read DJ, Newsham KK (2007) Widespread association between the ericoid mycorrhizal fungus *Rhizoscyphus ericae* and a leafy liverwort in the maritime and sub-Antarctic. *New Phytol* 176: 460–471.

- Upton R, Newsham KK, Bridge PD, Pearce DA, Read DJ (2009) Taxonomic affinities of dark septate root endophytes of *Colobanthus quitensis* and *Deschampsia antarctica*, the two native Antarctic vascular plant species. *Fungal Ecol* 2: 184–196.
- Usuki F, Abe JP, Kakishima M (2003) Diversity of ericoid mycorrhizal fungi isolated from hair roots of *Rhododendron obtusum* var. *kaempferi* in a Japanese red pine forest. *Mycoscience* 44: 97–10.
- Vandenkoornhuyse P, Husband R, Daniell TJ, Watson IJ, Duck JM, Fitter AH, Young JPW (2002) Arbuscular mycorrhizal community composition associated with two plant species in a grassland ecosystem. *Mol Ecol* 11: 1555–1564.
- Větrovský T, Baldrian P (2013) Analysis of soil fungal communities by amplicon pyrosequencing: current approaches to data analysis and the introduction of the pipeline SEED. *Biol Fertil Soils* 49: 1027–1037.
- Villarreal-Ruiz L, Anderson IC, Alexander IJ (2004) Interaction between an isolate from the *Hymenoscyphus ericae* aggregate and roots of *Pinus* and *Vaccinium*. *New Phytol* 164: 183–192.
- Vohník M, Pánek M, Fehrer J, Selosse MA (2016) Experimental evidence of ericoid mycorrhizal potential within Serendipitaceae (Sebacinales). *Mycorrhiza* 26: 831–346.
- Vrålstad T, Fossheim T, Schumacher T (2000) *Piceirhiza bicolorata* – the ectomycorrhizal expression of the *Hymenoscyphus ericae* aggregate? *New Phytol* 145: 549–563.
- Vrålstad T, Schumacher T, Taylor AFS (2002) Mycorrhizal synthesis between fungal strains of the *Hymenoscyphus ericae* aggregate and potential ectomycorrhizal and ericoid hosts. *New Phytol* 153: 143–152.
- Walker JF, Aldrich-Wolfe L, Riffel A, Barbare H, Simpson NB, Trowbridge J, Jumpponen A (2011) Diverse Helotiales associated with the roots of three species of Arctic Ericaceae provide no evidence for host specificity. *New Phytol* 191: 515–527.
- Wang CJK, Wilcox HE (1985) New species of ectendomycorrhizal and pseudomycorrhizal fungi – *Phialophora finlandia*, *Chloridium paucisporum* and *Phialocephala fortinii*. *Mycologia* 77: 951–958.
- Waterman RJ, Bidartondo MI, Stofberg J, Combs JK, Gebauer G, Savolainen V, Barraclough TG, Pauw A (2011) The effects of above-and belowground mutualisms on orchid speciation and coexistence. *Am Nat* 117: 54–68.
- Weiss M, Selosse MA, Rexer KH, Urban A, Oberwinkler F (2004) Sebacinales a hitherto overlooked cosm of heterobasidiomycetes with a broad mycorrhizal potential. *Mycol Res* 108: 1003–1010.
- Weiss M, Sýkorová Z, Garnica S, Riess K, Martos F, Krause C, Oberwinkler F, Bauer R, Redecker D (2013) Sebacinales everywhere: previously overlooked ubiquitous fungal endophytes. *PLoS One* 6: e16793.
- Williams AF, Chambers SM, Davies PW, McLean CB, Cairney JWG (2004) Molecular investigation of sterile root-associated fungi from *Epacris microphylla* R. Br. (Ericaceae) and other epacrids at alpine, subalpine and coastal heathland sites. *Australas Mycol* 23: 94–104.
- Zhang C, Yin L, Dai S (2009) Diversity of root-associated fungal endophytes in *Rhododendron fortunei* in subtropical forests of China. *Mycorrhiza* 19: 417–423.

PUBLICATIONS

CURRICULUM VITAE

Name: Petr Kohout
Date of Birth: 26.06.1986
Citizenship: Czechia
Address: Institute of Ecology and Earth Sciences, 14a Ravila St, 50411, Tartu, Estonia
Phone: +420 728 228 263
E-mail: petr.kohout@natur.cuni.cz

Education:

1997–2005 Gymnasium Ústavní, Prague, Czechia
2005–2008 Bachelor Degree, Department of Experimental Plant Biology, Faculty of Science, Charles University in Prague, Czechia.
2008–2011 Master Degree, Department of Experimental Plant Biology, Faculty of Science, Charles University in Prague, Czechia. (Dr. Martin Vohník, prof. Jana Albrechtová)
2012– PhD Degree, Department of Botany, University of Tartu, Estonia (Dr. Leho Tedersoo, prof. Urmas Kõljalg)

Publications:

- Kohout P**, Sýkorová Z, Bahram M, Hadincová V, Albrechtová J, Tedersoo L and Vohník M. 2011. Understorey ericaceous shrubs affect ectomycorrhizal fungal community of the invasive *Pinus strobus* and native *Pinus sylvestris* in a pot experiment. *Mycorrhiza* 21: 403–412.
- Tedersoo L, Abarenkov K, Nilsson RH, Schüßler A, Grelet GA, **Kohout P**, Oja J, Bonito GM, Veldre V, Jairus T, Ryberg M, Larsson KH and Kõljalg U. 2011. Tidying up GenBank: quality and metadata annotation of ITS sequences of mycorrhizal fungi. *PLoS ONE* 6: e24904.
- Kohout P**, Sýkorová Z, Čvrtlíková M, Rydlová J, Suda J, Vohník M and Sudová R. 2012. Surprising spectra of root associated fungi in submerged aquatic plants. *FEMS Microbiology Ecology* 80: 216–235.
- Vohník M, Sadowsky JJ, **Kohout P**, Lhotáková Z, Nestby R and Kolařík M. 2012. Novel root-fungus symbiosis in Ericaceae: sheathed ericoid mycorrhiza formed by a hitherto undescribed lineage in Agaricomycetes. *PLoS ONE* 7: e39524.
- Kohout P**, Malinová T, Roy M, Vohník M and Jersáková J. 2013. Diverse fungal community associated with roots of *Pseudorchis albida* (Orchidaceae). *Fungal Ecology* 6: 50–64.
- Bahram M, Kõljalg U, **Kohout P**, Mirshahvaladi S and Tedersoo L. 2013. Ectomycorrhizal fungal community and diversity of exotic pine plantations in relation to native Fagales in Iran: evidence of host switch of local symbionts to distantly related exotic host taxa. *Mycorrhiza* 23: 11–19.

- Doubková P, **Kohout P**, Sudová R. 2013. Soil nutritional status, not inoculum identity primarily determines the effect of different arbuscular mycorrhizal fungi on the growth of *Knautia arvensis* plants. *Mycorrhiza* 23: 561–572.
- Vohník M, Mrnka L, Lukešová T, Bruzone MC, **Kohout P** and Fehrer J. 2013. The cultivable endophytic community of Norway spruce ectomycorrhizae from microhabitats lacking ericaceous hosts is dominated by ericoid mycorrhizal *Meliniomyces variabilis*. *Fungal Ecology* 6: 281–292.
- Kõljalg U, Nilsson RH, Abarenkov A, Tedersoo T, Taylor AFS, Bahram M, Bates ST, Bruns TD, Bengtsson-Palme J, Martin T, Callaghan, Douglas B, Drenkhan T, Eberhardt U, Dueñas M, Grebenc T, Griffith GW, Hartmann M, Kirk PM, **Kohout P**, Larsson E, Lindahl BD, Lücking R, Martín MP, Matheny PB, Nguyen NH, Niskanen T, Oja J, Peay KG, Peintner U, Peterson M, Põldmaa K, Saag L, Saar I, Schüßler A, Senés C, Smith ME, Suija A, Taylor L, Telleria MT, Weiß M and Larsson KH. 2013. Towards a unified paradigm for sequence-based identification of Fungi. *Molecular Ecology* 22: 271–277.
- Kohout P**, Sudová R, Janoušková M, Čtvrtlíková M, Hejda M, Pánková H, Slavíková R, Štajerová K, Vosátka M, Sýkorová Z. 2014. Comparison of commonly used primer sets for evaluation of arbuscular mycorrhizal fungal communities: Is there a universal solution? *Soil Biology and Biochemistry* 68: 482–493.
- Tedersoo L, Bahram M, Põlme S, Kõljalg U, Yorou NS, Wijesundera R, Abell S, Ruiz LV, Thu PQ, Suija A, Smith ME, Sharp S, Saluveer E, Saitta A, Rosas M, Riit T, Ratkowsky D, Pritsch K, Põldmaa K, Piepenbring M, Phosri C, Peterson M, Parts K, Pärtel K, Palacios AV, Otsing E, Nouhra E, Njouonkou AL, Nilsson RH, Mayor J, May TW, Majuakim L, Lee SS, Larsson KH, Kohout P, Hosaka K, Hiiesalu I, Henkel TW, Harend H, Guo L, Greslebin A, Grelet G, Geml J, Gates G, Dunstan W, Dunk C, Drenkhan R, Dearnaley J, Kesel A, Dang T, Chen X, Buegger F, Brearley F, Bonito G, Anslan S, Abarenkov K. 2014. Global diversity and geography of soil fungi. *Science* 346: 1256688.
- Lukešová T, **Kohout P**, Větrovský T, Vohník M. 2015. The potential of Dark Septate Endophytes to form root symbioses with ectomycorrhizal and ericoid mycorrhizal middle European forest plants. *PLoS One* 10: e0124752.
- Kohout P**, Doubková P, Bahram M, Suda J, Tedersoo L, Voříšková J, Sudová R. 2015. Niche partitioning in arbuscular mycorrhizal communities in temperate grasslands: a lesson from adjacent serpentine and nonserpentine habitats. *Molecular Ecology* 24: 1831–1843.
- Oja J, **Kohout P**, Tedersoo L, Kull T, Kõljalg U. 2015. Temporal patterns of orchid mycorrhizal fungi in meadows and forests as revealed by 454 pyrosequencing. *New Phytologist* 205: 1608–1618.
- Põlme S, Öpik M, Moora M, Zobel M, **Kohout P**, Kõljalg U, Oja J, Tedersoo L. 2016. Arbuscular mycorrhizal fungi associating with roots of *Alnus* and *Rubus* in Europe and Middle East. *Fungal Ecology* 24: 27–34.

- Sýkorová Z, Rydlová J, Slavíková R, Ness T, **Kohout P**, Püschel D. 2016. Appraisal of mycorrhizal inoculation in forest reclamation of fly ash deposits. *Restoration Ecology* 24: 184–193.
- Bahram M, **Kohout P**, Anslan S, Harend H, Abarenkov K, Tedersoo L. 2016. Fine-scale spatial patterns of soil biota provide evidence for the predominance of stochastic processes in community assembly. *ISME Journal* 10: 885–896.
- Kohout P**, Bahram M, Põlme S, Tedersoo L. 2017. Altitude, space and host plant biogeography origin structure Ericaceae root associated fungal communities on Papua New Guinea. *Fungal Ecology* 30: 112–121.
- Harantová L, Mudrák O, **Kohout P**, Elhotová D, Frouz J, Baldrian P. 2017. Development of microbial community along primary succession in areas degraded by mining activities. *Land Degradation and Development* 28: 2574–2584.
- Knoblochová T, **Kohout P**, Püschel D, Doubková P, Frouz J, Cajthaml T, Kukla J, Vosátka M, Rydlová J. 2017. Asymmetric response of root-associated fungal communities of an arbuscular mycorrhizal grass and an ectomycorrhizal tree to their coexistence in primary succession. *Mycorrhiza* 27: 775–789.
- Kolaříková Z*, **Kohout P***, Krüger C, Janoušková M, Mrnka L, Rydlová J. 2017. Root-associated fungal communities along a primary successional chronosequence: different ecological guilds assemble differently. *Soil Biology and Biochemistry* 113: 143–152. *Joint first authorship
- Baldrian P, **Kohout P**. 2017. Interactions of saprotrophic fungi with tree roots: Can we observe the emergence of novel ectomycorrhizal fungi? *New Phytologist* 215: 511–513.
- Krüger C*, **Kohout P***, Janoušková M, Püschel D, Frouz J, Rydlová J. 2017. Plant communities rather than soil factors contribute to assembly of arbuscular mycorrhizal fungal communities along primary succession on a mine spoil. *Frontiers in Microbiology* 8: 719 *Joint first authorship
- Kohout P**. 2017. Biogeography of ericoid mycorrhiza. Chapter in *Ecological Studies: Biogeography of Mycorrhizal Symbiosis*, ed. by L. Tedersoo.
- Oja J, Vahta J, Bahram M, **Kohout P**, Kull T, Rannap R, Koljalg U, Tedersoo L. 2017. Local-scale spatial distribution and community composition of orchid mycorrhizal fungi in relation to grazing and environmental effects in semi-natural grasslands. *Mycorrhiza* 27: 355–367.
- Kohout P**, Tedersoo L. 2017. Effect of soil moisture on root-associated fungal communities of *Erica dominans* in Drakensberg mountains in South Africa. *Mycorrhiza* 27: 397–406.
- Janoušková M, **Kohout P**, Moradi J, Doubková P, Frouz J, Vosolsobě S, Rydlová J. 2018. Microarthropods influence the composition of rhizospheric fungal communities by stimulating specific taxa. *Soil Biology and Biochemistry* 122: 120–130.
- Kohout P**, Charvátová M, Štursová M, Mašíňová T, Tomšovský M, Baldrian P. 2018. Clearcutting alters decomposition processes and initiates complex

restructuring of fungal communities in soil and tree roots. *ISME Journal* 12: 692–703.

Põlme S, Bahram M, Jacquemyn H, Kennedy P, **Kohout P**, Moora M, Oja J, Öpik M, Pecoraro L, Tedersoo L. 2018. Host preference and network properties in biotrophic symbiotic plant-fungal associations. *New Phytologist* 217: 1230–1239.

ELULOOKIRJELDUS

Nimi: Petr Kohout
Sünniaeg: 26.06.1986
Kodadondsus: Tšehhi
Address: Ökoloogia ja Maateaduste Instituut, 14a Ravila St, 50411,
Tartu, Eesti
Telefon: +420 728 228 263
E-mail: petr.kohout@natur.cuni.cz

Hariduskäik:

1997–2005 Gymnasium Ústavní, Praha, Tšehhi
2005–2008 B.Sc bioloogia, Department of Experimental Plant Biology,
Faculty of Science, Charles Ülikool, Tšehhi.
2008–2011 M.Sc bioloogia, Department of Experimental Plant Biology,
Faculty of Science, Charles Ülikool, Tšehhi. (Dr. Martin
Vohník, prof. Jana Albrechtová)
2012– Botaanika osakond, Ökoloogia ja Maateaduste Instituut, Tartu
Ülikool, doktorant (Dr. Leho Tedersoo, prof. Urmas Kõljalg)

Publikatsioonid:

- Kohout P**, Sýkorová Z, Bahram M, Hadincová V, Albrechtová J, Tedersoo L and Vohník M. 2011. Understorey ericaceous shrubs affect ectomycorrhizal fungal community of the invasive *Pinus strobus* and native *Pinus sylvestris* in a pot experiment. *Mycorrhiza* 21: 403–412.
- Tedersoo L, Abarenkov K, Nilsson RH, Schüßler A, Grelet GA, **Kohout P**, Oja J, Bonito GM, Veldre V, Jairus T, Ryberg M, Larsson KH and Kõljalg U. 2011. Tidying up GenBank: quality and metadata annotation of ITS sequences of mycorrhizal fungi. *PLoS ONE* 6: e24904.
- Kohout P**, Sýkorová Z, Čtvrtlíková M, Rydlová J, Suda J, Vohník M and Sudová R. 2012. Surprising spectra of root associated fungi in submerged aquatic plants. *FEMS Microbiology Ecology* 80: 216–235.
- Vohník M, Sadowsky JJ, **Kohout P**, Lhotáková Z, Nestby R and Kolařík M. 2012. Novel root-fungus symbiosis in Ericaceae: sheathed ericoid mycorrhiza formed by a hitherto undescribed lineage in Agaricomycetes. *PLoS ONE* 7: e39524.
- Kohout P**, Malinová T, Roy M, Vohník M and Jersáková J. 2013. Diverse fungal community associated with roots of *Pseudorchis albida* (Orchidaceae). *Fungal Ecology* 6: 50–64.
- Bahram M, Kõljalg U, **Kohout P**, Mirshahvaladi S and Tedersoo L. 2013. Ectomycorrhizal fungal community and diversity of exotic pine plantations in relation to native Fagales in Iran: evidence of host switch of local symbionts to distantly related exotic host taxa. *Mycorrhiza* 23: 11–19.

- Doubková P, **Kohout P**, Sudová R. 2013. Soil nutritional status, not inoculum identity primarily determines the effect of different arbuscular mycorrhizal fungi on the growth of *Knautia arvensis* plants. *Mycorrhiza* 23: 561–572.
- Vohník M, Mrnka L, Lukešová T, Bruzone MC, **Kohout P** and Fehrer J. 2013. The cultivable endophytic community of Norway spruce ectomycorrhizae from microhabitats lacking ericaceous hosts is dominated by ericoid mycorrhizal *Meliniomyces variabilis*. *Fungal Ecology* 6: 281–292.
- Kõljalg U, Nilsson RH, Abarenkov A, Tedersoo T, Taylor AFS, Bahram M, Bates ST, Bruns TD, Bengtsson-Palme J, Martin T, Callaghan, Douglas B, Drenkhan T, Eberhardt U, Dueñas M, Grebenc T, Griffith GW, Hartmann M, Kirk PM, **Kohout P**, Larsson E, Lindahl BD, Lücking R, Martín MP, Matheny PB, Nguyen NH, Niskanen T, Oja J, Peay KG, Peintner U, Peterson M, Põldmaa K, Saag L, Saar I, Schüßler A, Senés C, Smith ME, Suija A, Taylor L, Telleria MT, Weiß M and Larsson KH. 2013. Towards a unified paradigm for sequence-based identification of Fungi. *Molecular Ecology* 22: 271–277.
- Kohout P**, Sudová R, Janoušková M, Čtvrtlíková M, Hejda M, Pánková H, Slavíková R, Štajerová K, Vosátka M, Sýkorová Z. 2014. Comparison of commonly used primer sets for evaluation of arbuscular mycorrhizal fungal communities: Is there a universal solution? *Soil Biology and Biochemistry* 68: 482–493.
- Tedersoo L, Bahram M, Põlme S, Kõljalg U, Yorou NS, Wijesundera R, Abell S, Ruiz LV, Thu PQ, Suija A, Smith ME, Sharp S, Saluveer E, Saitta A, Rosas M, Riit T, Ratkowsky D, Pritsch K, Põldmaa K, Piepenbring M, Phosri C, Peterson M, Parts K, Pärtel K, Palacios AV, Otsing E, Nouhra E, Njouonkou AL, Nilsson RH, Mayor J, May TW, Majuakim L, Lee SS, Larsson KH, Kohout P, Hosaka K, Hiiesalu I, Henkel TW, Harend H, Guo L, Greslebin A, Grelet G, Geml J, Gates G, Dunstan W, Dunk C, Drenkhan R, Dearnaley J, Kesel A, Dang T, Chen X, Buegger F, Brearley F, Bonito G, Anslan S, Abarenkov K. 2014. Global diversity and geography of soil fungi. *Science* 346: 1256688.
- Lukešová T, **Kohout P**, Větrovský T, Vohník M. 2015. The potential of Dark Septate Endophytes to form root symbioses with ectomycorrhizal and ericoid mycorrhizal middle European forest plants. *PLoS One* 10: e0124752.
- Kohout P**, Doubková P, Bahram M, Suda J, Tedersoo L, Voříšková J, Sudová R. 2015. Niche partitioning in arbuscular mycorrhizal communities in temperate grasslands: a lesson from adjacent serpentine and nonserpentine habitats. *Molecular Ecology* 24: 1831–1843.
- Oja J, **Kohout P**, Tedersoo L, Kull T, Kõljalg U. 2015. Temporal patterns of orchid mycorrhizal fungi in meadows and forests as revealed by 454 pyrosequencing. *New Phytologist* 205: 1608–1618.
- Põlme S, Öpik M, Moora M, Zobel M, **Kohout P**, Kõljalg U, Oja J, Tedersoo L. 2016. Arbuscular mycorrhizal fungi associating with roots of *Alnus* and *Rubus* in Europe and Middle East. *Fungal Ecology* 24: 27–34.

- Sýkorová Z, Rydlová J, Slavíková R, Ness T, **Kohout P**, Püschel D. 2016. Appraisal of mycorrhizal inoculation in forest reclamation of fly ash deposits. *Restoration Ecology* 24: 184–193.
- Bahram M, **Kohout P**, Anslan S, Harend H, Abarenkov K, Tedersoo L. 2016. Fine-scale spatial patterns of soil biota provide evidence for the predominance of stochastic processes in community assembly. *ISME Journal* 10: 885–896.
- Kohout P**, Bahram M, Pöhlme S, Tedersoo L. 2017. Altitude, space and host plant biogeography origin structure Ericaceae root associated fungal communities on Papua New Guinea. *Fungal Ecology* 30: 112–121.
- Harantová L, Mudrák O, **Kohout P**, Elhotová D, Frouz J, Baldrian P. 2017. Development of microbial community along primary succession in areas degraded by mining activities. *Land Degradation and Development* 28: 2574–2584.
- Knoblochová T, **Kohout P**, Püschel D, Doubková P, Frouz J, Cajthaml T, Kukla J, Vosátka M, Rydlová J. 2017. Asymmetric response of root-associated fungal communities of an arbuscular mycorrhizal grass and an ectomycorrhizal tree to their coexistence in primary succession. *Mycorrhiza* 27: 775–789.
- Kolaříková Z*, **Kohout P***, Krüger C, Janoušková M, Mrnka L, Rydlová J. 2017. Root-associated fungal communities along a primary successional chronosequence: different ecological guilds assemble differently. *Soil Biology and Biochemistry* 113: 143–152. * ühine esimene autorsus
- Baldrian P, **Kohout P**. 2017. Interactions of saprotrophic fungi with tree roots: Can we observe the emergence of novel ectomycorrhizal fungi? *New Phytologist* 215: 511–513.
- Krüger C*, **Kohout P***, Janoušková M, Püschel D, Frouz J, Rydlová J. 2017. Plant communities rather than soil factors contribute to assembly of arbuscular mycorrhizal fungal communities along primary succession on a mine spoil. *Frontiers in Microbiology* 8: 719 * ühine esimene autorsus
- Kohout P**. 2017. Biogeography of ericoid mycorrhiza. Chapter in *Ecological Studies: Biogeography of Mycorrhizal Symbiosis*, ed. by L. Tedersoo.
- Oja J, Vahta J, Bahram M, **Kohout P**, Kull T, Rannap R, Koljalg U, Tedersoo L. 2017. Local-scale spatial distribution and community composition of orchid mycorrhizal fungi in relation to grazing and environmental effects in semi-natural grasslands. *Mycorrhiza* 27: 355–367.
- Kohout P**, Tedersoo L. 2017. Effect of soil moisture on root-associated fungal communities of *Erica dominans* in Drakensberg mountains in South Africa. *Mycorrhiza* 27: 397–406.
- Janoušková M, **Kohout P**, Moradi J, Doubková P, Frouz J, Vosolsobě S, Rydlová J. 2018. Microarthropods influence the composition of rhizospheric fungal communities by stimulating specific taxa. *Soil Biology and Biochemistry* 122: 120–130.
- Kohout P**, Charvátová M, Štursová M, Mašíňová T, Tomšovský M, Baldrian P. 2018. Clearcutting alters decomposition processes and initiates complex

restructuring of fungal communities in soil and tree roots. *ISME Journal* 12: 692–703.

Põlme S, Bahram M, Jacquemyn H, Kennedy P, **Kohout P**, Moora M, Oja J, Öpik M, Pecoraro L, Tedersoo L. 2018. Host preference and network properties in biotrophic symbiotic plant-fungal associations. *New Phytologist* 217: 1230–1239.

DISSERTATIONES BIOLOGICAE UNIVERSITATIS TARTUENSIS

1. **Toivo Maimets.** Studies of human oncoprotein p53. Tartu, 1991, 96 p.
2. **Enn K. Seppet.** Thyroid state control over energy metabolism, ion transport and contractile functions in rat heart. Tartu, 1991, 135 p.
3. **Kristjan Zobel.** Epifüütsete makrosamblike väärtus õhu saastuse indikaatoritena Hamar-Dobani boreaalsetes mägimetsades. Tartu, 1992, 131 lk.
4. **Andres Mäe.** Conjugal mobilization of catabolic plasmids by transposable elements in helper plasmids. Tartu, 1992, 91 p.
5. **Maia Kivisaar.** Studies on phenol degradation genes of *Pseudomonas* sp. strain EST 1001. Tartu, 1992, 61 p.
6. **Allan Nurk.** Nucleotide sequences of phenol degradative genes from *Pseudomonas* sp. strain EST 1001 and their transcriptional activation in *Pseudomonas putida*. Tartu, 1992, 72 p.
7. **Ülo Tamm.** The genus *Populus* L. in Estonia: variation of the species biology and introduction. Tartu, 1993, 91 p.
8. **Jaanus Remme.** Studies on the peptidyltransferase centre of the *E.coli* ribosome. Tartu, 1993, 68 p.
9. **Ülo Langel.** Galanin and galanin antagonists. Tartu, 1993, 97 p.
10. **Arvo Käärnd.** The development of an automatic online dynamic fluorescence-based pH-dependent fiber optic penicillin flowthrough biosensor for the control of the benzylpenicillin hydrolysis. Tartu, 1993, 117 p.
11. **Lilian Järvekülg.** Antigenic analysis and development of sensitive immunoassay for potato viruses. Tartu, 1993, 147 p.
12. **Jaak Palumets.** Analysis of phytomass partition in Norway spruce. Tartu, 1993, 47 p.
13. **Arne Sellin.** Variation in hydraulic architecture of *Picea abies* (L.) Karst. trees grown under different environmental conditions. Tartu, 1994, 119 p.
13. **Mati Reeben.** Regulation of light neurofilament gene expression. Tartu, 1994, 108 p.
14. **Urmas Tartes.** Respiration rhythms in insects. Tartu, 1995, 109 p.
15. **Ülo Puurand.** The complete nucleotide sequence and infections *in vitro* transcripts from cloned cDNA of a potato A potyvirus. Tartu, 1995, 96 p.
16. **Peeter Hõrak.** Pathways of selection in avian reproduction: a functional framework and its application in the population study of the great tit (*Parus major*). Tartu, 1995, 118 p.
17. **Erkki Truve.** Studies on specific and broad spectrum virus resistance in transgenic plants. Tartu, 1996, 158 p.
18. **Illar Pata.** Cloning and characterization of human and mouse ribosomal protein S6-encoding genes. Tartu, 1996, 60 p.
19. **Ülo Niinemets.** Importance of structural features of leaves and canopy in determining species shade-tolerance in temperature deciduous woody taxa. Tartu, 1996, 150 p.

20. **Ants Kurg.** Bovine leukemia virus: molecular studies on the packaging region and DNA diagnostics in cattle. Tartu, 1996, 104 p.
21. **Ene Ustav.** E2 as the modulator of the BPV1 DNA replication. Tartu, 1996, 100 p.
22. **Aksel Soosaar.** Role of helix-loop-helix and nuclear hormone receptor transcription factors in neurogenesis. Tartu, 1996, 109 p.
23. **Maido Remm.** Human papillomavirus type 18: replication, transformation and gene expression. Tartu, 1997, 117 p.
24. **Tiiu Kull.** Population dynamics in *Cypripedium calceolus* L. Tartu, 1997, 124 p.
25. **Kalle Olli.** Evolutionary life-strategies of autotrophic planktonic microorganisms in the Baltic Sea. Tartu, 1997, 180 p.
26. **Meelis Pärtel.** Species diversity and community dynamics in calcareous grassland communities in Western Estonia. Tartu, 1997, 124 p.
27. **Malle Leht.** The Genus *Potentilla* L. in Estonia, Latvia and Lithuania: distribution, morphology and taxonomy. Tartu, 1997, 186 p.
28. **Tanel Tenson.** Ribosomes, peptides and antibiotic resistance. Tartu, 1997, 80 p.
29. **Arvo Tuvikene.** Assessment of inland water pollution using biomarker responses in fish *in vivo* and *in vitro*. Tartu, 1997, 160 p.
30. **Urmas Saarma.** Tuning ribosomal elongation cycle by mutagenesis of 23S rRNA. Tartu, 1997, 134 p.
31. **Henn Ojaveer.** Composition and dynamics of fish stocks in the gulf of Riga ecosystem. Tartu, 1997, 138 p.
32. **Lembi Lõugas.** Post-glacial development of vertebrate fauna in Estonian water bodies. Tartu, 1997, 138 p.
33. **Margus Pooga.** Cell penetrating peptide, transportan, and its predecessors, galanin-based chimeric peptides. Tartu, 1998, 110 p.
34. **Andres Saag.** Evolutionary relationships in some cetrarioid genera (Lichenized Ascomycota). Tartu, 1998, 196 p.
35. **Aivar Liiv.** Ribosomal large subunit assembly *in vivo*. Tartu, 1998, 158 p.
36. **Tatjana Oja.** Isoenzyme diversity and phylogenetic affinities among the eurasian annual bromes (*Bromus* L., Poaceae). Tartu, 1998, 92 p.
37. **Mari Moora.** The influence of arbuscular mycorrhizal (AM) symbiosis on the competition and coexistence of calcareous grassland plant species. Tartu, 1998, 78 p.
38. **Olavi Kurina.** Fungus gnats in Estonia (*Diptera: Bolitophilidae, Keroplattidae, Macroceridae, Ditomyiidae, Diadocidiidae, Mycetophilidae*). Tartu, 1998, 200 p.
39. **Andrus Tasa.** Biological leaching of shales: black shale and oil shale. Tartu, 1998, 98 p.
40. **Arnold Kristjuhan.** Studies on transcriptional activator properties of tumor suppressor protein p53. Tartu, 1998, 86 p.
41. **Sulev Ingerpuu.** Characterization of some human myeloid cell surface and nuclear differentiation antigens. Tartu, 1998, 163 p.

42. **Veljo Kisand.** Responses of planktonic bacteria to the abiotic and biotic factors in the shallow lake Võrtsjärv. Tartu, 1998, 118 p.
43. **Kadri Põldmaa.** Studies in the systematics of hypomyces and allied genera (Hypocreales, Ascomycota). Tartu, 1998, 178 p.
44. **Markus Vetemaa.** Reproduction parameters of fish as indicators in environmental monitoring. Tartu, 1998, 117 p.
45. **Heli Talvik.** Prepatent periods and species composition of different *Oesophagostomum* spp. populations in Estonia and Denmark. Tartu, 1998, 104 p.
46. **Katrin Heinsoo.** Cuticular and stomatal antechamber conductance to water vapour diffusion in *Picea abies* (L.) karst. Tartu, 1999, 133 p.
47. **Tarmo Annilo.** Studies on mammalian ribosomal protein S7. Tartu, 1998, 77 p.
48. **Indrek Ots.** Health state indices of reproducing great tits (*Parus major*): sources of variation and connections with life-history traits. Tartu, 1999, 117 p.
49. **Juan Jose Cantero.** Plant community diversity and habitat relationships in central Argentina grasslands. Tartu, 1999, 161 p.
50. **Rein Kalamees.** Seed bank, seed rain and community regeneration in Estonian calcareous grasslands. Tartu, 1999, 107 p.
51. **Sulev Kõks.** Cholecystokinin (CCK) – induced anxiety in rats: influence of environmental stimuli and involvement of endopioid mechanisms and serotonin. Tartu, 1999, 123 p.
52. **Ebe Sild.** Impact of increasing concentrations of O₃ and CO₂ on wheat, clover and pasture. Tartu, 1999, 123 p.
53. **Ljudmilla Timofejeva.** Electron microscopical analysis of the synaptone-mal complex formation in cereals. Tartu, 1999, 99 p.
54. **Andres Valkna.** Interactions of galanin receptor with ligands and G-proteins: studies with synthetic peptides. Tartu, 1999, 103 p.
55. **Taavi Virro.** Life cycles of planktonic rotifers in lake Peipsi. Tartu, 1999, 101 p.
56. **Ana Rebane.** Mammalian ribosomal protein S3a genes and intron-encoded small nucleolar RNAs U73 and U82. Tartu, 1999, 85 p.
57. **Tiina Tamm.** Cocksfoot mottle virus: the genome organisation and translational strategies. Tartu, 2000, 101 p.
58. **Reet Kurg.** Structure-function relationship of the bovine papilloma virus E2 protein. Tartu, 2000, 89 p.
59. **Toomas Kivisild.** The origins of Southern and Western Eurasian populations: an mtDNA study. Tartu, 2000, 121 p.
60. **Niilo Kaldalu.** Studies of the TOL plasmid transcription factor XylS. Tartu, 2000, 88 p.
61. **Dina Lepik.** Modulation of viral DNA replication by tumor suppressor protein p53. Tartu, 2000, 106 p.

62. **Kai Vellak**. Influence of different factors on the diversity of the bryophyte vegetation in forest and wooded meadow communities. Tartu, 2000, 122 p.
63. **Jonne Kotta**. Impact of eutrophication and biological invasions on the structure and functions of benthic macrofauna. Tartu, 2000, 160 p.
64. **Georg Martin**. Phytobenthic communities of the Gulf of Riga and the inner sea the West-Estonian archipelago. Tartu, 2000, 139 p.
65. **Silvia Sepp**. Morphological and genetical variation of *Alchemilla L.* in Estonia. Tartu, 2000. 124 p.
66. **Jaan Liira**. On the determinants of structure and diversity in herbaceous plant communities. Tartu, 2000, 96 p.
67. **Priit Zingel**. The role of planktonic ciliates in lake ecosystems. Tartu, 2001, 111 p.
68. **Tiit Teder**. Direct and indirect effects in Host-parasitoid interactions: ecological and evolutionary consequences. Tartu, 2001, 122 p.
69. **Hannes Kollist**. Leaf apoplastic ascorbate as ozone scavenger and its transport across the plasma membrane. Tartu, 2001, 80 p.
70. **Reet Marits**. Role of two-component regulator system PehR-PehS and extracellular protease PrtW in virulence of *Erwinia Carotovora* subsp. *Carotovora*. Tartu, 2001, 112 p.
71. **Vallo Tilgar**. Effect of calcium supplementation on reproductive performance of the pied flycatcher *Ficedula hypoleuca* and the great tit *Parus major*, breeding in Northern temperate forests. Tartu, 2002, 126 p.
72. **Rita Hõrak**. Regulation of transposition of transposon Tn4652 in *Pseudomonas putida*. Tartu, 2002, 108 p.
73. **Liina Eek-Piirsoo**. The effect of fertilization, mowing and additional illumination on the structure of a species-rich grassland community. Tartu, 2002, 74 p.
74. **Krõõt Aasamaa**. Shoot hydraulic conductance and stomatal conductance of six temperate deciduous tree species. Tartu, 2002, 110 p.
75. **Nele Ingerpuu**. Bryophyte diversity and vascular plants. Tartu, 2002, 112 p.
76. **Neeme Tõnisson**. Mutation detection by primer extension on oligonucleotide microarrays. Tartu, 2002, 124 p.
77. **Margus Pensa**. Variation in needle retention of Scots pine in relation to leaf morphology, nitrogen conservation and tree age. Tartu, 2003, 110 p.
78. **Asko Lõhmus**. Habitat preferences and quality for birds of prey: from principles to applications. Tartu, 2003, 168 p.
79. **Viljar Jaks**. p53 – a switch in cellular circuit. Tartu, 2003, 160 p.
80. **Jaana Männik**. Characterization and genetic studies of four ATP-binding cassette (ABC) transporters. Tartu, 2003, 140 p.
81. **Marek Sammul**. Competition and coexistence of clonal plants in relation to productivity. Tartu, 2003, 159 p.
82. **Ivar Ilves**. Virus-cell interactions in the replication cycle of bovine papillomavirus type 1. Tartu, 2003, 89 p.

83. **Andres Männik.** Design and characterization of a novel vector system based on the stable replicator of bovine papillomavirus type 1. Tartu, 2003, 109 p.
84. **Ivika Ostonen.** Fine root structure, dynamics and proportion in net primary production of Norway spruce forest ecosystem in relation to site conditions. Tartu, 2003, 158 p.
85. **Gudrun Veldre.** Somatic status of 12–15-year-old Tartu schoolchildren. Tartu, 2003, 199 p.
86. **Ülo Väli.** The greater spotted eagle *Aquila clanga* and the lesser spotted eagle *A. pomarina*: taxonomy, phylogeography and ecology. Tartu, 2004, 159 p.
87. **Aare Abroi.** The determinants for the native activities of the bovine papillomavirus type 1 E2 protein are separable. Tartu, 2004, 135 p.
88. **Tiina Kahre.** Cystic fibrosis in Estonia. Tartu, 2004, 116 p.
89. **Helen Orav-Kotta.** Habitat choice and feeding activity of benthic suspension feeders and mesograzers in the northern Baltic Sea. Tartu, 2004, 117 p.
90. **Maarja Öpik.** Diversity of arbuscular mycorrhizal fungi in the roots of perennial plants and their effect on plant performance. Tartu, 2004, 175 p.
91. **Kadri Tali.** Species structure of *Neotinea ustulata*. Tartu, 2004, 109 p.
92. **Kristiina Tambets.** Towards the understanding of post-glacial spread of human mitochondrial DNA haplogroups in Europe and beyond: a phylogeographic approach. Tartu, 2004, 163 p.
93. **Arvi Jõers.** Regulation of p53-dependent transcription. Tartu, 2004, 103 p.
94. **Lilian Kadaja.** Studies on modulation of the activity of tumor suppressor protein p53. Tartu, 2004, 103 p.
95. **Jaak Truu.** Oil shale industry wastewater: impact on river microbial community and possibilities for bioremediation. Tartu, 2004, 128 p.
96. **Maire Peters.** Natural horizontal transfer of the *pheBA* operon. Tartu, 2004, 105 p.
97. **Ülo Maiväli.** Studies on the structure-function relationship of the bacterial ribosome. Tartu, 2004, 130 p.
98. **Merit Otsus.** Plant community regeneration and species diversity in dry calcareous grasslands. Tartu, 2004, 103 p.
99. **Mikk Heidemaa.** Systematic studies on sawflies of the genera *Dolerus*, *Empria*, and *Caliroa* (Hymenoptera: Tenthredinidae). Tartu, 2004, 167 p.
100. **Ilmar Tõnno.** The impact of nitrogen and phosphorus concentration and N/P ratio on cyanobacterial dominance and N₂ fixation in some Estonian lakes. Tartu, 2004, 111 p.
101. **Lauri Saks.** Immune function, parasites, and carotenoid-based ornaments in greenfinches. Tartu, 2004, 144 p.
102. **Siiri Rootsi.** Human Y-chromosomal variation in European populations. Tartu, 2004, 142 p.
103. **Eve Vedler.** Structure of the 2,4-dichloro-phenoxyacetic acid-degradative plasmid pEST4011. Tartu, 2005. 106 p.

104. **Andres Tover.** Regulation of transcription of the phenol degradation *pheBA* operon in *Pseudomonas putida*. Tartu, 2005, 126 p.
105. **Helen Udras.** Hexose kinases and glucose transport in the yeast *Hansenula polymorpha*. Tartu, 2005, 100 p.
106. **Ave Suija.** Lichens and lichenicolous fungi in Estonia: diversity, distribution patterns, taxonomy. Tartu, 2005, 162 p.
107. **Piret Lõhmus.** Forest lichens and their substrata in Estonia. Tartu, 2005, 162 p.
108. **Inga Lips.** Abiotic factors controlling the cyanobacterial bloom occurrence in the Gulf of Finland. Tartu, 2005, 156 p.
109. **Kaasik, Krista.** Circadian clock genes in mammalian clockwork, metabolism and behaviour. Tartu, 2005, 121 p.
110. **Juhan Javoš.** The effects of experience on host acceptance in ovipositing moths. Tartu, 2005, 112 p.
111. **Tiina Sedman.** Characterization of the yeast *Saccharomyces cerevisiae* mitochondrial DNA helicase Hmi1. Tartu, 2005, 103 p.
112. **Ruth Aguraiuja.** Hawaiian endemic fern lineage *Diellia* (Aspleniaceae): distribution, population structure and ecology. Tartu, 2005, 112 p.
113. **Riho Teras.** Regulation of transcription from the fusion promoters generated by transposition of Tn4652 into the upstream region of *pheBA* operon in *Pseudomonas putida*. Tartu, 2005, 106 p.
114. **Mait Metspalu.** Through the course of prehistory in india: tracing the mtDNA trail. Tartu, 2005, 138 p.
115. **Elin Lõhmussaar.** The comparative patterns of linkage disequilibrium in European populations and its implication for genetic association studies. Tartu, 2006, 124 p.
116. **Priit Kupper.** Hydraulic and environmental limitations to leaf water relations in trees with respect to canopy position. Tartu, 2006, 126 p.
117. **Heili Ilves.** Stress-induced transposition of Tn4652 in *Pseudomonas Putida*. Tartu, 2006, 120 p.
118. **Silja Kuusk.** Biochemical properties of Hmi1p, a DNA helicase from *Saccharomyces cerevisiae* mitochondria. Tartu, 2006, 126 p.
119. **Kersti Püssa.** Forest edges on medium resolution landsat thematic mapper satellite images. Tartu, 2006, 90 p.
120. **Lea Tummeleht.** Physiological condition and immune function in great tits (*Parus major* L.): Sources of variation and trade-offs in relation to growth. Tartu, 2006, 94 p.
121. **Toomas Esperk.** Larval instar as a key element of insect growth schedules. Tartu, 2006, 186 p.
122. **Harri Valdmann.** Lynx (*Lynx lynx*) and wolf (*Canis lupus*) in the Baltic region: Diets, helminth parasites and genetic variation. Tartu, 2006. 102 p.
123. **Priit Jõers.** Studies of the mitochondrial helicase Hmi1p in *Candida albicans* and *Saccharomyces cerevisia*. Tartu, 2006. 113 p.
124. **Kersti Lilleväli.** Gata3 and Gata2 in inner ear development. Tartu, 2007, 123 p.

125. **Kai Rünk.** Comparative ecology of three fern species: *Dryopteris carthusiana* (Vill.) H.P. Fuchs, *D. expansa* (C. Presl) Fraser-Jenkins & Jermy and *D. dilatata* (Hoffm.) A. Gray (Dryopteridaceae). Tartu, 2007, 143 p.
126. **Aveliina Helm.** Formation and persistence of dry grassland diversity: role of human history and landscape structure. Tartu, 2007, 89 p.
127. **Leho Tedersoo.** Ectomycorrhizal fungi: diversity and community structure in Estonia, Seychelles and Australia. Tartu, 2007, 233 p.
128. **Marko Mägi.** The habitat-related variation of reproductive performance of great tits in a deciduous-coniferous forest mosaic: looking for causes and consequences. Tartu, 2007, 135 p.
129. **Valeria Lulla.** Replication strategies and applications of Semliki Forest virus. Tartu, 2007, 109 p.
130. **Ülle Reier.** Estonian threatened vascular plant species: causes of rarity and conservation. Tartu, 2007, 79 p.
131. **Inga Jüriado.** Diversity of lichen species in Estonia: influence of regional and local factors. Tartu, 2007, 171 p.
132. **Tatjana Krama.** Mobbing behaviour in birds: costs and reciprocity based cooperation. Tartu, 2007, 112 p.
133. **Signe Saumaa.** The role of DNA mismatch repair and oxidative DNA damage defense systems in avoidance of stationary phase mutations in *Pseudomonas putida*. Tartu, 2007, 172 p.
134. **Reedik Mägi.** The linkage disequilibrium and the selection of genetic markers for association studies in european populations. Tartu, 2007, 96 p.
135. **Priit Kilgas.** Blood parameters as indicators of physiological condition and skeletal development in great tits (*Parus major*): natural variation and application in the reproductive ecology of birds. Tartu, 2007, 129 p.
136. **Anu Albert.** The role of water salinity in structuring eastern Baltic coastal fish communities. Tartu, 2007, 95 p.
137. **Kärt Padari.** Protein transduction mechanisms of transportans. Tartu, 2008, 128 p.
138. **Siiri-Lii Sandre.** Selective forces on larval colouration in a moth. Tartu, 2008, 125 p.
139. **Ülle Jõgar.** Conservation and restoration of semi-natural floodplain meadows and their rare plant species. Tartu, 2008, 99 p.
140. **Lauri Laanisto.** Macroecological approach in vegetation science: generality of ecological relationships at the global scale. Tartu, 2008, 133 p.
141. **Reidar Andreson.** Methods and software for predicting PCR failure rate in large genomes. Tartu, 2008, 105 p.
142. **Birgot Paavel.** Bio-optical properties of turbid lakes. Tartu, 2008, 175 p.
143. **Kaire Torn.** Distribution and ecology of charophytes in the Baltic Sea. Tartu, 2008, 98 p.
144. **Vladimir Vimberg.** Peptide mediated macrolide resistance. Tartu, 2008, 190 p.
145. **Daima Örd.** Studies on the stress-inducible pseudokinase TRB3, a novel inhibitor of transcription factor ATF4. Tartu, 2008, 108 p.

146. **Lauri Saag.** Taxonomic and ecologic problems in the genus *Lepraria* (*Stereocaulaceae*, lichenised *Ascomycota*). Tartu, 2008, 175 p.
147. **Ulvi Karu.** Antioxidant protection, carotenoids and coccidians in greenfinches – assessment of the costs of immune activation and mechanisms of parasite resistance in a passerine with carotenoid-based ornaments. Tartu, 2008, 124 p.
148. **Jaanus Remm.** Tree-cavities in forests: density, characteristics and occupancy by animals. Tartu, 2008, 128 p.
149. **Epp Moks.** Tapeworm parasites *Echinococcus multilocularis* and *E. granulosus* in Estonia: phylogenetic relationships and occurrence in wild carnivores and ungulates. Tartu, 2008, 82 p.
150. **Eve Eensalu.** Acclimation of stomatal structure and function in tree canopy: effect of light and CO₂ concentration. Tartu, 2008, 108 p.
151. **Janne Pullat.** Design, functionlization and application of an *in situ* synthesized oligonucleotide microarray. Tartu, 2008, 108 p.
152. **Marta Putrinš.** Responses of *Pseudomonas putida* to phenol-induced metabolic and stress signals. Tartu, 2008, 142 p.
153. **Marina Semtšenko.** Plant root behaviour: responses to neighbours and physical obstructions. Tartu, 2008, 106 p.
154. **Marge Starast.** Influence of cultivation techniques on productivity and fruit quality of some *Vaccinium* and *Rubus* taxa. Tartu, 2008, 154 p.
155. **Age Tats.** Sequence motifs influencing the efficiency of translation. Tartu, 2009, 104 p.
156. **Radi Tegova.** The role of specialized DNA polymerases in mutagenesis in *Pseudomonas putida*. Tartu, 2009, 124 p.
157. **Tsipe Aavik.** Plant species richness, composition and functional trait pattern in agricultural landscapes – the role of land use intensity and landscape structure. Tartu, 2009, 112 p.
158. **Kaja Kiiver.** Semliki forest virus based vectors and cell lines for studying the replication and interactions of alphaviruses and hepaciviruses. Tartu, 2009, 104 p.
159. **Meelis Kadaja.** Papillomavirus Replication Machinery Induces Genomic Instability in its Host Cell. Tartu, 2009, 126 p.
160. **Pille Hallast.** Human and chimpanzee Luteinizing hormone/Chorionic Gonadotropin beta (*LHB/CGB*) gene clusters: diversity and divergence of young duplicated genes. Tartu, 2009, 168 p.
161. **Ain Vellak.** Spatial and temporal aspects of plant species conservation. Tartu, 2009, 86 p.
162. **Triinu Remmel.** Body size evolution in insects with different colouration strategies: the role of predation risk. Tartu, 2009, 168 p.
163. **Jaana Salujõe.** Zooplankton as the indicator of ecological quality and fish predation in lake ecosystems. Tartu, 2009, 129 p.
164. **Ele Vahtmäe.** Mapping benthic habitat with remote sensing in optically complex coastal environments. Tartu, 2009, 109 p.

165. **Liisa Metsamaa**. Model-based assessment to improve the use of remote sensing in recognition and quantitative mapping of cyanobacteria. Tartu, 2009, 114 p.
166. **Pille Säälük**. The role of endocytosis in the protein transduction by cell-penetrating peptides. Tartu, 2009, 155 p.
167. **Lauri Peil**. Ribosome assembly factors in *Escherichia coli*. Tartu, 2009, 147 p.
168. **Lea Hallik**. Generality and specificity in light harvesting, carbon gain capacity and shade tolerance among plant functional groups. Tartu, 2009, 99 p.
169. **Mariliis Tark**. Mutagenic potential of DNA damage repair and tolerance mechanisms under starvation stress. Tartu, 2009, 191 p.
170. **Riinu Rannap**. Impacts of habitat loss and restoration on amphibian populations. Tartu, 2009, 117 p.
171. **Maarja Adojaan**. Molecular variation of HIV-1 and the use of this knowledge in vaccine development. Tartu, 2009, 95 p.
172. **Signe Altmäe**. Genomics and transcriptomics of human induced ovarian folliculogenesis. Tartu, 2010, 179 p.
173. **Triin Suvil**. Mycorrhizal fungi of native and introduced trees in the Seychelles Islands. Tartu, 2010, 107 p.
174. **Velda Lauringson**. Role of suspension feeding in a brackish-water coastal sea. Tartu, 2010, 123 p.
175. **Eero Talts**. Photosynthetic cyclic electron transport – measurement and variably proton-coupled mechanism. Tartu, 2010, 121 p.
176. **Mari Nelis**. Genetic structure of the Estonian population and genetic distance from other populations of European descent. Tartu, 2010, 97 p.
177. **Kaarel Krjutškov**. Arrayed Primer Extension-2 as a multiplex PCR-based method for nucleic acid variation analysis: method and applications. Tartu, 2010, 129 p.
178. **Egle Köster**. Morphological and genetical variation within species complexes: *Anthyllis vulneraria* s. l. and *Alchemilla vulgaris* (coll.). Tartu, 2010, 101 p.
179. **Erki Õunap**. Systematic studies on the subfamily Sterrhinae (Lepidoptera: Geometridae). Tartu, 2010, 111 p.
180. **Merike Jõesaar**. Diversity of key catabolic genes at degradation of phenol and *p*-cresol in pseudomonads. Tartu, 2010, 125 p.
181. **Kristjan Herkül**. Effects of physical disturbance and habitat-modifying species on sediment properties and benthic communities in the northern Baltic Sea. Tartu, 2010, 123 p.
182. **Arto Pulk**. Studies on bacterial ribosomes by chemical modification approaches. Tartu, 2010, 161 p.
183. **Maria Põllupüü**. Ecological relations of cladocerans in a brackish-water ecosystem. Tartu, 2010, 126 p.
184. **Toomas Silla**. Study of the segregation mechanism of the Bovine Papillomavirus Type 1. Tartu, 2010, 188 p.

185. **Gyaneshwer Chaubey**. The demographic history of India: A perspective based on genetic evidence. Tartu, 2010, 184 p.
186. **Katrin Kepp**. Genes involved in cardiovascular traits: detection of genetic variation in Estonian and Czech populations. Tartu, 2010, 164 p.
187. **Virve Sõber**. The role of biotic interactions in plant reproductive performance. Tartu, 2010, 92 p.
188. **Kersti Kangro**. The response of phytoplankton community to the changes in nutrient loading. Tartu, 2010, 144 p.
189. **Joachim M. Gerhold**. Replication and Recombination of mitochondrial DNA in Yeast. Tartu, 2010, 120 p.
190. **Helen Tammert**. Ecological role of physiological and phylogenetic diversity in aquatic bacterial communities. Tartu, 2010, 140 p.
191. **Elle Rajandu**. Factors determining plant and lichen species diversity and composition in Estonian *Calamagrostis* and *Hepatica* site type forests. Tartu, 2010, 123 p.
192. **Paula Ann Kivistik**. ColR-ColS signalling system and transposition of Tn4652 in the adaptation of *Pseudomonas putida*. Tartu, 2010, 118 p.
193. **Siim Sõber**. Blood pressure genetics: from candidate genes to genome-wide association studies. Tartu, 2011, 120 p.
194. **Kalle Kipper**. Studies on the role of helix 69 of 23S rRNA in the factor-dependent stages of translation initiation, elongation, and termination. Tartu, 2011, 178 p.
195. **Triinu Siibak**. Effect of antibiotics on ribosome assembly is indirect. Tartu, 2011, 134 p.
196. **Tambet Tõnissoo**. Identification and molecular analysis of the role of guanine nucleotide exchange factor RIC-8 in mouse development and neural function. Tartu, 2011, 110 p.
197. **Helin Räägel**. Multiple faces of cell-penetrating peptides – their intracellular trafficking, stability and endosomal escape during protein transduction. Tartu, 2011, 161 p.
198. **Andres Jaanus**. Phytoplankton in Estonian coastal waters – variability, trends and response to environmental pressures. Tartu, 2011, 157 p.
199. **Tiit Nikopensius**. Genetic predisposition to nonsyndromic orofacial clefts. Tartu, 2011, 152 p.
200. **Signe Värv**. Studies on the mechanisms of RNA polymerase II-dependent transcription elongation. Tartu, 2011, 108 p.
201. **Kristjan Välk**. Gene expression profiling and genome-wide association studies of non-small cell lung cancer. Tartu, 2011, 98 p.
202. **Arno Põllumäe**. Spatio-temporal patterns of native and invasive zooplankton species under changing climate and eutrophication conditions. Tartu, 2011, 153 p.
203. **Egle Tammeleht**. Brown bear (*Ursus arctos*) population structure, demographic processes and variations in diet in northern Eurasia. Tartu, 2011, 143 p.

205. **Teele Jairus**. Species composition and host preference among ectomycorrhizal fungi in Australian and African ecosystems. Tartu, 2011, 106 p.
206. **Kessy Abarenkov**. PlutoF – cloud database and computing services supporting biological research. Tartu, 2011, 125 p.
207. **Marina Grigorova**. Fine-scale genetic variation of follicle-stimulating hormone beta-subunit coding gene (*FSHB*) and its association with reproductive health. Tartu, 2011, 184 p.
208. **Anu Tiitsaar**. The effects of predation risk and habitat history on butterfly communities. Tartu, 2011, 97 p.
209. **Elin Sild**. Oxidative defences in immunoeological context: validation and application of assays for nitric oxide production and oxidative burst in a wild passerine. Tartu, 2011, 105 p.
210. **Irja Saar**. The taxonomy and phylogeny of the genera *Cystoderma* and *Cystodermella* (Agaricales, Fungi). Tartu, 2012, 167 p.
211. **Pauli Saag**. Natural variation in plumage bacterial assemblages in two wild breeding passerines. Tartu, 2012, 113 p.
212. **Aleksei Lulla**. Alphaviral nonstructural protease and its polyprotein substrate: arrangements for the perfect marriage. Tartu, 2012, 143 p.
213. **Mari Järve**. Different genetic perspectives on human history in Europe and the Caucasus: the stories told by uniparental and autosomal markers. Tartu, 2012, 119 p.
214. **Ott Scheler**. The application of tmRNA as a marker molecule in bacterial diagnostics using microarray and biosensor technology. Tartu, 2012, 93 p.
215. **Anna Balikova**. Studies on the functions of tumor-associated mucin-like leukosialin (CD43) in human cancer cells. Tartu, 2012, 129 p.
216. **Triinu Kõressaar**. Improvement of PCR primer design for detection of prokaryotic species. Tartu, 2012, 83 p.
217. **Tuul Sepp**. Hematological health state indices of greenfinches: sources of individual variation and responses to immune system manipulation. Tartu, 2012, 117 p.
218. **Rya Ero**. Modifier view of the bacterial ribosome. Tartu, 2012, 146 p.
219. **Mohammad Bahram**. Biogeography of ectomycorrhizal fungi across different spatial scales. Tartu, 2012, 165 p.
220. **Anneli Lorents**. Overcoming the plasma membrane barrier: uptake of amphipathic cell-penetrating peptides induces influx of calcium ions and downstream responses. Tartu, 2012, 113 p.
221. **Katrin Männik**. Exploring the genomics of cognitive impairment: whole-genome SNP genotyping experience in Estonian patients and general population. Tartu, 2012, 171 p.
222. **Marko Prouš**. Taxonomy and phylogeny of the sawfly genus *Empria* (Hymenoptera, Tenthredinidae). Tartu, 2012, 192 p.
223. **Triinu Visnapuu**. Levansucrases encoded in the genome of *Pseudomonas syringae* pv. tomato DC3000: heterologous expression, biochemical characterization, mutational analysis and spectrum of polymerization products. Tartu, 2012, 160 p.

224. **Nele Tamberg.** Studies on Semliki Forest virus replication and pathogenesis. Tartu, 2012, 109 p.
225. **Tõnu Esko.** Novel applications of SNP array data in the analysis of the genetic structure of Europeans and in genetic association studies. Tartu, 2012, 149 p.
226. **Timo Arula.** Ecology of early life-history stages of herring *Clupea harengus membras* in the northeastern Baltic Sea. Tartu, 2012, 143 p.
227. **Inga Hiiesalu.** Belowground plant diversity and coexistence patterns in grassland ecosystems. Tartu, 2012, 130 p.
228. **Kadri Koorem.** The influence of abiotic and biotic factors on small-scale plant community patterns and regeneration in boreonemoral forest. Tartu, 2012, 114 p.
229. **Liis Andresen.** Regulation of virulence in plant-pathogenic pectobacteria. Tartu, 2012, 122 p.
230. **Kaupo Kohv.** The direct and indirect effects of management on boreal forest structure and field layer vegetation. Tartu, 2012, 124 p.
231. **Mart Jüssi.** Living on an edge: landlocked seals in changing climate. Tartu, 2012, 114 p.
232. **Riina Klais.** Phytoplankton trends in the Baltic Sea. Tartu, 2012, 136 p.
233. **Rauno Veeroja.** Effects of winter weather, population density and timing of reproduction on life-history traits and population dynamics of moose (*Alces alces*) in Estonia. Tartu, 2012, 92 p.
234. **Marju Keis.** Brown bear (*Ursus arctos*) phylogeography in northern Eurasia. Tartu, 2013, 142 p.
235. **Sergei Põlme.** Biogeography and ecology of *alnus*- associated ectomycorrhizal fungi – from regional to global scale. Tartu, 2013, 90 p.
236. **Liis Uusküla.** Placental gene expression in normal and complicated pregnancy. Tartu, 2013, 173 p.
237. **Marko Lõoke.** Studies on DNA replication initiation in *Saccharomyces cerevisiae*. Tartu, 2013, 112 p.
238. **Anne Aan.** Light- and nitrogen-use and biomass allocation along productivity gradients in multilayer plant communities. Tartu, 2013, 127 p.
239. **Heidi Tamm.** Comprehending phylogenetic diversity – case studies in three groups of ascomycetes. Tartu, 2013, 136 p.
240. **Liina Kangur.** High-Pressure Spectroscopy Study of Chromophore-Binding Hydrogen Bonds in Light-Harvesting Complexes of Photosynthetic Bacteria. Tartu, 2013, 150 p.
241. **Margus Leppik.** Substrate specificity of the multisite specific pseudouridine synthase RluD. Tartu, 2013, 111 p.
242. **Lauris Kaplinski.** The application of oligonucleotide hybridization model for PCR and microarray optimization. Tartu, 2013, 103 p.
243. **Merli Pärnoja.** Patterns of macrophyte distribution and productivity in coastal ecosystems: effect of abiotic and biotic forcing. Tartu, 2013, 155 p.
244. **Tõnu Margus.** Distribution and phylogeny of the bacterial translational GTPases and the Mqsr/YgiT regulatory system. Tartu, 2013, 126 p.

245. **Pille Mänd**. Light use capacity and carbon and nitrogen budget of plants: remote assessment and physiological determinants. Tartu, 2013, 128 p.
246. **Mario Plaas**. Animal model of Wolfram Syndrome in mice: behavioural, biochemical and psychopharmacological characterization. Tartu, 2013, 144 p.
247. **Georgi Hudjašov**. Maps of mitochondrial DNA, Y-chromosome and tyrosinase variation in Eurasian and Oceanian populations. Tartu, 2013, 115 p.
248. **Mari Lepik**. Plasticity to light in herbaceous plants and its importance for community structure and diversity. Tartu, 2013, 102 p.
249. **Ede Leppik**. Diversity of lichens in semi-natural habitats of Estonia. Tartu, 2013, 151 p.
250. **Ülle Saks**. Arbuscular mycorrhizal fungal diversity patterns in boreo-nemoral forest ecosystems. Tartu, 2013, 151 p.
251. **Eneli Oitmaa**. Development of arrayed primer extension microarray assays for molecular diagnostic applications. Tartu, 2013, 147 p.
252. **Jekaterina Jutkina**. The horizontal gene pool for aromatics degradation: bacterial catabolic plasmids of the Baltic Sea aquatic system. Tartu, 2013, 121 p.
253. **Helen Vellau**. Reaction norms for size and age at maturity in insects: rules and exceptions. Tartu, 2014, 132 p.
254. **Randel Kreitsberg**. Using biomarkers in assessment of environmental contamination in fish – new perspectives. Tartu, 2014, 107 p.
255. **Krista Takkis**. Changes in plant species richness and population performance in response to habitat loss and fragmentation. Tartu, 2014, 141 p.
256. **Liina Nagirnaja**. Global and fine-scale genetic determinants of recurrent pregnancy loss. Tartu, 2014, 211 p.
257. **Triin Triisberg**. Factors influencing the re-vegetation of abandoned extracted peatlands in Estonia. Tartu, 2014, 133 p.
258. **Villu Soon**. A phylogenetic revision of the *Chrysis ignita* species group (Hymenoptera: Chrysididae) with emphasis on the northern European fauna. Tartu, 2014, 211 p.
259. **Andrei Nikonov**. RNA-Dependent RNA Polymerase Activity as a Basis for the Detection of Positive-Strand RNA Viruses by Vertebrate Host Cells. Tartu, 2014, 207 p.
260. **Eele Õunapuu-Pikas**. Spatio-temporal variability of leaf hydraulic conductance in woody plants: ecophysiological consequences. Tartu, 2014, 135 p.
261. **Marju Männiste**. Physiological ecology of greenfinches: information content of feathers in relation to immune function and behavior. Tartu, 2014, 121 p.
262. **Katre Kets**. Effects of elevated concentrations of CO₂ and O₃ on leaf photosynthetic parameters in *Populus tremuloides*: diurnal, seasonal and inter-annual patterns. Tartu, 2014, 115 p.

263. **Küllil Lokko**. Seasonal and spatial variability of zoopsammon communities in relation to environmental parameters. Tartu, 2014, 129 p.
264. **Olga Žilina**. Chromosomal microarray analysis as diagnostic tool: Estonian experience. Tartu, 2014, 152 p.
265. **Kertu Lõhmus**. Colonisation ecology of forest-dwelling vascular plants and the conservation value of rural manor parks. Tartu, 2014, 111 p.
266. **Anu Aun**. Mitochondria as integral modulators of cellular signaling. Tartu, 2014, 167 p.
267. **Chandana Basu Mallick**. Genetics of adaptive traits and gender-specific demographic processes in South Asian populations. Tartu, 2014, 160 p.
268. **Riin Tamme**. The relationship between small-scale environmental heterogeneity and plant species diversity. Tartu, 2014, 130 p.
269. **Liina Remm**. Impacts of forest drainage on biodiversity and habitat quality: implications for sustainable management and conservation. Tartu, 2015, 126 p.
270. **Tiina Talve**. Genetic diversity and taxonomy within the genus *Rhinanthus*. Tartu, 2015, 106 p.
271. **Mehis Rohtla**. Otolith sclerochronological studies on migrations, spawning habitat preferences and age of freshwater fishes inhabiting the Baltic Sea. Tartu, 2015, 137 p.
272. **Alexey Reshchikov**. The world fauna of the genus *Lathrolestes* (Hymenoptera, Ichneumonidae). Tartu, 2015, 247 p.
273. **Martin Pook**. Studies on artificial and extracellular matrix protein-rich surfaces as regulators of cell growth and differentiation. Tartu, 2015, 142 p.
274. **Mai Kukumägi**. Factors affecting soil respiration and its components in silver birch and Norway spruce stands. Tartu, 2015, 155 p.
275. **Helen Karu**. Development of ecosystems under human activity in the North-East Estonian industrial region: forests on post-mining sites and bogs. Tartu, 2015, 152 p.
276. **Hedi Peterson**. Exploiting high-throughput data for establishing relationships between genes. Tartu, 2015, 186 p.
277. **Priit Adler**. Analysis and visualisation of large scale microarray data, Tartu, 2015, 126 p.
278. **Aigar Niglas**. Effects of environmental factors on gas exchange in deciduous trees: focus on photosynthetic water-use efficiency. Tartu, 2015, 152 p.
279. **Silja Laht**. Classification and identification of conopeptides using profile hidden Markov models and position-specific scoring matrices. Tartu, 2015, 100 p.
280. **Martin Kesler**. Biological characteristics and restoration of Atlantic salmon *Salmo salar* populations in the Rivers of Northern Estonia. Tartu, 2015, 97 p.
281. **Pratyush Kumar Das**. Biochemical perspective on alphaviral nonstructural protein 2: a tale from multiple domains to enzymatic profiling. Tartu, 2015, 205 p.

282. **Priit Palta**. Computational methods for DNA copy number detection. Tartu, 2015, 130 p.
283. **Julia Sidorenko**. Combating DNA damage and maintenance of genome integrity in pseudomonads. Tartu, 2015, 174 p.
284. **Anastasiia Kovtun-Kante**. Charophytes of Estonian inland and coastal waters: distribution and environmental preferences. Tartu, 2015, 97 p.
285. **Ly Lindman**. The ecology of protected butterfly species in Estonia. Tartu, 2015, 171 p.
286. **Jaanis Lodjak**. Association of Insulin-like Growth Factor I and Corticosterone with Nestling Growth and Fledging Success in Wild Passerines. Tartu, 2016, 113 p.
287. **Ann Kraut**. Conservation of Wood-Inhabiting Biodiversity – Semi-Natural Forests as an Opportunity. Tartu, 2016, 141 p.
288. **Tiit Örd**. Functions and regulation of the mammalian pseudokinase TRIB3. Tartu, 2016, 182. p.
289. **Kairi Käiro**. Biological Quality According to Macroinvertebrates in Streams of Estonia (Baltic Ecoregion of Europe): Effects of Human-induced Hydromorphological Changes. Tartu, 2016, 126 p.
290. **Leidi Laurimaa**. *Echinococcus multilocularis* and other zoonotic parasites in Estonian canids. Tartu, 2016, 144 p.
291. **Helerin Margus**. Characterization of cell-penetrating peptide/nucleic acid nanocomplexes and their cell-entry mechanisms. Tartu, 2016, 173 p.
292. **Kadri Runnel**. Fungal targets and tools for forest conservation. Tartu, 2016, 157 p.
293. **Urmo Võsa**. MicroRNAs in disease and health: aberrant regulation in lung cancer and association with genomic variation. Tartu, 2016, 163 p.
294. **Kristina Mäemets-Allas**. Studies on cell growth promoting AKT signaling pathway – a promising anti-cancer drug target. Tartu, 2016, 146 p.
295. **Janeli Viil**. Studies on cellular and molecular mechanisms that drive normal and regenerative processes in the liver and pathological processes in Dupuytren's contracture. Tartu, 2016, 175 p.
296. **Ene Kook**. Genetic diversity and evolution of *Pulmonaria angustifolia* L. and *Myosotis laxa sensu lato* (Boraginaceae). Tartu, 2016, 106 p.
297. **Kadri Peil**. RNA polymerase II-dependent transcription elongation in *Saccharomyces cerevisiae*. Tartu, 2016, 113 p.
298. **Katrin Ruisu**. The role of RIC8A in mouse development and its function in cell-matrix adhesion and actin cytoskeletal organisation. Tartu, 2016, 129 p.
299. **Janely Pae**. Translocation of cell-penetrating peptides across biological membranes and interactions with plasma membrane constituents. Tartu, 2016, 126 p.
300. **Argo Ronk**. Plant diversity patterns across Europe: observed and dark diversity. Tartu, 2016, 153 p.

301. **Kristiina Mark.** Diversification and species delimitation of lichenized fungi in selected groups of the family Parmeliaceae (Ascomycota). Tartu, 2016, 181 p.
302. **Jaak-Albert Metsoja.** Vegetation dynamics in floodplain meadows: influence of mowing and sediment application. Tartu, 2016, 140 p.
303. **Hedvig Tamman.** The GraTA toxin-antitoxin system of *Pseudomonas putida*: regulation and role in stress tolerance. Tartu, 2016, 154 p.
304. **Kadri Pärtel.** Application of ultrastructural and molecular data in the taxonomy of helotialean fungi. Tartu, 2016, 183 p.
305. **Maris Hindrikson.** Grey wolf (*Canis lupus*) populations in Estonia and Europe: genetic diversity, population structure and -processes, and hybridization between wolves and dogs. Tartu, 2016, 121 p.
306. **Polina Degtjarenko.** Impacts of alkaline dust pollution on biodiversity of plants and lichens: from communities to genetic diversity. Tartu, 2016, 126 p.
307. **Liina Pajusalu.** The effect of CO₂ enrichment on net photosynthesis of macrophytes in a brackish water environment. Tartu, 2016, 126 p.
308. **Stoyan Tankov.** Random walks in the stringent response. Tartu, 2016, 94 p.
309. **Liis Leitsalu.** Communicating genomic research results to population-based biobank participants. Tartu, 2016, 158 p.
310. **Richard Meitern.** Redox physiology of wild birds: validation and application of techniques for detecting oxidative stress. Tartu, 2016, 134 p.
311. **Kaie Lokk.** Comparative genome-wide DNA methylation studies of healthy human tissues and non-small cell lung cancer tissue. Tartu, 2016, 127 p.
312. **Mihhail Kurašin.** Processivity of cellulases and chitinases. Tartu, 2017, 132 p.
313. **Carmen Tali.** Scavenger receptors as a target for nucleic acid delivery with peptide vectors. Tartu, 2017, 155 p.
314. **Katarina Oganjan.** Distribution, feeding and habitat of benthic suspension feeders in a shallow coastal sea. Tartu, 2017, 132 p.
315. **Taavi Paal.** Immigration limitation of forest plants into wooded landscape corridors. Tartu, 2017, 145 p.
316. **Kadri Õunap.** The Williams-Beuren syndrome chromosome region protein WBSCR22 is a ribosome biogenesis factor. Tartu, 2017, 135 p.
317. **Riin Tamm.** In-depth analysis of factors affecting variability in thiopurine methyltransferase activity. Tartu, 2017, 170 p.
318. **Keiu Kask.** The role of RIC8A in the development and regulation of mouse nervous system. Tartu, 2017, 184 p.
319. **Tiia Möller.** Mapping and modelling of the spatial distribution of benthic macrovegetation in the NE Baltic Sea with a special focus on the eelgrass *Zostera marina* Linnaeus, 1753. Tartu, 2017, 162 p.
320. **Silva Kasela.** Genetic regulation of gene expression: detection of tissue- and cell type-specific effects. Tartu, 2017, 150 p.

321. **Karmen Süld.** Food habits, parasites and space use of the raccoon dog *Nyctereutes procyonoides*: the role of an alien species as a predator and vector of zoonotic diseases in Estonia. Tartu, 2017, p.
322. **Ragne Oja.** Consequences of supplementary feeding of wild boar – concern for ground-nesting birds and endoparasite infection. Tartu, 2017, 141 p.
323. **Riin Kont.** The acquisition of cellulose chain by a processive cellobiohydrolase. Tartu, 2017, 117 p.
324. **Liis Kasari.** Plant diversity of semi-natural grasslands: drivers, current status and conservation challenges. Tartu, 2017, 141 p.
325. **Sirgi Saar.** Belowground interactions: the roles of plant genetic relatedness, root exudation and soil legacies. Tartu, 2017, 113 p.
326. **Sten Anslan.** Molecular identification of Collembola and their fungal associates. Tartu, 2017, 125 p.
327. **Imre Taal.** Causes of variation in littoral fish communities of the Eastern Baltic Sea: from community structure to individual life histories. Tartu, 2017, 118 p.
328. **Jürgen Jalak.** Dissecting the Mechanism of Enzymatic Degradation of Cellulose Using Low Molecular Weight Model Substrates. Tartu, 2017, 137 p.
329. **Kairi Kiik.** Reproduction and behaviour of the endangered European mink (*Mustela lutreola*) in captivity. Tartu, 2018, 112 p.
330. **Ivan Kuprijanov.** Habitat use and trophic interactions of native and invasive predatory macroinvertebrates in the northern Baltic Sea. Tartu, 2018, 117 p.
331. **Hendrik Meister.** Evolutionary ecology of insect growth: from geographic patterns to biochemical trade-offs. Tartu, 2018, 147 p.
332. **Ilja Gaidutšik.** Irc3 is a mitochondrial branch migration enzyme in *Saccharomyces cerevisiae*. Tartu, 2018, 161 p.
333. **Lena Neuenkamp.** The dynamics of plant and arbuscular mycorrhizal fungal communities in grasslands under changing land use. Tartu, 2018, 241 p.
334. **Laura Kasak.** Genome structural variation modulating the placenta and pregnancy maintenance. Tartu, 2018, 181 p.
335. **Kersti Riibak.** Importance of dispersal limitation in determining dark diversity of plants across spatial scales. Tartu, 2018, 133 p.
336. **Liina Saar.** Dynamics of grassland plant diversity in changing landscapes. Tartu, 2018, 206 p.
337. **Hanna Ainelo.** Fis regulates *Pseudomonas putida* biofilm formation by controlling the expression of *lapA*. Tartu, 2018, 143 p.
338. **Natalia Pervjakova.** Genomic imprinting in complex traits. Tartu, 2018, 176 p.
339. **Andrio Lahesaare.** The role of global regulator Fis in regulating the expression of *lapF* and the hydrophobicity of soil bacterium *Pseudomonas putida*. Tartu, 2018, 124 p.

340. **Märt Roosaare.** *K*-mer based methods for the identification of bacteria and plasmids. Tartu, 2018, 117 p.
341. **Maria Abakumova.** The relationship between competitive behaviour and the frequency and identity of neighbours in temperate grassland plants. Tartu, 2018, 104 p.
342. **Margus Vilbas.** Biotic interactions affecting habitat use of myrmecophilous butterflies in Northern Europe. Tartu, 2018, 142 p.
343. **Liina Kinkar.** Global patterns of genetic diversity and phylogeography of *Echinococcus granulosus* sensu stricto – a tapeworm species of significant public health concern. Tartu, 2018, 147 p.
344. **Teivi Laurimäe.** Taxonomy and genetic diversity of zoonotic tapeworms in the species complex of *Echinococcus granulosus* sensu lato. Tartu, 2018, 143 p.
345. **Tatjana Jatsenko.** Role of translesion DNA polymerases in mutagenesis and DNA damage tolerance in *Pseudomonads*. Tartu, 2018, 216 p.
346. **Katrin Viigand.** Utilization of α -glucosidic sugars by *Ogataea (Hansenula) polymorpha*. Tartu, 2018, 148 p.
347. **Andres Ainelo.** Physiological effects of the *Pseudomonas putida* toxin grat. Tartu, 2018, 146 p.
348. **Killu Timm.** Effects of two genes (DRD4 and SERT) on great tit (*Parus major*) behaviour and reproductive traits. Tartu, 2018, 117 p.