

KADRI ÕUNAP

The Williams-Beuren syndrome  
chromosome region protein WBSCR22  
is a ribosome biogenesis factor





**KADRI ÕUNAP**

The Williams-Beuren syndrome  
chromosome region protein WBSCR22  
is a ribosome biogenesis factor



1632

UNIVERSITY OF TARTU  
Press

Institute of Molecular and Cell Biology, University of Tartu, Estonia

This dissertation was accepted for the commencement of the degree of Doctor of Philosophy (in Molecular Biology) on April 11, 2017 by the Council of the Institute of Molecular and Cell Biology at the University of Tartu.

Supervisors: Reet Kurg, PhD  
Senior Research Fellow in Gene Technology  
University of Tartu, Tartu, Estonia

Ants Kurg, PhD  
Professor of Molecular Biotechnology  
University of Tartu, Tartu, Estonia

Opponent: Pierre-Emmanuel Gleizes, PhD  
Professor of Cell Biology  
University of Toulouse-Paul Sabatier, Toulouse, France

Commencement: Room No. 105, 23B Riia St., Tartu, on May 22, 2017, at 10:15 am.

Publication of this dissertation is granted by the by the Institute of Molecular and Cell Biology, University of Tartu.

ISSN 1024-6479  
ISBN 978-9949-77-417-3 (print)  
ISBN 978-9949-77-418-0 (pdf)

Copyright: Kadri Õunap, 2017

University of Tartu Press  
[www.tyk.ee](http://www.tyk.ee)

# CONTENTS

LIST OF ORIGINAL PUBLICATIONS .....	7
LIST OF ABBREVIATIONS .....	8
1. INTRODUCTION.....	9
2. LITERATURE REVIEW.....	10
2.1 Williams-Beuren syndrome.....	10
2.2. The 7q11.23 duplication syndrome .....	11
2.3. The structure of the Williams-Beuren syndrome chromosomal region .....	11
2.4. The contributions of individual genes of the WBS locus to the clinical outcome.....	14
2.5. The WBSCR22 protein.....	14
2.5.1. The structure and function of WBSCR22 protein .....	14
2.5.2. The expression pattern of WBSCR22.....	15
2.5.3. The functions of non-mammalian WBSCR22 homologs .....	16
2.6. Eukaryotic ribosome biogenesis.....	17
2.6.1. The structure and function of ribosomes .....	17
2.6.2. Factors required for the production of eukaryotic ribosomes..	17
2.6.3. The processing of pre-rRNA in eukaryotes .....	18
2.6.3.1. The organization of human 47S pre-rRNA and its processing into 45S pre-rRNA.....	18
2.6.3.2. 45S pre-rRNA processing within ITS1 separates the biogenesis pathways of small and large ribosomal subunits .....	21
2.4.3.3. The formation of mature 18S rRNA .....	21
2.4.3.4. The formation of mature 28S and 5.8S rRNA .....	22
2.4.3.5. pre-rRNA processing in the eukaryotic model organism budding yeast .....	23
2.7. rRNA modifications .....	25
2.7.1. rRNA pseudouridylation and ribose 2'-O-methylation are carried out by ribonucleoprotein complexes.....	26
2.7.2. Stand-alone enzymes catalyse the formation of rRNA base modifications .....	27
2.7.2.1. 28S/25S rRNA methyltransferases involved in ribosome biogenesis.....	28
2.7.2.2. 18S rRNA methyltransferases in ribosome biogenesis.....	29
2.8. Trm112/TRMT112 is an interaction partner for several methyltransferases .....	32
2.9 Extracellular vesicles as carriers of biological information.....	33

3. AIMS OF THE STUDY.....	34
4. MATERIALS AND METHODS .....	35
5. RESULTS AND DISCUSSION .....	36
5.1. WBSCR22 protein is required for cell growth, ribosome small subunit biogenesis and pre-rRNA processing (Papers I and II).....	36
5.1.1. The level of WBSCR22 protein is reduced in cell lines derived from Williams-Beuren syndrome patients .....	36
5.1.2. WBSCR22 depletion reduces cell growth .....	36
5.1.3. WBSCR22 is required for ribosome small subunit biogenesis and pre-rRNA processing .....	37
5.1.4. WBSCR22 sediments with ribosome 40S subunits.....	39
5.1.5. WBSCR22 is a nuclear protein that can accumulate within nucleoli .....	39
5.2. A principal interaction partner of WBSCR22 in human cells is TRMT112 (Paper II).....	40
5.2.1. WBSCR22 interacts with TRMT112.....	40
5.2.2. TRMT112 regulates the stability of WBSCR22.....	42
5.2.3. WBSCR22 regulates the subcellular localization of TRMT112 .....	43
5.2.4. WBSCR22 mutants defective in TRMT112 binding are less stable than the wild-type protein.....	44
5.2.5. The level of WBSCR22 protein is regulated by the ubiquitin-proteasome pathway.....	46
5.2.6. WBSCR22 complements the growth and ribosome biogenesis defect of budding yeast $\Delta$ <i>Bud23</i> mutant strain .....	47
5.3. Ribosomal proteins and tRNA synthetases are incorporated into MLV Gag induced extracellular vesicles (Paper III).....	48
6. CONCLUSIONS.....	50
SUMMARY IN ESTONIAN .....	51
REFERENCES.....	53
ACKNOWLEDGEMENTS .....	66
PUBLICATIONS.....	67
CURRICULUM VITAE .....	118
ELULOOKIRJELDUS.....	119

## LIST OF ORIGINAL PUBLICATIONS

This dissertation is based on the following original publications, which will be referred to by the corresponding Roman numerals in the text:

- I **Õunap K**, Käsper L, Kurg A, Kurg R (2013) The human WBSR22 protein is involved in the biogenesis of the 40S ribosomal subunits in mammalian cells. PLoS ONE 8(9): e75686.  
doi:10.1371/journal.pone.0075686
- II **Õunap K**, Leetsi L, Matsoo M, Kurg R (2015) The stability of ribosome biogenesis factor WBSR22 is regulated by interaction with TRMT112 via ubiquitin-proteasome pathway. PLoS ONE 10(7): e0133841.  
doi:10.1371/journal.pone.0133841
- III Kurg R, Reinsalu O, Jagur S, **Õunap K**, Võsa L, Kasvandik S, Padari K, Gildemann K, Ustav M (2016). Biochemical and proteomic characterization of retrovirus Gag based microparticles carrying melanoma antigens. Sci. Rep. 6, 29425; doi: 10.1038/srep29425

These articles are reprinted with the permission of the copyright holders.

My contributions to the publications are as follows:

- I Participated in designing the experiments, carried out most of the experiments, analysed the data, prepared the figures and wrote some parts of the manuscript.
- II Participated in experimental design, carried out most of the experiments, analysed the data, prepared the figures and wrote some parts of the manuscript.
- III Performed some of the subcellular localization studies of melanoma antigens, carried out the validation of mass spectrometry results by immunoblotting, contributed in preparing the figures 3 and 5.

## LIST OF ABBREVIATIONS

ac <sup>4</sup> C	N <sup>4</sup> -acetylcytidine
C1QBP	complement C1q binding protein
CTD	C-terminal domain
EGFP	enhanced green fluorescent protein
ETS	external transcribed spacer
EV	extracellular vesicle
FBL	fibrillarin
FC	fibrillar centre
FISH	fluorescence <i>in situ</i> hybridization
ITS	internal transcribed spacer
LCR	low-copy repeats
LSU	large ribosomal subunit
m <sup>6</sup> A	N <sup>6</sup> , N <sup>6</sup> -dimethyladenosine
m <sup>1</sup> A	N <sup>1</sup> -methyladenosine
m <sup>1</sup> acp <sup>3</sup> Ψ	N <sup>1</sup> -methyl-N <sup>3</sup> -(amino-3-carboxypropyl) pseudouridine
m <sup>3</sup> U	N <sup>3</sup> -methyluridine
m <sup>5</sup> C	N <sup>5</sup> -methylcytosine
m <sup>7</sup> G	N <sup>7</sup> -methylguanosine
MLV	murine leukaemia virus
MTD	methyltransferase domain
NAHR	non-allelic homologous recombination
NLS	nuclear localization signal
NOR	nucleolar organizing regions
PTC	peptidyl transferase centre
SAM	S-adenosyl-L-methionine
SILAC	stable isotope labelling by amino acids in the cell culture
snoRNA	small nucleolar RNA
snoRNP	small ribonucleoprotein particle
SSU	small ribosomal subunit
SV40	simian virus 40
SVAS	supravalvular aortic stenosis
TRMT112	tRNA methyltransferase 11–2 homolog ( <i>S. cerevisiae</i> )
tRNA	transfer RNA
UTR	untranslated region
VLP	virus-like particle
WBS	Williams-Beuren syndrome
WBSCR22	Williams-Beuren syndrome chromosome region 22
YBX1	Y-box binding protein 1
Ψ	pseudouridine



# 1. INTRODUCTION

In human cells, almost 300 trans-acting factors are required for the production of ribosomes. Investigating the functions of individual proteins involved in the biogenesis of ribosomes is relevant because one third of these factors are related to genetic diseases and cancer.

Williams-Beuren syndrome is a multisystem developmental disorder caused by the contiguous deletion of 26–28 genes from chromosome region 7q11.23. This genomic disorder is characterized by cardiovascular abnormalities, connective tissue anomalies, a characteristic neurocognitive and behavioural profile, growth delay, and a subtle but distinctive facial dysmorphism. The 7q11.23 chromosome locus has a complex genomic architecture, consisting of a single-copy region surrounded by segmental duplications that predispose the region to chromosomal rearrangements, including deletions, duplications, and inversions. Although the mutational mechanisms that lead to the development of Williams-Beuren syndrome and a reciprocal duplication syndrome have been thoroughly investigated, it is not known how alterations to the copy number of individual genes contribute to the disease phenotypes. It is likely that several genes contribute to the phenotype and therefore it is important to study the cellular functions of all proteins expressed from the region 7q11.23. The gene that encodes the WBSCR22 protein is deleted in WBS, and studying the function of WBSCR22 is also relevant for cancer biology. Several works have demonstrated that WBSCR22 is upregulated in various types of cancers and regulates the survival and metastatic potential of cancer cells.

The WBSCR22 protein contains an S-adenosyl-L-methionine binding motif that is present in many methyltransferases, and, prior this work, *in silico* predictions indicated that WBSCR22 is a DNA methyltransferase. Later, WBSCR22 was suggested to be involved in histone methylation; only recently, it was confirmed to act as an 18S rRNA base methyltransferase. The first part of this dissertation is focused on studying the function of the WBSCR22 protein in mammalian cells, and, revealed that WBSCR22 is important for cell growth, ribosome biogenesis and pre-rRNA processing. A cell line that stably expresses epitope-tagged WBSCR22 was generated with the aim to study the interaction partners of WBSCR22, revealing that WBSCR22 interacts with several proteins, including the TRMT112. Subsequently, the importance of the interaction between WBSCR22 and TRMT112 proteins was analysed in detail.

The second part of this dissertation is focused on analysing the protein content of virus-like particles (VLPs) induced by the expression of murine leukaemia virus (MLV) Gag protein. The results revealed that these VLPs consist of different groups of cellular proteins, including ribosomal proteins from both subunits.

## 2. LITERATURE REVIEW

### 2.1. Williams-Beuren syndrome

Williams-Beuren syndrome (WBS, also referred to as Williams syndrome; OMIM 194050) is a genomic disorder caused by hemizygous deletion of 26–28 genes on chromosome region 7q11.23 (Pober, 2010). The contiguous gene deletion typically ranges from 1.55 to 1.84 Mb in size, but smaller and larger atypical deletions have also been described (Bayés et al., 2003; Ferrero et al., 2010; Schubert, 2009). The prevalence of WBS is relatively rare and has been estimated to occur sporadically in 1/7,500 to 1/20,000 live births (Strømme et al., 2002).

WBS is a multisystem disorder: patients usually have a subtle but distinctive facial dysmorphism, cardiovascular abnormalities, connective tissue anomalies, a characteristic neurocognitive and behavioural profile, and growth delay (Merla et al., 2010). The syndrome is named after two doctors, Dr. Williams and Dr. Beuren, who independently characterized a syndrome with specific facial features, supravalvular aortic stenosis (SVAS) and mental retardation in the 1960s (Beuren et al., 1962; Williams et al., 1961). The main facial features of WBS are a broad forehead, periorbital fullness, a low nasal root, a long philtrum, a wide mouth and full lips (Merla et al., 2010; Popowski et al., 2011).

The prevalence of cardiovascular diseases is high (~80%) in people diagnosed with WBS (Eronen et al., 2002; Pober et al., 2008). The most common cardiovascular symptoms include vascular stenosis, such as SVAS or pulmonary artery stenosis, and hypertension, however, patients often have multiple problems and a lifelong assessment of the cardiovascular system is recommended (Lin et al., 2008). The cardiovascular phenotype of WBS is similar to that of familial SVAS, and both are caused by haploinsufficiency of the elastin gene (*ELN*), which leads to reduced elastin synthesis (Ewart et al., 1993). Individuals with WBS often have endocrine abnormalities related to glucose metabolism, hypothyroidism, infantile hypercalcemia, hypercalciuria and decreased growth spurt (Pober, 2010).

WBS causes a special neurocognitive and behavioural profile that is characterized by developmental delay and mild to moderate intellectual disability with IQ scores typically in the range of 50 to 60 (Merla et al., 2010; Pober, 2010). The language development of people with WBS is below age expectations, however, compared to other groups that display intellectual disabilities, children with WBS have a higher expressive vocabulary level, albeit similar receptive vocabulary levels (Van Den Heuvel et al., 2016). The characteristic pattern of cognitive strengths and weaknesses of WBS include hypersociability, friendliness, and empathy towards others, but also the presence of anxieties, phobias, and a visuospatial construction deficit (Merla et al., 2010; Pober, 2010). Individuals with WBS often have strong attraction to sounds and music but on the other hand, they can be hypersensitive to certain noises like fireworks or thunderstorms (Gothelf et al., 2006; Pober, 2010). It has been shown that high

frequency sensorineural hearing loss has developed in ~75% of WBS adults over the age of thirty (Cherniske et al., 2004).

In addition to clinical observation, a diagnosis of WBS is confirmed either by fluorescence in situ hybridisation analysis (FISH), using the elastin gene as a marker, or by other molecular biology methods, e.g., restriction fragment length polymorphism, microsatellite analysis or array comparative genomic hybridization (Merla et al., 2010). WBS has also been diagnosed prenatally, following an initial diagnosis of intra-uterine growth retardation and nasal bone hypoplasia (Popowski et al., 2011).

## **2.2. The 7q11.23 duplication syndrome**

In addition to deletions of the WBS chromosomal region, a 1.5 Mb duplication in the corresponding region has also been characterized, and causes 7q11.23 duplication syndrome (also referred to as Williams-Beuren region duplication syndrome; OMIM 609757) (Somerville et al., 2005). Compared to WBS, the phenotype of 7q11.23 duplication syndrome is less severe and the syndrome may be underdiagnosed (Merla et al., 2010). The milder phenotype is probably the reason why the first case of 7q11.23 duplication was documented much later (Somerville et al., 2005). Some features of the duplication syndrome are opposite to WBS – patients with 7q11.23 duplication do not have impaired visuospatial cognitive skills but have deficits in social interactions and very poor expressive language (Merla et al., 2010). Although it is not yet fully defined, the phenotype of the WBS region duplication syndrome is also characterized by subtle but recognizable facial features, developmental delay and an increase in the prevalence of autistic-like features (Pober, 2010; Velleman and Mervis, 2011). Abbas et al. reported that ~91% of individuals diagnosed with WBS locus duplication have speech delay and ~75% have at least one anxiety disorder (Abbas et al., 2016). Among the psychiatric features, social anxiety, attention deficit/hyperactivity disorder, selective mutism, autism, and special phobias are the most prominent (Abbas et al., 2016). The diagnosis of WBS duplication syndrome often occurs as the result of an untargeted genomic analysis method such as chromosomal microarray analysis on patients with autistic-like features.


## **2.3. The structure of the Williams-Beuren syndrome chromosomal region**

The structure of 7q11.23, the WBS chromosomal region, is complex (**Fig. 1A**), and consists of a single copy region (~1.2 Mb in size) that is flanked by segmental duplications or low copy repeats (LCRs) (Bayés et al., 2003). Three large LCR sequences in 7q11.23, the centromeric, middle, and telomeric LCRs (ordered by their location in the chromosomal band), are composed of DNA

blocks termed A, B, and C (Bayés et al., 2003; Schubert, 2009). The DNA blocks have a length of ~10–400 bp and they display ~97% sequence identity within each block class, which predisposes to strand misalignment and unequal crossing-over in meiosis, causing deletions, duplications, and inversions (Stankiewicz and Lupski, 2002).

The centromeric and middle LCR blocks in the WBS chromosomal share the same orientation; however, the telomeric LCRs occur in the opposite direction compared with the centromeric and middle LCRs, which leads to the possibility of deletion, duplication, or inversion events in the WBS chromosomal region (**Fig. 1C and D**) (Bayés et al., 2003). The B-cen and B-mid LCRs share the highest degree of sequence identity (99.6%) and therefore form the most frequent site of meiotic non-allelic homologous recombination (NAHR), thereby generating the 1.55 Mb deletion seen in ~95% of WBS patients (Bayés et al., 2003). NAHR within A-cen and A-mid, generates a 1.84 Mb deletion, which occurs in ~3–5% of WBS patients, and the frequency of atypical deletions are even lower (~2–3% of WBS cases) (Bayés et al., 2003; Merla et al., 2010; Schubert, 2009). In addition to unequal meiotic crossing-over between homologous chromosomes (Baumer et al., 1998; Cuscó et al., 2008; Urbán et al., 1996), WBS deletions can also occur as a result of intrachromatid recombination (**Fig. 1D**) (Bayés et al., 2003; Cuscó et al., 2008).

Duplications of the WBS chromosomal region arise by interchromosomal NAHR, but not by intrachromosomal rearrangements (**Fig. 1C**) (Schubert, 2009). In addition, NAHR can cause inversions (**Fig. 1E**) that range between 1.79 to 2.56 Mb, but these genomic rearrangements do not appear to affect the expression of genes from the WBS region (Bayés et al., 2003; Tam et al., 2008). Inversions can occur in meiosis or mitosis due to intrachromatid misalignment (Schubert, 2009).



**Figure 1.** A schematic representation of the Williams-Beuren syndrome chromosome region (7q11.23) and chromosomal rearrangements. A) The Williams-Beuren chromosome region consists of a 1.2 Mb single copy gene region (blue) and highly similar low-copy repeat (LCR) blocks A (grey), B (red), and C (black). Lower case letters (c-centromeric, m-medial, t-telomeric) mark the positions of LCRs within the chromosomal band. B) Protein-coding genes in the single copy gene region, in medial block C (Cm) and B (Bm) duplicons. These genes are commonly deleted in WBS patients. C) Development of deletion and duplication due to interchromosomal or interchromatid NAHR between directly oriented LCRs. Here we present the most frequent deletion and duplications with breakpoints in the centromeric and medial B-blocks. D) Intrachromatid NAHR leads to the deletion and generation of an acentric chromosome, the latter of which can be lost in segregation. E) Development of inversions in the WBS region caused by NAHR of inverted LCR blocks. Misalignment can occur between centromeric and telomeric blocks of type C, A, and B and therefore the size of the inverted region is variable.

## **2.4. The contributions of individual genes of the WBS locus to the clinical outcome**

Although the molecular mechanisms that lead to the development of genomic rearrangements in 7q11.23 have been known for many years, the role of individual genes to the WBS phenotype has not yet been solved. Genomic rearrangements in the WBS region affect the copy number of up to 28 genes (Merla et al., 2010; Pober, 2010). Quantitative PCR analysis from skin fibroblasts and lymphoblastoid cell lines of WBS patients demonstrate that the hemizygous deletion of 7q11.23 leads to a reduced expression of genes located within the deleted interval (Merla et al., 2006). Furthermore, the mRNA level of some genes that flank the deleted region is also decreased in these patients, showing that the number of genes that can influence the development of WBS phenotype is more numerous than previously expected (Merla et al., 2006). Deletion of elastin gene (ELN) is known to cause cardiovascular problems such as SVAS (Ewart et al., 1993), however, determining the role of many other genes requires further study.

## **2.5. The WBSCR22 protein**

### **2.5.1. The structure and function of WBSCR22 protein**

One of the protein coding genes that is mapped within single copy gene regions of the WBS locus is *WBSCR22* (Doll and Grzeschik, 2001; Merla et al., 2002). *WBSCR22* mRNA encodes a protein composed of 281 amino acid residues with a predicted molecular weight of 31.8 kDa and an isoelectric point of 8.95 (Doll and Grzeschik, 2001). The protein is conserved in eukaryotes (White et al., 2008). *In silico* analysis has shown that the *WBSCR22* protein contains a carboxy-terminal nuclear localization signal (NLS) and an amino-terminal S-adenosyl-L-methionine (SAM) binding motif that is common for methyltransferases (Merla et al., 2002). The human genome encodes 208 methyltransferases that methylate nucleic acids, proteins, lipids, or small molecules, and the *WBSCR22* protein belongs to the largest class of methyltransferases – seven- $\beta$ -strand superfamily or “class I” methyltransferases (Petrossian and Clarke, 2011). The core of these enzymes is composed of seven-stranded  $\beta$ -sheets, where the first six  $\beta$ -sheets are parallel and the seventh occurs in the opposite direction (Falnes et al., 2016; Petrossian and Clarke, 2011). The  $\beta$ -sheets (ordered 3214576) are flanked by  $\alpha$ -helices ( $\alpha$ Z and  $\alpha$ A- $\alpha$ E) – three helices (Z, A, B) are on one side and three helices (C, D and E) on the other (Martin and McMillan, 2002). The yeast orthologue of human *WBSCR22* is the Bud23 protein (Ebersberger et al., 2014). A crystal structure analysis of the methyltransferase domain of budding yeast Bud23 protein revealed that it contains a typical class I methyltransferase fold (Létoquart et al., 2014). The 50% sequence similarity of *WBSCR22* with yeast Bud23 leads us to believe that the structure of *WBSCR22* might be similar to the Bud23 protein.

Despite the presence of the methyltransferase fold, the enzymatic activity of WBSCR22 was unknown for a long time. Based on sequence alignment, it was first suggested to be a DNA methyltransferase, but so far, no DNA methyltransferase activity has been verified experimentally (Doll and Grzeschik, 2001). Several studies have analysed whether WBSCR22 might be a histone methyltransferase (Jangani et al., 2014; Nakazawa et al., 2011). The results of a comparative transcriptome analysis demonstrated a negative correlation between WBSCR22 and tumor-suppressor *Zac1* expression, which suggested that WBSCR22 might be a protein methyltransferase that renders transcriptionally repressive chromatin environment (Nakazawa et al., 2011). Chromatin-immunoprecipitation results indeed indicated the loss of H3-Lys<sup>9</sup> methylation in the *Zac1* promoter after the depletion of WBSCR22; however, an *in vitro* methylation assay could not confirm the histone methyltransferase activity of WBSCR22 (Nakazawa et al., 2011). In addition to regulating *Zac1* expression, downregulation of WBSCR22 expression influences the glucocorticoid receptor target gene *GILZ1* by decreasing its response to glucocorticoid receptor induction (Jangani et al., 2014). WBSCR22 affects glucocorticoid receptor binding to its response elements and furthermore, the intact methyltransferase domain and S-adenosyl-methionine binding sequence of WBSCR22 is necessary to support glucocorticoid receptor transactivation (Jangani et al., 2014). Although the downregulation of WBSCR22 causes changes in the status of histone methylation in the *GILZ1* promoter, the authors of the latter article could not prove that WBSCR22 possess histone methyltransferase activity (Jangani et al., 2014).

We and others have shown that WBSCR22 is involved in ribosome biogenesis and pre-rRNA processing (Paper I; Tafforeau et al., 2013). Recently, the methyltransferase activity of WBSCR22 and its molecular target were confirmed. As with yeast *Bud23*, WBSCR22 carries out ribosomal RNA methylation, namely the methylation of G1639 of 18S rRNA and is one of the few rRNA base methyltransferases in humans (Haag et al., 2015; Zorbas et al., 2015).

### **2.5.2. The expression pattern of WBSCR22**

This gene is composed of twelve exons and spans ~45 kbp on genomic DNA (Doll and Grzeschik, 2001; Merla et al., 2002). Using a panel consisting of twenty human cDNA pools from various tissues and developmental stages, Merla et al detected high expression of WBSCR22 mRNA in kidney, spleen, liver, lung, testis, and fetal heart (Merla et al., 2002). Another study, performed by Doll and Grzeschik, analyzed eight human tissues and showed that WBSCR22 mRNA was detected in all samples tested with high expression in skeletal muscle, heart, and kidney (Doll and Grzeschik, 2001). An analysis of human gene expression array data demonstrated that WBSCR22 is widely or ubiquitously expressed with increased expression in bronchial epithelium and activated immune cells – e.g., in B lymphocytes and CD8<sup>+</sup> T lymphocytes (Jangani et al.,

2014). Due to the observation that WBSCR22 expression is upregulated in primary plasma cells, Tiedemann et al. suggested that it might have a role in plasma cell biology (Tiedemann et al., 2012).

The expression of WBSCR22 is upregulated in different types of cancer – in invasive breast cancer (invasive ductal carcinoma), hepatocellular carcinoma and multiple myeloma (Nakazawa et al., 2011; Stefanska et al., 2014; Tiedemann et al., 2012). Using a mouse experimental metastasis model, it has been shown that both mouse *Wbscr22* and human WBSCR22 proteins are important for metastasis formation by regulating tumour cell survival in the mouse vasculature (Nakazawa et al., 2011). The metastatic potential of *Wbscr22* depends on its expression level – overexpression of *Wbscr22* enhances, and down-regulation by shRNAs reduces the metastatic ability of cancer cells (Nakazawa et al., 2011). Because the expression level of WBSCR22 seems to affect the oncogenic potential of the cell, its expression level might be tightly controlled. The importance of WBSCR22 in cancer biology was further confirmed by the results of a high throughput siRNA screen that targeted about one third of the human genome. This study showed that WBSCR22 is among 57 multiple myeloma survival genes (Tiedemann et al., 2012). Although the downregulation of WBSCR22 expression affects the survival of non-myeloma cells, as in the 293 and A549 cell-lines, the absence of WBSCR22 seems to be more detrimental to multiple myeloma cells (Tiedemann et al., 2012). The relevance of WBSCR22 for cell survival is supported by recent screens that employed CRISPR/Cas9 genome editing and gene trap methods to show that WBSCR22 is among the genes that are essential in human cells (Blomen et al., 2015; Wang et al., 2015a).

### **2.5.3. The functions of non-mammalian WBSCR22 homologs**

In the beginning of this work, the cellular function of the WBSCR22 protein was unknown. However, as already discussed, WBSCR22 protein is highly conserved in eukaryotes and the functions of some of its orthologs are better characterized. The budding yeast protein Bud23 is not essential, however, the deletion of *Bud23* gene causes a slow growth phenotype (Niewmierzycka and Clarke, 1999). *Saccharomyces cerevisiae* has a bipolar budding pattern which means that it can divide either from its distal or proximal polar end. However, a *bud23* deletion mutant has a random budding pattern, meaning that Bud23 is important for bud-site selection (Ni and Snyder, 2001). In addition, Bud23 is known to be important for ribosome biogenesis, 18S pre-rRNA processing and for the formation of m7G1575 of 18S rRNA (White et al., 2008). Similarly, the Rid2 protein in *Arabidopsis thaliana*'s is important for cell proliferation (Ohbayashi et al., 2011). Rid2 was first characterized from a genetic screen that analysed temperature-sensitive mutants defective in root formation (Konishi and Sugiyama, 2003). The Rid2 protein is involved in rRNA processing while its mutation leads to the enlargement of nucleoli and the accumulation of various



pre-rRNA processing intermediates (Ohbayashi et al., 2011). Although RID2 and Bud23 are both involved in rRNA processing, expression of RID2 protein does not correct the slow growth phenotype of *bud23* mutant (Ohbayashi et al., 2011).

## **2.6. Eukaryotic ribosome biogenesis**

### **2.6.1. The structure and function of ribosomes**

Ribosomes are large ribonucleoprotein complexes essential for translating the genetic information encoded within mRNAs into proteins. These particles are composed of ribosomal RNA (rRNA) and proteins (r-proteins): the RNA component has catalytic activity and therefore the ribosomes are actually ribozymes. The eukaryotic and prokaryotic ribosomes differ in size – eukaryotes have more complex 80S ribosomes, whereas prokaryotes have smaller 70S ribosomes. Each ribosome is composed of two unequal subunits, the large subunit (LSU) and the small subunit (SSU) – e.g., 60S and 40S subunits in eukaryotes and 50S and 30S subunits in prokaryotes. The small subunit contains a decoding centre that monitors the complementarity between the anticodons in tRNA and the codons in mRNA. The large subunit contains a peptidyl-transferase centre that catalyses the formation of peptide bonds. Each ribosome contains aminoacyl-tRNA, peptidyl-tRNA, and deacylated tRNA binding regions termed ribosome A-, P- and E-sites, respectively. In contrast with prokaryotic ribosomes, eukaryotic ribosomes contain additional r-proteins and additional residues in both protein and rRNA content, termed ribosomal protein extensions and rRNA expansion segments, respectively (Wilson and Doudna Cate, 2012). The structural core of ribosomes is highly conserved and eukaryote specific rRNA expansion segments (5 in SSU and 16 in LSU) and ribosomal protein extensions are mostly localized to the surface of ribosomal subunits (Wilson and Doudna Cate, 2012). The deletion experiments conducted in budding yeast indicate that eukaryotic expansion segments affect ribosome biogenesis and are important for growth (Ramesh and Woolford, 2016).

The human 40S subunit has one rRNA (18S rRNA) and 33 proteins, whereas the 60S subunit has three rRNAs (28S, 5.8S and 5S rRNA) and 47 proteins (Anger et al., 2013). The yeast 60S subunit has 25S rRNA and 46 ribosomal proteins, meaning that the total number of proteins in the yeast ribosomes is 79 (Yusupova and Yusupov, 2014).

### **2.6.2. Factors required for the production of eukaryotic ribosomes**

Eukaryotic ribosome biogenesis is a complex and energy demanding process that begins in the nucleolus, proceeds in the nucleoplasm, and is completed in the cytoplasm (Woolford and Baserga, 2013). Eukaryotic ribosomes are subject to more complex regulation and biogenesis pathways compared with prokaryotes and moreover, major differences occur in ribosome biogenesis of higher

and lower eukaryotes (Henras et al., 2015; Woolford and Baserga, 2013). Ribosome biogenesis involves the synthesis of pre-rRNAs and ribosomal proteins, pre-rRNA processing, generation of covalent rRNA modifications – e.g., pseudouridylations and methylations, the assembly and transport of pre-ribosomal particles, as well as quality control and surveillance mechanisms (Sharma and Lafontaine, 2015). The production of yeast ribosomes requires more than 200 synthesis factors and ~80 small nucleolar RNAs (snoRNAs) that transiently associate with pre-ribosomes at certain stages during the maturation process (Henras et al., 2015; Mullineux and Lafontaine, 2012; Woolford and Baserga, 2013). In human cells, more than 200 snoRNAs guide rRNA modifications and pre-rRNA processing (Lestrade and Weber, 2006) and it has been shown by a siRNA screen that at least 286 trans-acting factors are important for pre-rRNA processing (Tafforeau et al., 2013). Understanding the role of individual ribosome biogenesis transacting factors is relevant because about one third of the factors are related to genetic diseases and/or cancer (Tafforeau et al., 2013).

### **2.6.3. The processing of pre-rRNA in eukaryotes**

#### **2.6.3.1. The organization of human 47S pre-rRNA and its processing into 45S pre-rRNA**


In human cells, the rDNA that encodes 47S pre-rRNA is tandemly repeated within nucleolar organizing regions (NOR) on five acrocentric chromosomes (chromosomes 13–15, 21 and 22) and the rDNA repeats that encode 5S rRNA are localized on chromosome 1 (Henderson et al., 1973; Sørensen and Frederiksen, 1991). The 47S pre-rRNA is a single polycistronic transcript composed of three rRNAs (18S, 5.8S and 28S) and it is synthesised by RNA polymerase I (Pol I) in the nucleolus. In *Saccharomyces cerevisiae*, the RNA Pol I synthesizes the 35S pre-rRNA with a general architecture similar to mammalian 47S pre-rRNA (Henras et al., 2015; Mullineux and Lafontaine, 2012).

The nucleolus is a non-membranous nuclear compartment that forms around the NOR and it is the major site for ribosome biogenesis (Raska, 2003). The cells of most eukaryotes, including plants, invertebrates, and anamniote vertebrates contain bipartite nucleoli; whereas the nucleoli of amniotic reptiles, birds, and mammals consist of three parts (Thiry and Lafontaine, 2005; Thiry et al., 2011). In the inner part of tripartite nucleolus is the electron-microscope transparent area, fibrillar centre (FC), that contains transcriptionally active rDNA, transcription factors and RNA Pol I (Derenzini et al., 2006). rRNA transcription occurs in the interface between the FC and the dense fibrillar centre, which consists of an electron-dense area that contains nascent rRNA transcripts and ribonucleoprotein complexes that participate in pre-rRNA processing and rRNA modification (Denissov et al., 2011). Maturing ribosomes, which consist of rRNA, ribosomal proteins and transacting factors, move towards the nuclear periphery to the granular component of the nucleolus (Boisvert et al., 2007).

5S rRNA, is transcribed separately by RNA polymerase III in the nucleoplasm; however, it forms a 5S RNP complex with ribosomal proteins RPL5 and RPL11, followed by binding to the pre-60S particles in the nucleolus (Ciganda and Williams, 2011; Pelava et al., 2016). In eukaryotes, both rRNAs are processed by nucleases, and, in *S. cerevisiae*, the exonucleases Rex1p, Rex2p, and Rex3p participate in the processing of the 3' end of 5S rRNA (van Hoof et al., 2000).

The overall architecture of the precursor rRNA processing pathway is conserved through evolution (Henras et al., 2015; Mullineux and Lafontaine, 2012). In a series of exo- and endonucleolytic cleavages, the external (5'ETS and 3'ETS) and internal transcribed spacers (ITS1 and ITS2) are removed from the primary transcript and the mature 18S, 5.8S and 28S/25S rRNAs are released (Eichler and Craig, 1994; Henras et al., 2015; Mullineux and Lafontaine, 2012). The schemes for pre-rRNA processing in human and budding yeast are provided in Figures 2 and 3, respectively. Pre-rRNA processing in yeast can occur co-transcriptionally and usually starts with cleavages within 5'ETS prior to ITS1 processing (Kos and Tollervy, 2010; Mougey et al., 1993). In higher eukaryotes, pre-rRNA processing can follow alternative pathways and there is no evidence of co-transcriptional release of pre-40S particles (Henras et al., 2015).

The processing of human 47S rRNA transcripts begins within 5'ETS at site 01, mapped to the conserved region in close proximity to the box C/D snoRNA U3 binding region (Kass et al., 1987; Mullineux and Lafontaine, 2012). The U3 snoRNA is required for the ribosome small subunit biogenesis and 5'ETS processing at sites 01 and A0 in humans (Langhendries et al., 2016) but also for the initial processing events in mouse (Kass et al., 1990) and yeast (Hughes and Ares, 1991). The human 01 cleavage can occur within two alternative sites, either within C414-C416 or G420-U422 (nucleotide numbering corresponds to the primary transcript) (Kass et al., 1987). The other processing event that occurs shortly after transcription, occurs within 3'ETS at site 02, thereby producing 45S pre-rRNA (Mullineux and Lafontaine, 2012). The human 02 site is located in close proximity to the 3' end of 28S rRNA, and 02 processing is thought to happen either concomitantly with 01 cleavage or soon after 01 cleavage (Mullineux and Lafontaine, 2012).



**Figure 2.** Pre-rRNA processing in human cells. 47S pre-rRNA, transcribed by RNA polymerase I in the nucleolus, is composed of mature 18S (green), 5.8S (grey) and 28S (blue) rRNA. External (5'ETS and 3'ETS) and internal transcribed spacers (ITS1 and ITS2) flank the sequences of mature rRNAs and are removed by exo- and endoribonucleolytic cleavage sites. Individual cleavage sites are marked with blue. The 47S pre-rRNA is cleaved at 5'ETS (site 01) and 3'ETS (site 02), to produce 45S pre-rRNA that can be cleaved by alternative pathways, depending if the next cleavage occurs within ITS1 at site 2 (major pathway) or within 5'ETS at sites A0 and 1 (minor pathway). The cleavage at site 2 produces 30S and 32S pre-rRNA and the further processing of 30S within 5'ETS produces the 21S pre-rRNA. In the minor pathway, the cleavage within 5'ETS produces 41S pre-rRNA that is further cleaved at site 2 to produce 21S and 32S pre-rRNA. The cleavages at A0 and 1 are tightly coupled; however, their uncoupling can produce 26S and 43S pre-rRNA (not marked on scheme). The 21S pre-rRNA is cleaved at sites C and E to produce 18S-E and at site 3 to produce mature 18S rRNA. The 32S pre-rRNA is cleaved at site 3' to separate 12S rRNA and 28S rRNA. The 5'-3' and 3'-5' exoribonucleases are involved in the production of 5.8S<sub>SL</sub> rRNA and 5'-3' exoribonucleases and are required to produce the 5' end of mature 28S rRNA. There are two types of 5.8S rRNA in human cells, 5.8<sub>S</sub> and 5.8<sub>L</sub>, that differing in size ~7–8 nucleotides. The 5' extension of 5.8<sub>L</sub> is marked with pink.

### **2.6.3.2. 45S pre-rRNA processing within ITS1 separates the biogenesis pathways of small and large ribosomal subunits**

In mammalian cells, the processing of 45S pre-rRNA is flexible and is performed by alternative pathways depending if the processing at site O2 is followed by the cleavage within ITS1 or 5'ETS region (Hadjiolova et al., 1993). In the major pathway of HeLa cells, pre-45S rRNA cleavage occurs within ITS1 at site 2 (C6469-6476), thereby producing 30S and 32S pre-rRNAs (Idol et al., 2007; Mullineux and Lafontaine, 2012). The 30S pre-rRNA that contains mature 18S rRNA is further cleaved at sites A0 (producing 26S rRNA) and at site 1 (producing 21S pre-rRNA) (Mullineux and Lafontaine, 2012). In the minor pathway, 45S rRNA cleavage first occurs within 5'ETS, at site A0 (producing 43S pre-rRNA) and at site 1 (producing 41S pre-rRNA) (Mullineux and Lafontaine, 2012). The 5'ETS processing site A0, which is mapped to position G1643, is located between sites O1 and 1 (Rouquette et al., 2005). It has been shown that processing at sites A0 and 1 is tightly coupled and therefore the 26S and 43S pre-rRNA are present at very low amounts under normal conditions (Rouquette et al., 2005). However, the uncoupling of cleavages at sites A0 and 1 cause the accumulation of 43S pre-rRNA in an aggressive breast cancer cell line (Belin et al., 2009). The cleavage of 41S rRNA within ITS1 at site 2 produces 21S and 32S pre-rRNAs; the same processing intermediates were produced in the major pathway and therefore, the downstream processing of 21S and 32S pre-rRNA are identical in both pathways (Mullineux and Lafontaine, 2012). The primary transcript contains rRNAs of both ribosomal subunits; however, the cleavage in ITS1 leads to the separation of the rRNA processing pathways of the small and large ribosomal subunits.

### **2.4.3.3. The formation of mature 18S rRNA**

The processing of the 21S pre-rRNA continues with 3'-5' exoribonucleolytic digestion that stops at a conserved region, termed region C (Henras et al., 2015). Removal of ~250 nucleotides that separate site 2 and site C, yields 21S-C rRNA intermediates (Henras et al., 2015). The 21S-C pre-rRNA accumulates after the knock-down of Rps19 (Idol et al., 2007) and ENP1/bystin (Carron et al., 2011), and it is also detected in normal HeLa cells, albeit at a very low level (Carron et al., 2011).

In the nucleolus, further processing of 21S pre-rRNA yields the 18S-E pre-rRNA that is exported to the cytoplasm where the final maturation of 18S rRNA occurs (Rouquette et al., 2005). Processing of 18S-E pre-rRNA is a multistep process in which the first cleavage occurs at site E, located 79 and 82 nucleotides downstream of the 3' end of 18S rRNA (at G5606 and G5609 of the 47S pre-rRNA) (Preti et al., 2013). Cleavage at site E is followed by rapid 3'-5' exoribonucleolytic trimming (Preti et al., 2013). The exosome complex, namely the EXOSC10/RRP6, exosome cofactors SKIV2L2 and MPP6, exosome core protein RRP46/EXOSC5 and MTR4 helicase, participate in the

production of the 3' end of 18S rRNA (Sloan et al., 2013; Tafforeau et al., 2013). Under normal conditions, ~70-80% of the 18S-E rRNA is found in the cytoplasm (Rouquette et al., 2005). The exonucleolytic processing of 18S-E rRNA occurs during its transport and, as a result, 18S-E pre-rRNA of various lengths can be detected in the nucleus and cytoplasm (Preti et al., 2013). The longest 18S-E rRNA variants contain more than 40 nucleotides of ITS1 and are predominantly localized in the nucleolus, whereas the shorter forms that contain between 14–24 nucleotides of ITS1 are found in the cytoplasm (Preti et al., 2013). The presence of two pools of 18S-E pre-rRNA that differ in size by ~25 nucleotides has also shown by others (Haag et al., 2015).

The final maturation of 18S rRNA occurs in the cytoplasm and requires the cleavage at site 3, which is performed by the conserved endonuclease NOB1 (Preti et al., 2013; Sloan et al., 2013). A recent article provides the three-dimensional structure of human pre-40S particles at 19Å resolution and notes the locations of ribosome biogenesis factors LTV1, ENP1/bystin, TSR1, RIO2, and NOB1 (Larburu et al., 2016). In this model, NOB1 is separated from the 18S rRNA 3' end, and, in order to act at site 3, NOB1 should either change its position or undergo a change in conformation (Larburu et al., 2016). Alternatively, it has been suggested that mature 18S rRNA can be produced by 3'-5' exoribonucleolytic trimming of 18S-E to site 3 (Henras et al., 2015).

#### **2.4.3.4. The formation of mature 28S and 5.8S rRNA**

The 5' end of 32S pre-rRNA, which consists of ITS1, 5.8S rRNA, ITS2 and 28S rRNA, is further processed by exonucleolytic digestion (Mullineux and Lafontaine, 2012). In mouse and human cells, the 5'-3' exonuclease XRN2, which is an ortholog of the Rat1 protein in budding yeast (Schillewaert et al., 2012), is required for the production of the 5' end of 5.8S pre-rRNA (Wang and Pestov, 2011). As in yeast, the 5' end of human 5.8S rRNA is heterogeneous as seen by the presence of shorter (5.8S<sub>S</sub> rRNA) and longer forms (5.8S<sub>L</sub> rRNA) that differ in size by ~7–8 nucleotides (Heindl and Martinez, 2010; Morello et al., 2011). The ratio of 5.8S<sub>S</sub> and 5.8S<sub>L</sub> is between 60:40 to 70:30 in HeLa cells (Heindl and Martinez, 2010; Schillewaert et al., 2012) and does not change after XRN2 depletion (Schillewaert et al., 2012).


Endonucleolytic cleavage of ITS2 at site 3' (C7849 on the 47S rRNA) separates the 12S and 28S rRNA (Hadjiolova et al., 1993) and requires the Las1L protein in humans (Schillewaert et al., 2012). 12S pre-rRNA is further cleaved within ITS2 at site 4a (C6947 on the 47S rRNA), thereby producing 7S pre-rRNA (Farrar et al., 2008). The processing of the 3' end of human 5.8S rRNA requires the exosome complex and is performed via 3'-5' exoribonucleolytic trimming (Schillewaert et al., 2012; Tafforeau et al., 2013). The accumulation of 7S pre-rRNA (~340 nucleotides in size) appear after the depletion of exosome components, including EXOSC2, EXOSC5, EXOSC10, SKIV2L2, and MPHOSPH6 (Tafforeau et al., 2013). However, depletion of EXOSC10 also causes the accumulation of another 5.8S rRNA intermediate, the 5.8S+40

(~190–200 nucleotides in size) (Schillewaert et al., 2012; Tafforeau et al., 2013). The final precursor of 5.8S rRNA is transported to the cytoplasm, where ERI1 is required to remove the last few nucleotides of ITS2 by 3′-5′ exoribonucleolytic digestion (Ansel et al., 2008; Schillewaert et al., 2012).

The 3′ end of 28S rRNA is generated in an early processing event, namely the 02 cleavage of 47S rRNA (Mullineux and Lafontaine, 2012). The vertebrate specific snoRNA U8 is important for ribosome 60S subunit biogenesis and its depletion leads to the accumulation of 3′ extended processing products (45S-L, 43S-L, 41S-L, 36S-L and 32S-L) (Langhendries et al., 2016). As discussed above, the production of the 5′ end of 28S rRNA requires the processing of ITS2 by the 5′-3′ exonuclease XRN2 (Wang and Pestov, 2011). In addition to producing mature rRNA ends, XRN2 is involved in the degradation of 5′ETS, ITS1, and ITS2 spacer fragments (Prete et al., 2013; Schillewaert et al., 2012). As with other rRNAs in mammals, the last intermediate of 28S rRNA, which is extended by 4–6 nucleotides at its 5′ end, is transported to the cytoplasm, where its final maturation occurs (Mullineux and Lafontaine, 2012).

#### **2.4.3.5. pre-rRNA processing in the eukaryotic model organism budding yeast**

For decades, *Saccharomyces cerevisiae* has been the model organism in the field of eukaryotic ribosome biogenesis. This makes it important to introduce the main points gained from studies of budding yeast and compare these with human cells. The 35S pre-rRNA of budding yeast consists of 18S, 5.8S, and 25S rRNAs that are interspersed with external and internal transcribed spacers; however, compared to the human 47S pre-rRNA, the spacers of 35S pre-rRNA, are smaller and the processing of 35S pre-rRNA generates the precursors that are similar, but not identical, to those of higher eukaryotes (**Fig. 3**) (Henras et al., 2015; Woolford and Baserga, 2013). In budding yeast, ~70% of nascent transcripts are cleaved co-transcriptionally at the early processing sites A<sub>0</sub>, A<sub>1</sub>, and A<sub>2</sub>, thereby generating 20S and 27SA<sub>2</sub> pre-rRNAs while remaining pre-rRNA transcripts, which are processed post-transcriptionally within the 90S pre-ribosome complex (Kos and Tollervy, 2010). Another study demonstrates that the frequency of co-transcriptional processing ranges from 41% to 79% and is dependent on the growth phase of budding yeast (Osheim et al., 2004). Co-transcriptional processing requires the formation of the SSU processome, which consists of several processing factors including the core U3 snoRNP complex that base pairs via U3 snoRNA with sequences of 5′ETS and 18S rRNA (Dragon et al., 2002). These complexes are visible as terminal knobs at the 5′ ends of nascent pre-rRNA transcripts using cryo-electron microscopy (Miller and Beatty, 1969; Mougey et al., 1993; Osheim et al., 2004). The 5′ terminal knobs on nascent transcripts appear in most eukaryotes, however, co-transcriptional separation of the precursors of the small and large subunits of the ribosome is rare (Osheim et al., 2004).



**Figure 3.** Schematic representation of pre-rRNA processing in *Saccharomyces cerevisiae*. The 35S pre-rRNA consists of 18S (red), 5.8S (grey), and 25S (blue) rRNAs, flanked by external (5'ETS and 3'ETS) and internal transcribed spacers (ITS1 and ITS2). The processing sites are marked. Pre-rRNA processing can occur co-transcriptionally or after transcription. During post-transcriptional processing, cleavage in 3'ETS at site B<sub>0</sub> generates the 35S pre-rRNA that is processed by coupled digestions at A<sub>0</sub>, A<sub>1</sub>, and A<sub>2</sub>, yielding 20S and 27SA<sub>2</sub> pre-rRNAs. The processing of 20S pre-rRNA continues with nuclear export and cleavage at site D, thereby generating mature 18S rRNA. There are two alternative routes for 27SA<sub>2</sub> processing, either at site A<sub>3</sub> followed by 5'-3' exonucleolytic digestion to site B<sub>1s</sub>, or at B<sub>1L</sub>, yielding 27SB<sub>s</sub> or 27SB<sub>L</sub>, respectively. 27SB<sub>s</sub> and 27SB<sub>L</sub> differ in their 5' ends; however, their exonucleolytic digestions in ITS2 lead to the production of 25S rRNA and 5.8S<sub>s</sub> or 5.8S<sub>L</sub> rRNA, respectively. The 5' extension present in 5.8S<sub>L</sub> is marked with pink.

During post-transcriptional processing, the cleavage in 3'ETS at site B<sub>0</sub> is performed by Rnt1 and releases the 90S pre-ribosome (Kufel et al., 1999; Venema and Tollervey, 1999). The initial cleavage is followed by cleavages at site A<sub>0</sub> of 5'ETS and at site A<sub>1</sub>, generating the 33S and 32S pre-rRNAs, respectively (Venema and Tollervey, 1999). The A<sub>1</sub> cleavage generates the 5' end of mature 18S rRNA and the cleavage at site A<sub>2</sub> in ITS1 results in the separation of the 32S pre-rRNA into the 20S and 27SA<sub>2</sub> pre-rRNA molecules, which are the precursors of SSU and LSU (Henras et al., 2015; Mullineux and Lafontaine, 2012; Woolford and Baserga, 2013). Processing at sites A<sub>0</sub>, A<sub>1</sub>, and A<sub>2</sub> is tightly coupled; however, their uncoupling or inhibition leads to cleavage



at site A<sub>3</sub>, thereby yielding aberrant 23S, 22S, and 21S pre-rRNAs that are probably degraded by the nucleolar surveillance pathway (Mullineux and Lafontaine, 2012). However, 23S pre-rRNA is detected in low amounts in wild type cells and can be a normal rRNA precursor (Granneman and Baserga, 2004).

Unlike the processing of 21S pre-rRNA in higher eukaryotes, which require both exo- and endonucleolytic cleavages, the processing of 20S pre-rRNA in budding yeast requires a single cleavage at site D, which is performed by the endonuclease Nob1, releasing ITS1 fragment D-A<sub>2</sub>, which is degraded by the 5'-3' exoribonuclease Xrn1 (Mullineux and Lafontaine, 2012). Site D corresponds to site 3 in humans and in both organisms, the final processing step is performed in the cytoplasm by Nob1/NOB1 (Larburu et al., 2016; Lebaron et al., 2012; Pertschy et al., 2009; Preti et al., 2013; Sloan et al., 2013).

The processing of 27SA<sub>2</sub> pre-rRNA can occur in two alternative pathways either at site A<sub>3</sub> or B1<sub>L</sub> (Woolford and Baserga, 2013). About 80% of 27SA<sub>2</sub> pre-rRNA is cleaved by MRP RNase at site A<sub>3</sub> in ITS1 (Chu et al., 1994; Lygerou et al., 1996), thereby generating 27SA<sub>3</sub>, which is further trimmed by 5'-3' exoribonucleases Rat1-Rail heterodimer (Henry et al., 1994), Rrp17 (Oeffinger et al., 2009) and Xrn1 (Henry et al., 1994) and requires the presence of A<sub>3</sub> processing factors (Granneman et al., 2011). In the major pathway, 5'-3' exonucleases stop at B1<sub>S</sub>, yielding the 27SB<sub>S</sub> pre-rRNA (Woolford and Baserga, 2013). The remaining 20% of 27SA<sub>2</sub> pre-rRNA is processed by direct cleavage at site B1<sub>L</sub>, producing the 27SB<sub>L</sub> (Faber et al., 2006). The processing of 27SB<sub>S</sub> and 27SB<sub>L</sub> is identical and the cleavage in ITS2 at site C<sub>2</sub> separates the processing pathways of 5.8S and 25S rRNAs, thereby generating the 7S<sub>S</sub>/7S<sub>L</sub> and 26S pre-rRNAs (Woolford and Baserga, 2013). The 26S pre-rRNA is processed by a two-step 5'-3' exonucleolytic digestion that includes the formation of 25.5S pre-rRNA, and is rapidly digested to 25S rRNA by 5'-3' exonuclease Rat1 or Xrn1 (El Hage et al., 2008; Geerlings et al., 2000). The 7S<sub>S</sub>/7S<sub>L</sub> pre-rRNA is processed to 6S<sub>S</sub>/6S<sub>L</sub> pre-rRNA by the exosome complex and is then transported to the cytoplasm where Ngl2 is required to produce mature 5.8S rRNA (Faber et al., 2006; Thomson and Tollervey, 2010). Due to the different pathways and processing sites, the two types of 5.8S rRNA molecules generated, the major 5.8S<sub>S</sub> and the minor 5.8S<sub>L</sub>, differ in their 5' ends by either 7 or 8 nucleotides (Henry et al., 1994).

## 2.7. rRNA modifications

Ribosome biogenesis involves the formation of modified nucleotides in pre-rRNA, which allows changing the biochemical and biophysical properties of certain nucleotides (Motorin and Helm, 2011). ~2–3% of the nucleotides in rRNA are modified (Sloan et al., 2016) and the majority of rRNA modifications are introduced already in the nucleolus (Watkins and Bohnsack, 2012). The most abundant posttranscriptional modifications in eukaryotes are pseudo-

uridylation ( $\Psi$ ) and ribose methylation – namely, there are 44  $\Psi$ s and 55 ribose methylations in the ribosome of budding yeast (Machnicka et al., 2013; Piekna-Przybylska et al., 2008) and 95  $\Psi$ s and 106 ribose methylations in the human ribosome (Krogh et al., 2016). In addition, approximately ten base modifications are introduced in rRNA. The number of modified bases in rRNA is similar in budding yeast and human and most, but not all, modifications are introduced at equivalent positions (Maden and Hughes, 1997; Piekna-Przybylska et al., 2008; Sharma and Lafontaine, 2015). The rRNA modifications are often conserved and clustered at functionally important regions of the ribosome, namely the peptidyl transferase centre in the LSU and the decoding centre in the SSU as well as the A, P, and E sites of tRNA and mRNA binding, the polypeptide exit tunnel, and intersubunit bridges (Decatur and Fournier, 2002). Posttranscriptional modification of rRNA can affect the functions of the ribosome, most probably the efficiency and fidelity of translation (Decatur and Fournier, 2002; Motorin and Helm, 2011).

For decades, it was assumed that each organism contains only one type of ribosome and all positions in the rRNA are always fully methylated; however, measuring the fraction of molecules modified at particular sites demonstrates that in human cells only  $\sim 2/3$  of methylation sites are fully or close to fully methylated, and the remainder are partially methylated, thereby supporting the concept that ribosomes can be heterogeneous (Krogh et al., 2016). In the rRNA of budding yeast, 94 nucleotides, including the majority of base modifications, are close to fully modified, whereas 18 nucleotides are modified in less than 85% of ribosomes (Taoka et al., 2016). In actively growing yeast cells, approximately 68% of adenosines at position 100 of 18S rRNA in both monosomes and polysomes are methylated, which shows that the ribosomes that lack A100 methylation are engaged in translation (Buchhaupt et al., 2014). The extent of 25S rRNA A100 methylation depends on the expression level of the guide RNA, the box C/D snoRNA snR51, which is required for introducing this modification (Buchhaupt et al., 2014).

### **2.7.1. rRNA pseudouridylation and ribose 2'-O-methylation are carried out by ribonucleoprotein complexes**

In eukaryotes, the pseudouridylation and ribose methylation reactions are performed by small ribonucleoprotein (snoRNP) complexes, and guided by box H/ACA (Ganot et al., 1997; Ni et al., 1997) and box C/D snoRNAs (Kiss-László et al., 1996), respectively. The formation of ribose methylation requires the addition of a methyl group to the 2'-oxygen of ribose, catalysed by a methyltransferase fibrillarin (FBL) in humans and by Nop1 in yeast (Tollervey et al., 1993). The ribose methylation reaction takes place five nucleotides upstream of the conserved D or D' box of snoRNA and one box C/D snoRNA can guide the methylation at two alternative sites in the rRNA (Kiss-László et al., 1996; Maden and Hughes, 1997). The formation of  $\Psi$  occurs via iso-

merization of uridine and is performed by  $\Psi$  synthase dyskerin in humans and by Cbf5 in yeast cells (Lafontaine et al., 1998a). The precise role of pseudouridine formation in ribosomes remains unknown, however it offers an additional hydrogen bond donor site and may be required to stabilize the structure and function of rRNA, as shown with the  $\Psi$ s located in human 28S rRNA helix 69, which are part of the intersubunit bridge B2a (Sumita et al., 2005).

Fibrillarin/Nop1 and dyskerin/Cbf5 belong to the box C/D and box H/ACA snoRNP complexes, respectively (Balakin et al., 1996). In addition to enzymes, the box C/D snoRNP complex is composed of RNA binding protein 15.5K/NHPX, NOP56, and NOP58, whereas the box H/ACA snoRNP complex consists of RNA binding protein NHP2, GAR1, and NOP10 (Watkins and Bohnsack, 2012). The genes that encode the components of the snoRNP complexes are essential for the viability of human cells (Blomen et al., 2015) and their down-regulation causes defects in early steps of pre-rRNA processing (Tafforeau et al., 2013)

Although variations in ribosome structure occur under normal conditions, differential expression of ribosome biogenesis factors, including rRNA modification enzymes, can lead to several diseases (Sloan et al., 2016; Tafforeau et al., 2013). The expression level of both snoRNAs and FBL are upregulated in different types of malignancies (Marcel et al., 2013; Martens-Uzunova et al., 2012; Su et al., 2014) and mutations in the DKC1 are the cause of X-linked dyskeratosis congenita, characterized by bone-marrow failure and increased susceptibility to cancer (Heiss et al., 1998). It has been shown using the xenograft model that decreasing the expression of fibrillarin reduces the tumorigenicity of human breast cancer cell line MCF-7 (Su et al., 2014).

### **2.7.2. Stand-alone enzymes catalyse the formation of rRNA base modifications**

In eukaryotes, the formation of base methylation and acetylation are catalysed by stand-alone enzymes. The 25S rRNA of budding yeast contains six methylated bases, located in the peptidyl transferase centre ( $m^1A645$ ,  $m^5C2870$ , and  $m^3U2634$ ), in the subunit interface ( $m^1A2142$  and  $m^5C2278$ ) and in the phosphostalk ( $m^3U2843$ ) (Sharma and Lafontaine, 2015). The 18S rRNA of budding yeast contains six modified bases, located in the decoding site ( $ac^4C1773$ ,  $m_2^6A1781$ , and  $m_2^6A1782$ ), mRNA channel ( $ac^4C1280$ ) and near the tRNA binding sites ( $m^1acp^3\Psi1191$  and  $m^7G1575$ ) (Sharma and Lafontaine, 2015). In addition to performing rRNA modifications, several rRNA methyltransferases are also required for ribosome biogenesis and pre-rRNA processing; however, only the presence of these proteins, and not their catalytic activity is important for ribosome biogenesis (Sharma and Lafontaine, 2015).

### **2.7.2.1. 28S/25S rRNA methyltransferases involved in ribosome biogenesis**

Rrp8 of budding yeast catalyses the formation of m<sup>1</sup>A645 of 25S rRNA (Peifer et al., 2013), while nucleomethylin (NML) is responsible for the formation of 28S rRNA m<sup>1</sup>A1309 in human (Waku et al., 2016). Rrp8 is not essential, however, it affects the biogenesis of both ribosomal subunits (Peifer et al., 2013; Schilling et al., 2012) and pre-rRNA processing within ITS1 at site A<sub>2</sub> (Bousquet-Antonelli et al., 2000). The methyltransferase activity of Rrp8 is not required to process ITS2, which indicates that the roles of Rrp8 in rRNA methylation and ribosome biogenesis are separated (Peifer et al., 2013). Human NML is not required for pre-rRNA processing (Tafforeau et al., 2013), however, it is important for ribosome 60S subunit biogenesis and its depletion suppresses cell proliferation, by activating the p53 pathway (Waku et al., 2016). Under nutrient deficiency, NML inhibits rDNA synthesis and protects cells from energy deprivation-dependent apoptosis (Murayama et al., 2008; Yang et al., 2013).

In budding yeast, the essential nucleolar protein Nop2/Nsun1 (de Beus et al., 1994) is responsible for the formation of m<sup>5</sup>C2870 of 25S rRNA (Sharma et al., 2013a). A decrease in the expression level of Nop2 (Nucleolar protein gene 2) leads to lengthened doubling time and defects in ribosome biogenesis (Hong et al., 1997). It has been shown that the presence of m<sup>5</sup>C2870 methylation is critical because the expression of a catalytically inactive mutant of Nop2 does not rescue either the growth defects, the impairment of ribosome biogenesis, nor the reduced amount of polysomes (Sharma et al., 2013a). However, another study demonstrated that a catalytically dead mutant of Nop2 rescued the growth defect of a  $\Delta$ Nop2 yeast strain, showing that the presence of m<sup>5</sup>C2870 is not important for ribosome biogenesis (Bourgeois et al., 2015). Based on sequence alignment, the homologous protein NSUN1 (also known as p120 and NOL1) is considered as a putative candidate for human 28S rRNA m<sup>5</sup>C4447 modification (Sharma et al., 2013a). NSUN1 is not required for rRNA processing; however, it possesses m<sup>5</sup>C methyltransferase activity and the expression of human NSUN1 in a  $\Delta$ Nop2 yeast strain completely restores the 25S rRNA C2870 methylation and partially rescues its growth defect (Bourgeois et al., 2015; Tafforeau et al., 2013).

In budding yeast, the nucleolar protein Bmt5 and the nucleocytoplasmic protein Bmt6 catalyse the formation of two m<sup>3</sup>U modifications in 25S rRNA (Sharma et al., 2014). Both proteins and methylated residues are conserved among lower eukaryotes, however, these enzymes are not essential and their absence does not affect ribosome biogenesis or cell growth (Sharma et al., 2014).

The budding yeast Bmt2 introduces the m<sup>1</sup>A2142 of 25S rRNA (Sharma et al., 2013b) and Rcm1 catalyses the formation m<sup>5</sup>C2278 of 25S rRNA (Gigova et al., 2014; Sharma et al., 2013a). Both methylations lie in the subunit interface of budding yeast ribosomes, however, neither the presence of these enzymes nor the methylations are essential for cell growth (Sharma et al., 2013b, 2013a).

Yeast 25S rRNA m<sup>1</sup>A2142 is conserved in lower eukaryotes and a *Bmt2* deletion does not cause defects in ribosome biogenesis (Sharma et al., 2013b).

25S rRNA m<sup>5</sup>C2278 is part of the eukaryote specific intersubunit bridge eB14 (Sharma and Lafontaine, 2015). In this bridge, ribosomal protein eL41 connects the LSU and modified nucleotides in the decoding site of SSU, namely the ac<sup>4</sup>C1773, m<sup>6</sup>A1781 and m<sup>6</sup>A1782 of 18S rRNA (Sharma and Lafontaine, 2015). Deletion of *Rcm1* does not interfere with ribosome 60S subunit synthesis (Schosserer et al., 2015; Sharma et al., 2013a). Downregulation of *Rcm1* in yeast and its orthologs in worms and flies lengthens the lifespan of these organisms and increases their resistance to different types of stress, including heat shock and oxidative stress (Schosserer et al., 2015). m<sup>5</sup>C2278 corresponds to 28S rRNA m<sup>5</sup>C3782 in humans and NSUN5/WBSCR20, a conserved protein in eukaryotes, is considered to be the enzyme that catalyses this reaction in human cells (Schosserer et al., 2015; Sharma et al., 2013a). The methyltransferase activity of human NSUN5 has not yet been elucidated, however, it is known to affect ribosome synthesis in the early steps of pre-rRNA processing (Tafforeau et al., 2013).

### **2.7.2.2. 18S rRNA methyltransferases in ribosome biogenesis**

*Dim1/DIMT1 catalyses the formation of dimethyladenosines  
in 3' end of 18S rRNA*

In the small subunit of the ribosome, the structure of 3' end of rRNA (either 16S and 18S rRNA) is highly conserved and contains two dimethylated adenosines, located in helix 45 (Piekna-Przybylska et al., 2008). The formation of these modifications is carried out by a family of methyltransferases that are present in all domains of life (O'Farrell et al., 2008). The Dim1 protein modifies 18S rRNA m<sup>6</sup>A1781 and m<sup>6</sup>A1782 in budding yeast (Lafontaine et al., 1995) and the homologous human protein DIMT1 (also known as DIMT1L and DIM1) (Stanchi et al., 2001) is involved in the formation of 18S rRNA m<sup>6</sup>A1850 and m<sup>6</sup>A1851 (Zorbas et al., 2015). The nucleolar proteins DIMT1 and Dim1 are essential for human and yeast cells, respectively (Blomen et al., 2015; Lafontaine et al., 1994). Downregulation of Dim1 causes severe defects in pre-rRNA processing, at sites A<sub>1</sub> and A<sub>2</sub>, and leads to both the upregulation of 22S pre-rRNA and the downregulation of 27SA and 20S pre-rRNAs and this, in turn, is followed by a depletion in the pool of mature 18S rRNA (Lafontaine et al., 1995). In HeLa cells, depletion of DIMT1 by RNA interference causes uncoupling of cleavages within sites A<sub>0</sub> and 1 which leads to the accumulation of 26S and 43S pre-rRNAs (Zorbas et al., 2015). DIMT1 depletion in other human cell lines (WI-38, RKO and HCT116) causes the level of 26S pre-rRNA to rise and also leads to the accumulation of 21S and 21S-C pre-rRNAs, which results in a reduction of mature 18S rRNA (Zorbas et al., 2015). The expression of catalytically inactive Dim1/DIMT1 mutants is able to rescue defects in pre-

rRNA processing, which demonstrates that the presence of 18S rRNA dimethylation is not required for ribosome biogenesis (Lafontaine et al., 1998b; Zorbas et al., 2015).

Dim1 is a nucleolar protein that binds to the pre-rRNA processing machinery and affects early pre-rRNA cleavages; however, it remains bound with pre-rRNA and catalyses the modification of two  $m_2^6A$  on 20S pre-rRNA in the cytoplasm (Lafontaine et al., 1998b). In human cells, the formation of  $m_2^6A1850$  and  $m_2^6A1851$  within 18S rRNA occurs earlier, already in the nucleus, but after the cleavages that are regulated by the presence of DIMT1 (Zorbas et al., 2015). Delay in rRNA processing and modification events support the hypothesis that the presence of Dim1 and DIMT1 within pre-ribosomes might serve as a quality control mechanism that regulates ribosome biogenesis (Lafontaine et al., 1998b; Zorbas et al., 2015).

#### *Kre33/NAT10 catalyses the formation N4-acetylcytidines*

The Kre33 protein within budding yeast and the human protein NAT10 are important for the introduction of two N4-acetylcytidines within 18S rRNA (Sharma et al., 2015). In humans, the vertebrate-specific box C/D snoRNA U13 (Tyc and Steitz, 1989) is also critical for the formation of  $ac^4C1842$  (Sharma et al., 2015). Expression of a catalytically inactive form of Kre33 demonstrates that 18S rRNA acetylation is not essential for the growth of budding yeast cells (Sharma et al., 2015). However, Kre33 interacts with pre-40S ribosomes and its presence is essential for the biogenesis of the ribosome small subunit, pre-rRNA processing, and cell growth (Ito et al., 2014a; Sharma et al., 2015). Kre33 affects the early nucleolar cleavages (at sites  $A_0$ - $A_2$ ), and, as with Dim1, down-regulation of Kre33 expression leads to the accumulation of 22S pre-rRNA (Lafontaine et al., 1995; Sharma et al., 2015). The depletion of NAT10 causes morphological changes in nucleoli and leads to a slow growth phenotype (Shen et al., 2009), increased apoptosis (Ito et al., 2014b), and reduces the level of 18S rRNA precursors (21S and 18S-E pre-rRNA), which results in a decrease in the production of ribosome 40S subunits (Ito et al., 2014b; Sharma et al., 2015; Tafforeau et al., 2013).

#### *snoRNP complex and stand-alone enzymes participate in the formation of hypermodified residue ( $m^1acp^3\Psi$ )*

The hypermodified residue,  $m^1acp^3\Psi1191$  of 18S rRNA is synthesized in three steps and requires the action of the snoRNP complex as well as stand-alone enzymes. The first step occurs in the nucleolus where the H/ACA snoRNP complex (snoRNP35) carries out a pseudouridylation reaction (Samarsky et al., 1995). Thereafter, the methyltransferase Nep1 (Emg1) catalyses N1-methylation of  $\Psi$  (Meyer et al., 2011) and in the cytoplasm, Tsr3 catalyses the transfer of aminocarboxypropyl (acp) to  $m^1\Psi$  thereby producing  $m^1acp^3\Psi$  (Meyer et al.,

2016). 18S rRNA hypermodification and the enzymes that catalyse its formation are conserved in eukaryotes and it has been shown that the H/ACA box snoRNP complex (contains snoRNA ACA13) and TSR3 are required for the formation of 18S rRNA m<sup>1</sup>acp<sup>3</sup>Ψ1248 in human cells (Kiss et al., 2004; Meyer et al., 2016). The results obtained from an *in vitro* methylation assay demonstrate that human NEP1 is able to methylate Ψ1248 (Wurm et al., 2010).

In budding yeast, deletion of either *snR35* or *Tsr3* causes minor growth defects (Meyer et al., 2016), while the presence of *Nep1* is essential (Eschrich et al., 2002). Like other 18S rRNA methyltransferases, yeast *Nep1* and human NEP1, are important for ribosome small subunit biogenesis (Eschrich et al., 2002; Tafforeau et al., 2013). Downregulation of NEP1 expression slightly decreases the levels of 47S, 45S, and 41S pre-rRNAs (Tafforeau et al., 2013) and in another study, upregulation of 30S, 21S, and 21S-C pre-rRNAs is also documented (Warda et al., 2016). A missense mutation in the gene encoding NEP1/EMG1 (D86G) is the cause of a severe autosomal recessive disease known as Bowen-Conradi syndrome (BCS; OMIM 211180) (Armistead et al., 2009). The BCS mutation does not abolish the methyltransferase activity of NEP1, however, it does disrupt a salt bridge which leads to protein aggregation (Armistead et al., 2009) and its subsequent degradation in a proteasome-dependent manner (Warda et al., 2016). As a result, a decrease in the level of NEP1 protein impairs ribosome SSU maturation in BCS (Warda et al., 2016). TSR3 affects several steps in ribosome small subunit maturation and its depletion in human cells causes the accumulation of 47S and 18S-E pre-rRNAs (Meyer et al., 2016). As with the human phenotype, deletion of *Tsr3* in budding yeast causes accumulation of 35S pre-rRNA as well as an increase in the level of the final precursor of 18S rRNA, and the 20S pre-rRNA (Meyer et al., 2016). Unlike most rRNA methyltransferases, the expression of catalytically inactive mutants of *Tsr3* does not rescue the processing defects (Meyer et al., 2016).

#### *Bud23/WBSCR22 catalyses the formation of N7-methylated guanosine*

Eukaryotic 18S rRNA contains another base methylation, N7-methylated guanosine (m<sup>7</sup>G), which, in the three-dimensional structure of the ribosome, is located in the ridge between the P-site and E-site tRNAs (Sharma and Lafontaine, 2015). The presence of a guanosine base at this location is highly conserved, and corresponds to G1338 in *Escherichia coli*, however, its m<sup>7</sup>G methylation is only present in eukaryotes (m<sup>7</sup>G1575 and m<sup>7</sup>G1639 in budding yeast and human 18S rRNA, respectively) (Piekna-Przybylska et al., 2008; White et al., 2008). Bud23 in yeast and its human ortholog WBSCR22 catalyse these methyl transfer reactions (Haag et al., 2015; White et al., 2008; Zorbas et al., 2015).

rRNA base methyltransferase Bud23 is not essential in yeast, however its absence leads to loss of 18S rRNA m<sup>7</sup>G1575 methylation, pre-rRNA processing defects, impaired ribosome small subunit biogenesis, and decreased cell growth (White et al., 2008). Bud23 interacts with components of the SSU processome, Utp2 and Utp14, but not with U3 snoRNA (Sardana et al., 2013). The loss of

Bud23 expression causes accumulation of the initial pre-rRNA transcript, 35S pre-rRNA, as well as a decrease in 27SA pre-rRNA, which demonstrates that the early pre-rRNA processing step of cleaving A<sub>2</sub> within ITS1 is impaired (Sardana and Johnson, 2012; White et al., 2008). Furthermore, after cleavage within A<sub>2</sub>, Bud23 remains bound to 20S pre-rRNA (Sardana et al., 2013). 20S pre-rRNA accumulation upon *Bud23* deletion shows that the protein is also engaged in late steps of 18S rRNA formation (White et al., 2008). Similarly, pre-rRNA processing defects occur in WBSCR22 depleted human cells (Haag et al., 2015; Tafforeau et al., 2013; Zorbas et al., 2015). On the other hand, the pre-rRNA intermediates that accumulate as a result of WBSCR22 depletion are not identical with those that increase after *Bud23* deletion. The details of this process are discussed in section 5.1.3.

## **2.8. Trm112/TRMT112 is an interaction partner for several methyltransferases**

Budding yeast Trm112 is a 15-kDa zinc-finger protein that interacts with several methyltransferases important for translation. This protein is conserved in all domains of life, however, the residues important for binding the zinc atom are not conserved in metazoans (Bourgeois et al., 2017a). In budding yeast, the interactions of Trm112 with four methyltransferases, i.e., Trm9, Trm11, Mtq2, and Bud23, are considered direct interactions because these proteins co-purify with Trm112 when they are co-expressed in bacterial cells (Figaro et al., 2012; Heurgué-Hamard et al., 2006; Mazauric et al., 2010). A recent review compares various crystal structures of Trm112-methyltransferase complexes and found that Trm112 interacts with its methyltransferase partners in a very similar way, and, methyltransferases probably compete with each other to interact with Trm112 (Bourgeois et al., 2017a). During binding, the  $\beta$ 3 of the methyltransferase forms a  $\beta$ -zipper interaction with  $\beta$ 4 of Trm112 and this complex formation buries a large hydrophobic region on the surface of both partners (Létoquart et al., 2014, 2015; Liger et al., 2011).

Trm9-Trm112 and Trm11-Trm112 complexes are involved in the modification of tRNAs. Trm9-Trm112 modifies the wobble uridine (U34) of some tRNAs and the Trm11-Trm112 complex catalyses the formation of 2-methylguanosine (m<sup>2</sup>G) at position 10 of several tRNAs (Mazauric et al., 2010; Purushothaman et al., 2005). The Mtq2-Trm112 complex catalyses the methylation of translation termination factor eRF1 (Heurgué-Hamard et al., 2006), whereas the Bud23-Trm112 complex catalyses the formation of m<sup>7</sup>G1575 within 18S rRNA and participates in ribosome 40S subunit biogenesis. In addition to direct interactions, co-immunoprecipitation analysis demonstrates that the Trm112 protein forms complexes with 25S rRNA methyltransferases Nop2 and Rcm1 and is bound to 27S pre-rRNA, 7S pre-rRNA, and 25S rRNA, which affects the biogenesis of the large ribosomal subunit (Sardana and Johnson, 2012).



## **2.9. Extracellular vesicles as carriers of biological information**

In order to facilitate intercellular communication in normal and pathophysiological processes, cells release various particles, collectively termed extracellular vesicles (EVs), to the extracellular environment (Abels and Breakefield, 2016). EVs present a heterogeneous family of small membranous particles that originate from either endosomes or from the plasma membrane, and these particles are divided into three groups – e.g., exosomes, microvesicles and apoptotic bodies (Kalra et al., 2012). Exosomes range from 40 to 120 nm in size, are produced inside multivesicular endosomes or multivesicular bodies, and their release occurs when these compartments fuse with the plasma membrane (Colombo et al., 2014). Microvesicles (50–1000 nm) are formed and released via budding from the plasma membrane of the cell (Colombo et al., 2014).

EVs are complex structures that are covered with a lipid bilayer that contains transmembrane proteins and encloses soluble hydrophilic components derived from the cytosol of the donor cell (Colombo et al., 2014). These particles are internalized via endocytosis or membrane fusion, which enables the release of the contents of the particle into the recipient cell (Mulcahy et al., 2014). Cell-derived membrane vesicles are endogenous carriers of cellular proteins, as well as genetic material (DNA, RNA and non-coding RNAs), lipids or sugars and thus, the contents of a particular EV can influence the phenotype of recipient cells (Yáñez-Mó et al., 2015). EVs are deregulated in cancer and the possibility of using EVs to deliver therapeutic cargo is being investigated (van Dommelen et al., 2012). Virus-like particles (VLPs) are produced by ectopic expression of retrovirus Gag protein and are biochemically very similar to EVs and could potentially be used in immunotherapy and for the delivery of therapeutic cargo.

In order to use EVs for medical purpose, it is important to precisely know their contents. Mass-spectrometry based analyses have shown that exosomes contain specific groups of cellular proteins, some of which vary with cell type; whereas others are found in most exosomes regardless of cell type (Colombo et al., 2014).

### 3. AIMS OF THE STUDY

Ribosome biogenesis is a complex process that, in human cells, requires at least 284 trans-acting factors that participate in pre-rRNA processing and modification, ribosome subunit assembly and transport of pre-ribosomal particles (Tafforeau et al., 2013). One third of these factors are medically important because they are related to genetic diseases and cancer (Tafforeau et al., 2013). The gene encoding the WBSCR22 protein is deleted in Williams-Beuren syndrome and its expression is increased in cancers (Doll and Grzeschik, 2001; Merla et al., 2002; Nakazawa et al., 2011; Tiedemann et al., 2012). Orthologs of WBSCR22 – e.g., the budding yeast's Bud23 and *Arabidopsis thaliana*'s Rid2 have nucleolar functions and participate in ribosome biogenesis (Ohbayashi et al., 2011; White et al., 2008). The cellular functions of WBSCR22 were unknown when this current work was undertaken and the aim of the first part of this dissertation is to investigate the functions of WBSCR22 protein in human cells.

The specific aims of the first part of this dissertation (Papers I and II) are the following:

- to compare the expression level of WBSCR22 protein in cell-lines derived from Williams-Beuren syndrome patients and healthy controls,
- to define the subcellular localization of WBSCR22 protein in order to become closer to clarifying its functions in the cells,
- to elucidate, whether WBSCR22 protein is important for cell growth, pre-rRNA processing and ribosome biogenesis,
- to determine the interaction partners of WBSCR22 protein,
- to elucidate whether the interacting proteins influence the functions of WBSCR22 protein.

The second part of this work includes the formation and characterization of MLV Gag based virus-like particles carrying melanoma antigens. These are chimeric enveloped VLPs, formed by outward budding at the plasma membrane of host cells and therefore, host proteins are always included in VLPs. The aim of Paper III is to study the protein content of MLV-Gag induced virus-like particles using mass-spectrometry.

## 4. MATERIALS AND METHODS

Detailed descriptions of the materials and methods used in this study are provided in the original publications and briefly reproduced here.

In general, the cellular functions of WBSCR22 protein were studied in human tissue culture cells in which the expression level of WBSCR22 was reduced using RNA interference. For WBSCR22 depletion, WBSCR22 siRNAs were transfected into cells and analysed 72 hours after transfection by various methods discussed below. In order to analyse the role of WBSCR22 downregulation on cell growth, the number of siWBSCR22 treated and control cells was counted up to 5 days. Studying the role of WBSCR22 in ribosome biogenesis and pre-rRNA processing required fractionating the cells into cytoplasmic and nuclear fractions. The cytoplasmic lysate from siWBSCR22 treated cells was used in polyribosome profile analysis. For pre-rRNA processing analysis, RNA from whole cells, cytoplasmic and nuclear fractions was extracted and separated by agarose gel electrophoresis and the abundance of different 18S rRNA processing precursors were evaluated by Northern blot. HeLa cell lysate ultracentrifugation through a sucrose gradient, followed by gradient fractionation, immunoblot, and RNA analysis enabled us to estimate the sedimentation properties of endogenous WBSCR22 protein. Immunoblot analysis was used to study the expression level of WBSCR22 protein in cell-lines derived from the Williams-Beuren syndrome patients.

In order to investigate which proteins interact with WBSCR22, a cell line that stably expresses epitope tagged WBSCR22 protein was generated and then studied using co-immunoprecipitation analysis. For this purpose, these cells, together with control cells, were metabolically labelled in SILAC media and analysed using mass-spectrometry. The most significant interactions found by analysing the SILAC data were then evaluated using immunoblot analysis. Based on these results, the immunofluorescence analysis was used to study the subcellular localization of WBSCR22, TRMT112, and C1QBP. For this, the fluorescent fusion proteins EGFP-WBSCR22 and TRMT112-mCherry were expressed in order to determine the subcellular localisation of the WBSCR22-TRMT112 complex in live cells. Both immunoblot analysis and flow cytometry were used to monitor the expression of ectopically expressed WBSCR22 and EGFP-WBSCR22, respectively. Treatment with a 26S proteasome inhibitor (MG132) followed by immunoblot or flow cytometry enabled us to analyse whether the expression WBSCR22 is regulated by proteasomes.

In order to produce MLV Gag based VLPs, the mouse fibroblast cell line COP5 was transfected with either plasmids that encode the MLV Gag protein to produce MLV Gag VLPs, or with plasmids that encode both the MLV Gag protein and the melanoma antigen (MART1, TRP1, MAGEA4, MAGEA10 or MCAM), to produce VLPs that carry one of the melanoma antigens. The VLPs were purified from cell culture media and characterized using various methods. To study the protein content of MLV Gag VLPs and VLPs that carry melanoma antigens, fractions that contain the VLPs were pooled and studied using mass-spectrometry.

## 5. RESULTS AND DISCUSSION

### 5.1. WBSCR22 protein is required for cell growth, ribosome small subunit biogenesis and pre-rRNA processing (Papers I and II)

#### 5.1.1. The level of WBSCR22 protein is reduced in cell lines derived from Williams-Beuren syndrome patients

*WBSCR22* is one of the genes expressed from the single-copy gene region of Williams-Beuren syndrome chromosome locus (Doll and Grzeschik, 2001; Merla et al., 2002). Expression of most of the transcripts encoded by the 7q11.23 chromosome region, including *WBSCR22* mRNA, is reduced in WBS patients (Merla et al., 2006). However, for two genes, dosage-compensation mechanisms are known to exist, and their mRNA levels remain unaffected compared to healthy controls (Merla et al., 2006). Crucially, the existence of possible compensatory mechanisms acting at the protein level remains largely unexplored. To analyse the expression level of WBSCR22 protein, the immunoblot analysis from lymphoblastoid cell lines (LCL) derived from Williams-Beuren syndrome patients, was performed (Paper I, Fig. 6A). This analysis demonstrates that the expression level of WBSCR22 protein is reduced to a level similar to the level of mRNA found in cell lines derived from WBS patients, which indicates that WBSCR22 is a dosage-sensitive gene and therefore studying its role in human cells is of great interest.

#### 5.1.2. WBSCR22 depletion reduces cell growth

Several works have shown that the mRNA of *WBSCR22* is upregulated in different types of malignancies, including invasive breast cancer, hepatocellular carcinoma, and multiple myeloma (Nakazawa et al., 2011; Stefanska et al., 2014; Tiedemann et al., 2012). In cancer cells, the expression level of WBSCR22 regulates cell survival and metastatic potential (Nakazawa et al., 2011; Tiedemann et al., 2012). In the cell lines derived from WBS patients, the levels of WBSCR22 mRNA (Merla et al., 2006) and protein (Paper I, Fig. 6) are reduced. The gene-dosage of multiple genes is altered in microdeletion syndromes (such as WBS) and also in malignancies, and one way to understand the function of individual genes is to observe cells deficient in the gene of interest. To determine the role of WBSCR22 in regulating cell growth, RNA interference (RNAi) was used to silence the expression of the *WBSCR22* gene in HeLa cells, and cell growth was monitored for 5 days (Paper I, Fig. 1B). Surprisingly, when compared to control cells, the cells transfected with siRNA targeting *WBSCR22* mRNA displayed significantly reduced growth at 72, 96, and 120 hours after transfection. The doubling time of control cells was 21 hours, whereas in WBSCR22-depleted cells, the doubling time was lengthened to 25 hours. This

clearly demonstrates that WBSR22 is important for cell growth. This observation is in accordance with the results obtained by gene-trap mutagenesis and CRIPR/Cas9-directed gene editing systems that both demonstrate that WBSR22 is essential for cell viability under laboratory experimental conditions (Blomen et al., 2015; Wang et al., 2015a). In comparison, the budding yeast protein Bud23, which is an ortholog of WBSR22, is not essential for cell viability, yet the deletion of *Bud23* causes a slow-growth phenotype (White et al., 2008). Similarly, in *Arabidopsis thaliana*, the presence of the WBSR22 ortholog RID2 is required for plant growth (Ohbayashi et al., 2011). Therefore, as with its orthologs RID2 and Bud23, WBSR22 is a factor that helps to regulate cell growth.

### **5.1.3. WBSR22 is required for ribosome small subunit biogenesis and pre-rRNA processing**

In order to determine whether human WBSR22 affects ribosome biogenesis, its expression was reduced by siRNAs and a polyribosome profile analysis of cytoplasmic extracts was performed. As shown in Paper I, Fig. 2A, down-regulation of *WBSR22* causes defects in ribosome small subunit biogenesis that are characterized by a decrease in the amount of free 40S subunits in the cytoplasm, together with an increase in the amount of free 60S subunits. This ribosome subunit imbalance results in a diminished amount of 80S ribosomes in WBSR22 depleted cells, compared to control cells. This, in turn, can reduce the capability to translate and may be the main cause of increased cell doubling time (Paper I, Fig. 1A). Compared to controls, the amount of mature 18S rRNA decreases in the cytoplasm of WBSR22 depleted cells, and the decreased ratio of 18S to 28S rRNA (Paper I, Fig. 2B) indicates that the observed defects in ribosome small subunit biogenesis may arise due to errors in the production of mature 18S rRNA.

Mature 18S rRNA is produced from 47S pre-rRNA transcript by exo- and endoribonucleolytic cleavages in a complex and hierarchical pre-rRNA processing pathway (Henras et al., 2015; Mullineux and Lafontaine, 2012). In human cells, at least 286 proteins affect pre-rRNA processing (Tafforeau et al., 2013). With the aim to analyse the role of WBSR22 in pre-rRNA processing, cells transfected with siRNAs against *WBSR22* were fractionated into cytoplasmic and nuclear parts, and the RNA from each fraction as well as whole cells was analysed by Northern blot (Paper I, Fig. 3B and C). In this experiment, a oligonucleotide probe that hybridizes to the 5' end of ITS1 and recognizes 18S rRNA processing precursors, including 45S, 41S, 30S, 21S and 18S-E pre-rRNAs, was used. As depicted in Paper I (Fig. 3B and C), WBSR22 does not affect the early pre-rRNA processing events because the relative amount of 45S, 41S and 30S pre-rRNAs remained unaffected in the HeLa and HEK293 cells used in this study. However, when the amount of WBSR22 was depleted, a slight decrease in 21S pre-rRNA occurs in the cell nucleus, together with a

3-fold accumulation of 18S-E pre-rRNA. This indicates that WBSCR22 is involved in late nuclear steps of pre-rRNA processing. Accumulation of 18S-E pre-rRNA upon WBSCR22 depletion was also uncovered in a siRNA screen that analysed the involvement of nucleolar proteins in pre-rRNA processing (Tafforeau et al., 2013) and later in other articles (Haag et al., 2015; Zorbas et al., 2015).

18S-E pre-rRNA is the final precursor of 18S rRNA that appears in the nucleus, and is then exported to the cytoplasm where the final maturation of pre-40S subunits take place (Henras et al., 2015; Mullineux and Lafontaine, 2012; Rouquette et al., 2005). In whole cell extract, the accumulation of 18S-E pre-rRNA does not indicate, whether the late nuclear processing events, or subsequent cytoplasmic cleavage at site, 3 is impaired. In WBSCR22 depleted cells, the diminished amount of free 40S particles and the nuclear accumulation of 18S-E pre-rRNA indicate that WBSCR22 can be important for ribosome small subunit export. Alternatively, the effect can be indirect, and instead of influencing transport to the cytoplasm, defective pre-40S particles can become trapped in the nucleus and accumulate. In mammalian cells, 18S-E pre-rRNA is comprised of mature 18S rRNA with an additional ~15 to 80 nucleotides of ITS1 sequence at its 3' end (Haag et al., 2015; Preti et al., 2013; Rouquette et al., 2005). This variation in ITS1 sequence length arises because 18S-E pre-rRNA processing is performed by 3'-5' exoribonucleolytic trimming that occurs during the transport of pre-40S particles (Preti et al., 2013). As a result of WBSCR22 depletion, the average length of 18S rRNA 3' end extensions increases, which indicates that WBSCR22 is required for proper 3'-5' trimming of 18S-E pre-rRNA prior to its export to the cytoplasm (Haag et al., 2015).

The expression of exogenous WBSCR22 is able to rescue the pre-rRNA processing defect present in WBSCR22 siRNA treated cells because the accumulation of 18S-E pre-rRNA disappears at an expression level similar to the endogenous protein (Paper I, **Fig. 3D and E**).

WBSCR22 is a methyltransferase that catalyses the formation of 18S rRNA m<sup>7</sup>G1639 methylation; however, as with the wild-type protein, it has been shown that the expression of catalytically inactive mutants, WBSCR22-G63E and WBSCR22-D82K, also rescue the ribosome biogenesis defect which implies that WBSCR22 protein also has a methyltransferase activity-independent role in ribosome biogenesis (Haag et al., 2015; Zorbas et al., 2015). Likewise, the methyltransferase activity of the budding yeast protein Bud23 is not critical for pre-rRNA processing because the exogenous expression of catalytically inactive forms of Bud23 rescues the pre-rRNA processing defects and the slow-growth phenotype of a *Bud23Δ* yeast strain (White et al., 2008). Similarly, the rRNA methyltransferase activities of human DIMT1 as well as yeast Dim1 and Emg1/Nep1 are dispensable for ribosome biogenesis and cell growth (Lafontaine et al., 1995; Leulliot et al., 2008; Zorbas et al., 2015). Thus, rRNA methyltransferases often have two independent roles in ribosome biogenesis, and their participation in pre-rRNA processing seems to be more critical than their enzymatic activity as rRNA methyltransferases. Nevertheless, evolution has

maintained the catalytic activity of enzymes that modify rRNA, and rRNA modifications are clustered at functionally important regions of the ribosome, thereby implying that they are beneficial for the function of ribosomes (Decatur and Fournier, 2002).

#### **5.1.4. WBSCR22 sediments with ribosome 40S subunits**

To both perform 18S rRNA G1639 methylation and participate in pre-rRNA processing, WBSCR22 should bind to pre-ribosomes. Therefore, the sedimentation behaviour of WBSCR22 protein in sucrose density gradient ultracentrifugation was analysed (Paper I, Fig. 5B and C). The results of this experiment revealed that WBSCR22 sediments at positions that correspond to ribosome 40S/pre-40S subunits. However, it did also sediment in fractions smaller than 40S subunits, which indicates that WBSCR22 is not exclusively associated with ribosomal particles. Compared to WBSCR22, Bud23 exhibits a broader sedimentation profile, namely in peaks corresponding to 40S/pre-40S and 80S/90S sized fractions (Figaro et al., 2012; L etoquart et al., 2014; Sardana and Johnson, 2012). The 90S pre-ribosome or SSU processome represents an early ribosome biogenesis complex that consists of pre-rRNA, U3 snoRNA, early binding transacting factors and ribosomal proteins (Dragon et al., 2002; Grandi et al., 2002). Bud23 sedimentation at 90S pre-ribosomes is in agreement with the genetic and physical interaction between Bud23 and SSU processome components Utp2 and Utp14 (Sardana et al., 2013). It has been proposed that Bud23 acts at the transition between the early and middle stages of 40S subunit synthesis at the time of A<sub>2</sub> cleavage, which separates the biogenesis pathways of LSU and SSU in budding yeast (Sardana et al., 2013). In contrast, WBSCR22 depletion affects processing events that occur after the cleavage within ITS1, which may explain why it does not sediment in large pre-ribosome fractions (Paper I, Fig. 3B and C).

#### **5.1.5. WBSCR22 is a nuclear protein that can accumulate within nucleoli**

The subcellular localization analysis of WBSCR22 in HeLa cells demonstrates that endogenous WBSCR22 exhibits a diffuse signal throughout the cell nucleus and it is neither accumulated nor excluded from the nucleoli (Paper I, Fig. 5A). A similar pattern of WBSCR22 staining is also reported for the human melanoma cell line A375M (Nakazawa et al., 2011). However, in some works, WBSCR22 displays a nuclear and cytoplasmic staining with enrichment in the nucleus (Jangani et al., 2014; Zorbas et al., 2015). WBSCR22 contains a C-terminal nuclear localization signal in addition to its N-terminal methyltransferase domain (Doll and Grzeschik, 2001; Merla et al., 2002), and probably moves through nuclear pore complexes (NPC) by active carrier-mediated transport (Marfori et al., 2011). In addition, because of its small size (~32 kDa), WBSCR22

can also enter the nucleus by passive diffusion (Marfori et al., 2011). Small variations in the results of indirect immunofluorescence analyses can be explained by the properties of the antibodies used – different antibodies can either recognize different epitopes of WBSCR22, or, furthermore, antibodies can also be unspecific and bind to epitopes of other proteins. One alternative to immunofluorescence analysis is to take advantage of recombinant proteins composed of an in-frame fusion between a fluorescent protein, such as EGFP, and the protein of interest. The use of fluorescent fusion proteins enables one to study the localisation of proteins and protein complexes in live cells. As seen in live cell microscopy analysis (Paper II, Fig. 3A), the fusion protein EGFP-WBSCR22 can be detected in the nucleoplasm as well as in nucleoli. The C-terminal domain of WBSCR22 (EGFP-WBSCR22-CTD), which contains the NLS, can be detected in both the nucleus and nucleoli. In contrast, the methyltransferase domain of WBSCR22 (EGFP-WBSCR22-MTD), which lacks the NLS, is enriched in the cytoplasm. The nucleolar accumulation of WBSCR22 indicates that WBSCR22 is probably involved in both nucleolar as well as nucleoplasmic processes. In *Sacchamycetes cerevisiae*, Bud23-GFP localizes mainly in the nucleolus (White et al., 2008). Compared to WBSCR22, the nucleolar accumulation of Bud23 is more pronounced and corresponds with its involvement in earlier pre-rRNA processing steps (Sardana et al., 2013; White et al., 2008). In *Arabidopsis thaliana*, RID2-GFP expressed from the *RID2* promoter localizes in the nucleus with accumulation in the nucleolus, which is in accordance with its role in early and intermediate pre-rRNA processing steps (Ohbayashi et al., 2011).

## **5.2. A principal interaction partner of WBSCR22 in human cells is TRMT112 (Paper II)**

### **5.2.1. WBSCR22 interacts with TRMT112**

For a better understanding of the cellular functions of WBSCR22, it is important to know its interaction partners. In order to investigate the protein complexes that contain WBSCR22, a SILAC (stable isotope labelling by amino acids in the cell culture) coupled co-immunoprecipitation analysis followed by identification by mass-spectrometry was performed. For this experiment, cells that stably express BPV1 E2-tagged WBSCR22 were generated from the U2OS cell line. As seen in Paper II, Fig. 1A, the expression level of epitope-tagged WBSCR22 in the U2OS-E2Tag-WBSCR22 cell line is similar to the endogenous protein, which enables one to study its function close to the physiological expression level. The same cell line was used for co-immunoprecipitation analysis and here the intensity ratio of heavy to light peptides reflects the relative abundance of peptides enriched in WBSCR22 pull-down fraction compared to control.



As seen in Fig. 1C within Paper II, the peptides of ribosomal proteins, as well as YBX1 (Y-box binding protein 1), C1QBP (complement C1q binding protein), TRMT112 (tRNA methyltransferase 11–2 homolog (*S. cerevisiae*)) and WBSR22 itself were enriched in WBSR22 pull-down fractions. C1QBP is a multifunctional protein involved in the complement pathway (Ghebrehiwet et al., 1994), mRNA splicing (Petersen-Mahrt et al., 1999), transcription regulation (Chattopadhyay et al., 2004), apoptosis (Itahana and Zhang, 2008) and also pre-rRNA processing (Yoshikawa et al., 2015). C1QBP localizes mainly in the cytoplasm and mitochondria; however, a fraction of this protein is found in nucleoli, where it interacts with proteins involved in ribosome biogenesis, including FBL and NOP52, and affects the early steps in pre-rRNA processing (Yoshikawa et al., 2011). Although WBSR22 is involved in the late steps of pre-rRNA processing, the enrichment of C1QBP in WBSR22 pull-down fractions suggests that WBSR22 might also associate with 90S pre-ribosomes. The interaction between C1QBP and WBSR22 has been detected in another mass-spectrometry based analysis (Zhang et al., 2013), but could not be confirmed by immunoblot analysis, which may reflect the transient nature of this interaction (Paper II, Fig. 1D). C1QBP interacts with and regulates the activity of YBX1 (Wang et al., 2015b), another protein found in the current study. Although the interaction between WBSR22 and YBX1 proteins was not analysed by immunoblotting in this work, its importance for ribosome biogenesis needs to be evaluated in future studies.

TRMT112 exhibits the highest enrichment (H/L ratio) in the WBSR22 pull-down fraction (Paper II, Fig. 1C) and is also confirmed as a WBSR22 interacting protein by immunoblot analysis (Paper II, Fig. 1B). In agreement with the results of this work, another study that investigated the interaction partners of over-expressed WBSR22 by mass-spectrometry also identified C1QBP, YBX1, and TRMT112 in WBSR22 pull-down fractions (Jangani et al., 2014). The budding yeast ortholog of TRMT112, Trm112, is a small zinc-finger protein (~15 kDa) that interacts with several methyltransferases, including the tRNA methyltransferases Trm9 and Trm11 (Purushothaman et al., 2005; Studte et al., 2008), the protein methyltransferase Mtq2 (Heurgué-Hamard et al., 2006) and the rRNA methyltransferases Nop2 and Rcm1 (Sardana and Johnson, 2012). Human TRMT112 protein also interacts with methyltransferases involved in translation, namely with ALKBH8 (ABH8) and HEMK2 $\alpha$ , which methylate the wobble uridine of certain tRNAs and the translation termination factor eRF1, respectively (Figaro et al., 2008; Fu et al., 2010). Thus, in complex with various methyltransferases, budding yeast Trm112 and human TRMT112 serve as cofactors. Notably, the interaction between WBSR22 and TRMT112 is required for pre-rRNA processing as well as facilitating the 18S rRNA m<sup>7</sup>G 1639 methylation (Zorbas et al., 2015).

### 5.2.2. TRMT112 regulates the stability of WBSCR22

In order to study the functional importance of the protein-protein interactions revealed in this work, the expression levels of WBSCR22, TRMT112, and C1QBP were reduced by siRNAs, and the efficiency of depletion was determined by immunoblotting. As seen in **Fig. 2A** within Paper II, the expression of all proteins analysed was reduced upon treatment with their respective siRNAs. Furthermore, transfection with siRNA that targets *TRMT112* reduced the expression level of TRMT112 itself and also decreased the expression level of WBSCR22 protein in HeLa cells. This indicates that the presence of TRMT112 is important for the stability of WBSCR22 protein. In contrast to TRMT112 depletion, downregulation of C1QBP expression by RNA interference did not alter the expression level of WBSCR22. In order to determine if the decrease in WBSCR22 protein level is specific to a reduced level of TRMT112 protein rather than an off-target effect of the particular siRNA sequence, alternative siRNA sequences that target *TRMT112* mRNA were designed and used in a comparable study. As with the original TRMT112 siRNA, other siRNA sequences were effective in decreasing the TRMT112 mRNA expression level and all led to a decrease in WBSCR22 expression in the human cell lines HeLa and U2OS (Paper II, **Fig. 2B and C**). In addition, in the African green monkey cell line COS-7, siRNAs that target *TRMT112* also reduce the expression of both TRMT112 and WBSCR22 (Paper II, **Fig. 2D**).

siRNAs complementary to the 3' untranslated region (3'UTR) of human *TRMT112* mRNA decrease the expression of endogenous TRMT112; however, these siRNAs do not target exogenously expressed TRMT112 that lacks the native 3'UTR, which makes it possible to perform rescue experiments. In complementation analysis, the expression of exogenous TRMT112 is able to restore the normal protein level of WBSCR22 in cells where the expression of endogenous TRMT112 is silenced by siRNA transfection (Paper II, **Fig. 2E**). These results reveal that the expression level of WBSCR22 protein is positively regulated by the expression of TRMT112 and indicate that the availability of TRMT112 is crucial for the stability of WBSCR22.

Unpublished results from our research group demonstrate that in *Escherichia coli*, recombinant histidine-tagged WBSCR22 protein is insoluble and forms inclusion bodies. On the other hand, co-expression of WBSCR22 and its interaction partner TRMT112 increases the solubility of WBSCR22 in bacterial cells which implies that TRMT112 regulates folding and thereby the stability of the WBSCR22 protein (Zorbas et al., 2015). Similarly, the budding yeast Bud23, Mtd2, and Trm9 proteins are largely insoluble in *E. coli*, and the co-expression of these methyltransferases with Trm112 increases their solubility (Figaro et al., 2012; Heurgué-Hamard et al., 2006; Mazauric et al., 2010). Crystal structure analysis has revealed that Bud23 interacts with Trm112 through the formation of a parallel  $\beta$ -zipper involving the Bud23 strand  $\beta$ 3 and the Trm112 strand  $\beta$ 4, thereby covering the hydrophobic residues of Bud23 (Létoquart et al., 2014). Thus, shielding the hydrophobic amino acid residues of Bud23 increases the

solubility and stability of this protein (Létoquart et al., 2014). In budding yeast cells, Trm112 is required for the stability of Bud23 and Mtq2, but not for Trm11, Nop2, and Rcm1 (Sardana and Johnson, 2012). The interaction mode of other complexes (Trm11-Trm112, Mtq2-Trm112, and Trm9-Trm112 heterodimers) is similar to the Bud23-Trm112 complex and only the tRNA methyltransferase Trm11 can be expressed and purified in *E. coli* without Trm112 co-expression (Bourgeois et al., 2017b).

### 5.2.3. WBSR22 regulates the subcellular localization of TRMT112

We and others have previously shown by indirect immunofluorescence analysis that endogenous WBSR22 localizes diffusely in the cell nucleus (Paper I, Fig. 5A; Nakazawa et al., 2011; Zorbas et al., 2015). *In silico* predictions indicate that ~50% of human TRMT112 protein localizes to the cytoplasm (Towns and Begley, 2012). In accordance with this, indirect immunofluorescence analyses reveal that human TRMT112 distributes to both the nucleus and the cytoplasm of HeLa cells, and, moreover, is excluded from the nucleolus (Paper II, Fig. S1). In another publication, human TRMT112 was found to be enriched in the nucleus and excluded from the nucleoli, with only a fraction of the protein found in the cytoplasm (Zorbas et al., 2015). Comparing the work of our group and D. Lafontaine's group, the sub-nuclear distribution of TRMT112 protein is identical; however, the amount and distribution of cytoplasmic TRMT112 protein is different. The same antibody against human TRMT112 protein as in our work was also used for the Human Protein Atlas database; however, in their work, TRMT112 protein co-localizes with microtubules, while only a small fraction of the protein is nuclear (Uhlén et al., 2015). Thus, the results of different groups are in partial disagreement and, therefore, the use of another method for studying the subcellular localization of TRMT112 would be beneficial. In order to further examine the subcellular distribution of TRMT112, a construct that encodes the fluorescent fusion protein TRMT112-mCherry was generated. As demonstrated in Fig. 3A of Paper II, the control protein mCherry and TRMT112-mCherry can be detected throughout the entire cell, both in the nucleus and in the cytoplasm. These data are in agreement with immunofluorescence analysis of the TRMT112 protein and indicate that TRMT112 can shuttle between the nucleus and cytoplasm of the cell.

In order to analyse the subcellular distribution of the WBSR22 and TRMT112 complex, EGFP-WBSR22 and TRMT112-mCherry fusion proteins were co-expressed in human cells and analysed using live cell microscopy. The fluorescence signals of EGFP-WBSR22 and TRMT112-mCherry were detected in the cell nucleus, indicating that WBSR22-TRMT112 complex localizes mainly in nucleoplasm (Paper II, Fig. 3B). When WBSR22 is exogenously expressed together with TRMT112, the latter protein redistributes to the cell nucleus, thereby implying that WBSR22 affects the subcellular localization of TRMT112. However, while TRMT112-mCherry in complex with EGFP-

WBSCR22 can be detected in the nucleolus it does not display enrichment towards this compartment.

Based on co-immunoprecipitation analysis, WBSCR22 interacts with TRMT112 via its N-terminal methyltransferase domain (MTD) (Paper II, Fig. 3D). The fusion protein EGFP-WBSCR22-MTD, which lacks the C-terminally located NLS, localizes within the cytoplasm of HeLa cells, and, crucially, when co-expressed with TRMT112-mCherry, both proteins localize within the cytoplasm (Paper II, Fig. 3B). On the other hand, the chimeric protein EGFP-WBSCR22-CTD, which is comprised of the EGFP protein fused with 38 amino acids from the C-terminus of WBSCR22, does not interact with TRMT112 (Paper II, Fig. 3E). In agreement with the other experiments, co-expression of EGFP-WBSCR22-CTD and TRMT112-mCherry does not change the sub-cellular localization of the latter protein – EGFP-WBSCR22-CTD is a nuclear protein that is enriched in the nucleoli, whereas in the same cells, TRMT112-mCherry can be detected in the cytoplasm and nucleus of the cell and is excluded from the nucleoli (Paper II, Fig. 3B). These results demonstrate that the methyltransferase domain of WBSCR22 protein is important for its interaction with TRMT112, and, furthermore, the localization of the WBSCR22-TRMT112 complex is determined by the WBSCR22 protein.

#### **5.2.4. WBSCR22 mutants defective in TRMT112 binding are less stable than the wild-type protein**

In order to gain more knowledge about the biological function of the WBSCR22-TRMT112 interaction, WBSCR22 mutants with substitutions in amino acids located on the interface of the WBSCR22-TRMT112 complex were generated. A crystal structure analysis of the budding yeast Bud23-Trm112 complex indicates that the interaction interface consists of a large hydrophobic core surrounded by polar residues (Létoquart et al., 2014). A sequence alignment between Bud23 and WBSCR22 proteins demonstrates that the amino acid residues involved in complex formation are conserved between budding yeast and human (Létoquart et al., 2014).

Based on the conservation data, two putative TRMT112 binding deficient mutants of WBSCR22 protein – WBSCR22-KT/AA and WBSCR22-D117A – were constructed in the current work (Paper II). In the WBSCR22-D117A mutant, the conserved aspartic acid at position 117 of WBSCR22 is replaced with alanine, and in the WBSCR22-KT/AA double-mutant, the lysine and threonine at positions 112 and 115, respectively, are replaced with alanines. In addition, these mutants contained either an E2Tag epitope tag in their N-terminus or were expressed as fusion proteins in-frame with EGFP.

The amino acid D112 of Bud23, corresponding to D117 in WBSCR22, is involved in the formation of a salt bridges between Bud23 and Trm112, and the Bud23 D112A and D112G mutants exhibit a slow-growth phenotype (Létoquart et al., 2014). The amino acids R107 and S110 of Bud23 are both involved in the

formation of hydrogen bonds in the Bud23-Trm112 complex, and correspond to K112 and T115 in the WBSCR22 protein (Létoquart et al., 2014). As shown by two hybrid assays in yeast, single-mutant R107L, S110P, and D112G forms of Bud23, as well as triple-mutant Bud23, have reduced binding to Trm112 (Sardana and Johnson, 2012; Sardana et al., 2014). Much less is known about the interaction between human WBSCR22 and TRMT112 proteins.

With the aim of tracking the expression of the WBSCR22 protein variants described above, HeLa cells were transiently transfected with plasmids that either encode for EGFP-WBSCR22, or one of its mutants KT/AA and D117A, and the number of EGFP positive cells was monitored up to three days after transfection using flow cytometry analysis. As shown in Fig. 4A of Paper II, 24 hours after transfection approximately 50% of the cells expressed the EGFP and EGFP-WBSCR22 proteins, whereas only about 40% of the cells expressed the mutant proteins EGFP-WBSCR22-D117A and EGFP-WBSCR22-KT/AA. After 48 hours, the percentage of cells expressing EGFP and EGFP-WBSCR22 proteins was still close to 50%, whereas the number of cells expressing mutant proteins EGFP-WBSCR22-D117A and EGFP-WBSCR22-KT/AA had declined to 15–20%. 72 hours after transfection, the difference between WBSCR22 and its mutants was even more pronounced. Similar results were seen in other human cell lines.

The expression of WBSCR22 protein and its mutants using flow cytometry was also confirmed using immunoblot analysis. For this experiment, the epitope-tagged WBSCR22 protein was expressed in HeLa cells, which were lysed 24 hours after transfection. As seen in Fig. 4B of Paper II, the expression level of wild-type WBSCR22 protein was elevated compared to the mutant proteins WBSCR22-KT/AA and D117A, which were barely detectable. The ectopic expression of epitope-tagged TRMT112 protein increased the abundance of wild-type WBSCR22 protein as well as the expression levels of the mutant proteins, which indicates that the mutants have possibly retained some ability to bind to TRMT112. Nevertheless, over-expression of TRMT112 had more influence on the protein level of wild-type WBSCR22 protein compared to its impact on the expression levels of the WBSCR22 mutants. The protein level of TRMT112 was also increased by co-expression with WBSCR22, indicating that WBSCR22 and TRMT112 proteins stabilize each other within the cells.

In order to investigate how strongly the WBSCR22 point-mutations affect the interaction with TRMT112 protein, a co-immunoprecipitation analysis was performed in HeLa cells. Epitope-tagged wild-type WBSCR22 protein was able to pull-down the endogenous TRMT112 protein (Paper II, Fig. 4C), and, reciprocally, epitope-tagged TRMT112 co-immunoprecipitated with the endogenous WBSCR22 protein (Paper II, Fig. 4D). Unfortunately, due to the low expression levels of the WBSCR22 mutants in HeLa cells, the results of the pull-down experiments did not provide insight into whether the interaction with TRMT112 was affected in these cells. Therefore, a similar co-immunoprecipitation analysis was performed in COS-7 cells. COS-7 cells express the large T-antigen of simian virus 40 (SV40), allowing plasmids containing the

SV40 origin of replication to replicate in these cells, thereby augmenting the level of plasmid-driven gene expression. The vector that encodes EGFP protein replicates in COS-7 cells (Paper II, Fig. 4A) and the expression levels of WBSCR22 protein and its mutants are high in COS-7 cells (Paper II, Fig. 4E). Furthermore, the number of cells expressing EGFP-WBSCR22 and its mutant variants remained constant for three days (Paper II, Fig. 4A). Fig. 4E within Paper II shows that the wild-type WBSCR22 protein and also both mutants (KT/AA and D117A) were able to pull-down endogenous TRMT112 protein. However, the resulting amount of TRMT112 protein was different between WBSCR22 and its mutants. Mutating WBSCR22 lysine 112 and threonine 115 results in about a 2-fold decrease of co-precipitated TRMT112, while mutating the highly conserved aspartic acid 117 results in an almost complete loss of interaction between WBSCR22 and TRMT112. This confirms that the residues K112, T115, and D117 within WBSCR22 are all important for interacting with TRMT112.

The subcellular localization of EGFP-WBSCR22-D117A is similar to the wild-type protein, i.e., the D117A mutant localises to the nucleus and nucleoli (Paper II, Fig. 3B). Surprisingly, the co-expression of EGFP-WBSCR22-D117A and TRMT112-mCherry caused an enrichment of TRMT112-mCherry in the nucleus. This result points to the possibility that the nuclear accumulation of TRMT112-mCherry is caused by endogenous WBSCR22 protein. In support of this, the depletion of endogenous WBSCR22 by RNA interference decreased the amount of WBSCR22-D117A and TRMT112 in the nucleus and increased the amount of cytoplasmic TRMT112-mCherry (Paper II, Fig. S2). Nevertheless, these results indicate that the WBSCR22-D117A mutant has retained some ability to bind to TRMT112.

### **5.2.5. The level of WBSCR22 protein is regulated by the ubiquitin-proteasome pathway**

The stability of WBSCR22 protein is controlled by its interaction partner TRMT112, and WBSCR22 mutants with reduced affinity towards TRMT112 are less stable than wild-type WBSCR22. Furthermore, ectopically expressed EGFP-WBSCR22 wild-type seems to be less stable than the control protein EGFP. In eukaryotic cells, the turnover of cellular proteins is chiefly regulated by the ubiquitin-proteasome pathway. In order to find out whether the amount of WBSCR22 protein and its mutants are controlled by proteasomal degradation, HeLa cells were transfected with plasmids encode for EGFP, EGFP-WBSCR22, and its mutants, and the number of cells expressing EGFP was monitored in the presence and absence of MG132, an inhibitor of the proteasome. MG132 was added to the cells 24 hours after transfection, and cells were subsequently incubated for 16 hours before analysis by flow cytometry (Paper II, Fig. 6A). Inhibition of the ubiquitin-proteasome pathway slightly increased the percentage of cells expressing EGFP-WBSCR22; however, the proportion

of cells expressing EGFP protein did not change. The percentage of cells expressing EGFP-WBSCR22-D117A and EGFP-WBSCR22-KT/AA increased to the level of wild-type WBSCR22. These results indicate that the turnover of wild-type WBSCR22 as well as its mutants is regulated by proteasomal degradation. Immunoblot analysis confirmed that the inhibition of proteasomes with MG132 increased the amount of ectopically expressed WBSCR22, whereas the amount of endogenous tubulin did not change (Paper II, Fig. 6B). In the same experiment, a slight increase in the expression level of endogenous TRMT112 protein was also detected, implying that the degradation of TRMT112 protein is also under the control of the ubiquitin-proteasome pathway. As shown in Fig. 6C within Paper II, the downregulation of TRMT112 expression decreased the amount of WBSCR22 protein, whereas the inhibition of proteasomes is able to increase the amount of WBSCR22 in TRMT112-depleted cells. This indicates that unbound WBSCR22 is degraded by the ubiquitin-proteasome pathway.

Proteasome-dependent degradation involves the addition of poly-ubiquitin chains to the target molecule. In order to study whether ubiquitin chains are added to WBSCR22 protein, a co-immunoprecipitation experiment was performed. In this experiment, HeLa cells were transfected with plasmids that encode HA-tagged ubiquitin and E2-tagged WBSCR22, and the antibody against E2 Tag was used to precipitate WBSCR22 complexes. As shown in Fig. 6D within Paper II, WBSCR22 and WBSCR22-D17A pull-down fractions contained poly-ubiquitinated molecules, which again confirm that the amount of WBSCR22 is controlled by proteasome-dependent degradation.

### **5.2.6. WBSCR22 complements the growth and ribosome biogenesis defect of budding yeast $\Delta$ Bud23 mutant strain**

The budding yeast  $\Delta$ Bud23 strain is characterized by a slow growth phenotype, which is caused by defects in ribosome biogenesis and pre-rRNA processing (White et al., 2008). The human WBSCR22 protein is also involved in the formation of 18S rRNA and the ribosome small subunit, and the decreased expression of WBSCR22 protein also affects the rate of cell growth (Paper I, Fig. 1 and 2). In order to investigate whether the functions of these two proteins are conserved, a complementation experiment was performed in the  $\Delta$ Bud23 yeast strain. As demonstrated in Paper I, Fig. 4A, the expression of the budding yeast protein Bud23 is able to restore a normal growth phenotype and defects in ribosome biogenesis. The expression of WBSCR22 protein in this yeast strain partially complements both defects, which indicates that the function of human WBSCR22 and budding yeast Bud23 are conserved, at least in part. In comparison, others have shown that the expression of wild-type RID2 protein of *Arabidopsis thaliana* does not rescue the slow growth phenotype of  $\Delta$ Bud23 (Ohbayashi et al., 2011). Compared to the negative control, the expression of WBSCR22 mutants D117A and KT/AA also enhanced the slow growth of  $\Delta$ Bud23, although to a much lesser extent than the expression of wild-type

WBSCR22 (Paper II, Fig. 5A). Further, the expression of WBSCR22 mutants with reduced TRMT112 binding is not able to restore the ribosome biogenesis defects of  $\Delta Bud23$  (Paper II, Fig. 5B). These mutants may have lost the ability to bind yeast Trm112, which can reduce their stability in yeast cells in a similar manner to what was observed in human cells.

### **5.3. Ribosomal proteins and tRNA synthetases are incorporated into MLV Gag induced extracellular vesicles (Paper III)**

Recombinant virus-like particles (VLPs) can be used to transfer biological information between cells and tissues. In the second part of this dissertation, the protein content of retrovirus Gag protein induced VLPs was analysed. This information is important to avoid unwanted side-effects, when using these VLPs in biological systems. This work is part of a larger project that aimed to produce VLPs that express melanoma antigens with the purpose of using them for immunotherapy of melanoma patients.

In mammalian cells, the expression of retroviral Gag protein causes the spontaneous formation and release of virus-like particles to the tissue culture media (Larson et al., 2005). In the current work, the budding properties of Gag protein of murine leukemia virus (MLV) were used to produce a family of chimeric VLPs that each carry one of the following cancer/testis antigens: MART1, TRP1, MAGEA4, MAGEA10, or MCAM. The results of dynamic light scattering and transmission electron microscopy experiments indicate that the average size of MLV Gag VLPs is ~140 nm and the size of VLPs that carry melanoma antigens are very similar to each other. The assembled particles have a spherical structure and regular shape.

In order to characterize the protein content of Gag VLPs and VLPs carrying melanoma antigens, LC-MS/MS-based label-free quantitative proteomics analysis was performed. The results of these proteomics tests are presented in Supplementary Dataset 1 and in Fig. 5 within Paper III. The results indicate that all six types of VLPs contained the MLV Gag protein, and in addition, the melanoma antigens were incorporated into their respective VLPs. An immunoblot analysis of the MLV Gag protein further confirms that this protein is included in all particles (Paper III, Fig. 5E). However, in addition to exogenously expressed proteins, the mass-spectrometry analysis enabled to detect that VLPs contained more than 500 proteins acquired from the host cells. Probably, most of these cellular proteins are encapsulated into VLPs. Nonetheless, some of the cellular proteins might bind to the surface of VLPs after the assembly of VLPs.

In order to better understand which proteins form the components of these VLPs, the identified cellular proteins were divided into several classes based on their function. Among the proteins identified by LC-MS/MS analysis were Tsg101, Alix, and Nedd4, which are all proteins that directly interact with MLV



Gag protein and are involved in the budding of VLPs (Segura-Morales et al., 2005). Tsg101 and Alix are components of extracellular vesicles and are considered as markers of exosomes (Colombo et al., 2014). In addition, all VLPs contained the clathrin adaptor complex subunit AP-1, which is involved in retroviral release (Camus et al., 2007).

In addition to proteins that facilitate the budding of particles, these VLPs contained cellular proteins from various groups, including cell adhesion, membrane and cytoskeleton proteins, ribosomal proteins, translation factors, tRNA synthetases, metabolic enzymes, RNA binding proteins, and RNA helicases. The results of immunoblot analyses (Paper III, Fig. 5E) support these LC-MS/MS findings and demonstrate that cytoskeleton protein  $\alpha$ -tubulin and ribosomal protein Rps6 are components of the MLV Gag based VLPs. Others have also reported that ribosomal proteins and rRNA can be incorporated into MLV VLPs (Muriaux et al., 2002); however, the significance of incorporating components of translational machinery (e.g., ribosomal proteins, tRNA synthetases, and translation factors) into VLPs remains unknown. Mutations in the MLV Gag protein cause the assembly of immature particles, where the viral genomic RNA is replaced with ribosomal RNA (Muriaux et al., 2002). These authors suggest that intact ribosomes or ribosomal subunits are packaged into the VLPs (Muriaux et al., 2002). Similarly, other proteomic characterizations of endogenous extracellular vesicles demonstrate that both microvesicles and exosomes contain ribosomal proteins and other proteins involved in translation (Kowal et al., 2016; Willms et al., 2016). The majority of proteins found in the exosomes of neuroblastoma cells are either cytoplasmic (~40%) or membrane proteins (~16%) (Marimpietri et al., 2013). It is possible that ribosomal proteins that naturally locate within the cytoplasm are passively incorporated into VLPs and other types of EVs. To support this, a comparison of the proteome of host cell membrane and HIV Gag induced particles found that the protein composition of VLPs and the plasma membrane is similar and the bulk of the host proteins in the plasma membrane are passively incorporated into the VLPs (Hammarstedt and Garoff, 2004).

## 6. CONCLUSIONS

Based on the results of Paper I and Paper II presented in the first part of this dissertation, the following conclusions can be made:

- The expression level of WBSCR22 protein is decreased in cell lines derived from Williams-Beuren syndrome patients. This result is in agreement with a reduced level of WBSCR22 mRNA in these cells.
- WBSCR22 is a nuclear protein that exhibits a diffuse signal throughout the nucleoplasm in human cells. When overexpressed, some of the WBSCR22 protein accumulates within the nucleoli, which reveals that WBSCR22 might have an affinity to bind with pre-ribosomes.
- WBSCR22 is important for cell growth, ribosome small subunit biogenesis, and pre-rRNA processing. Depletion of WBSCR22 causes the nuclear accumulation of the last precursor of 18S rRNA, the 18S-E pre-rRNA, thereby leading to reduced production of mature 18S rRNA and ribosomal 40S subunits.
- WBSCR22 interacts with TRMT112 protein in human cells and this interaction is important for the stability of WBSCR22 protein.
- The subcellular localization of WBSCR22-TRMT112 complexes are determined by WBSCR22.
- The expression level of WBSCR22 protein is tightly controlled. Under normal conditions, WBSCR22 protein is either bound to TRMT112 in the nucleoplasm or localizes to nucleolus where it may be associated with nucleolar proteins or rRNA. Under these conditions, when the amount of TRMT112 is limited, e.g., WBSCR22 is overexpressed or the expression level of TRMT112 is decreased, unbound WBSCR22 is tagged with ubiquitin and degraded by the 26S ubiquitin-proteasome pathway.

The results of the second part of this dissertation enabled to conclude that virus-like particles, induced by the exogenous expression of murine leukemia virus Gag protein, contain several groups of cellular proteins, including ribosomal proteins, tRNA synthetases, and translation initiation factors. It is currently unknown whether the incorporation of ribosomal proteins into VLPs are functionally relevant or are present by chance.

## SUMMARY IN ESTONIAN

### **Williams-Beureni sündroomi kromosoomiregiooni valk WBSCR22 kui ribosoomi biogeneesifaktor**

Ribosoomide biogenees on kompleksne protsess, milles osaleb ligi 300 valku ning neist valkudest umbes kolmandik on seotud genoomsete haiguste ja kasvaja tekkega. Williams-Beureni sündroom (WBS) on multisüsteemne arenguhäire, mida põhjustab on umbes 30 geeni puudumine seitsmenda kromosoomi pikast õlast, lookusest 7q11.23. WBS patsientidel esineb rida erinevaid probleeme, millest kõige sagedasemateks on kaasasündinud südame-veresoonkonna haigused, aga ka sidekoe arenguhäired, neuroloogilised probleemid, spetsiifiline kognitiivne profiil, kasvuprobleemid ja iseloomulikud näojooned. Genoomsed ümberkorraldused nagu deletsioonid, duplikatsioonid ja inversioonid WBS kromosoomiregioonis tekivad selle genoomipiirkonna kompleksuse tõttu. Nimelt ümbritsevad 7q11.23 piirkonna geenidega ala madalakoopiaarvulised kordusjärjestused, mis oma kõrge sarnasuse tõttu põhjustavad valepaardumisest tingitud ümberkorraldusi. Ehkki mehhanismid, mis viivad 7q11.23 piirkonna genoomsete häirete tekkimiseni, on suhteliselt hästi iseloomustatud, on vähem teada, kuidas üksikute geenide koopiaarvu muutus ja sellest tulenev avaldumistaseme erinevus põhjustab haigustunnuseid. Hetkel arvatakse, et paljud geenid on olulised WBS-i ning sama piirkonna duplikatsioonisündroomi kujunemisel, mistõttu on vajalik nendelt geenidelt ekspresseerivate valkude funktsioonide uurimine. Üheks WBS lookusest ekspresseeritavaks valguks, mille funktsioon enne antud doktoritöö tegemist oli teadmata, on WBSCR22. Nimetatud valgu iseloomustamine on oluline ka seetõttu, et on näidatud tema kõrget avaldumistaset mitmete kasvaja korral ning on teada, et WBSCR22 on oluline kasvajakude elulemise ja metastaatilise potentsiaali reguleermisel.

Oma struktuurilt on WBSCR22 valk metüültransferaas ning bioinformaatiliste meetoditega on talle ennustatud DNA metüleerimise aktiivsust. Uuritud on ka WBSCR22 valgu rolli histoonide metüleerijana, kuid alles hiljuti avastati, et tegelikult on WBSCR22 ensüüm, mis katalüüsib 18S ribosomaalse RNA (18S rRNA) metüleerimist. Antud doktoritöö esimese ning peamise osa eesmärgiks oli WBSCR22 funktsiooni uurimine imetaja rakkudes.

Antud töö raames leiti, et WBSCR22 valk asub hajusalt rakutuumas, kuid tema kõrgema avaldumistaseme juures on näha valgu kogunemist tuumakesse. Selles tuumasiseses struktuuris toimub rRNA süntees ja algab ribosoomide biogenees. WBSCR22 valgu paiknemine tuumakeses viitab, et ta võib seonduda varajaste pre-ribosoomidega. Eukarüootide ribosoomide biogenees hõlmab pre-rRNA ja ribosoomivalkude sünteesimist, pre-rRNA protsessimist, rRNA-de modifitseerimist, ribosoomi subühikute kokkupanemist, nende ekspordit tuumast tsütoplasmasse ning sünteesitavate ribosoomide kvaliteedi kontrollimist. Antud doktoritöös leiti, et WBSCR22 on oluline rakkude jagunemiseks, ribosoomi väikese subühiku biogeneesil ja pre-rRNA protsessingul. Täpsemalt, WBSCR22 ekspressiooni mahasurumine põhjustab 18S-E pre-rRNA kuhjumist raku tuumas.

18S-E pre-rRNA on viimane 18S rRNA eellane, mida leidub nii raku tuumas kui tsütoplasmas. Pärast 18S-E pre-rRNA eksporti toimub tema viimane endoribonukleelüütiline lõikamine, mille tulemusel moodustub küps 18S rRNA. WBSCR22 valgu avaldumistaseme vähenemine toob kaasa 18S rRNA hulga languse tsütoplasmas, mistõttu pärsitakse ribosoomi 40S subühikute teke ning see omakorda võib põhjustada rakkude aeglasemat jagunemist.

Selleks et uurida, millised on WBSCR22 valgu interaktsioonipartnerid, konstrueeriti inimese rakuliin, mis püsivalt tootis epitoopmärgisega WBSCR22 valku ning leiti, et üheks WBSCR22-ga seonduvaks valguks on TRMT112. Lisaks näidati, et TRMT112 paiknemine rakus on määratud WBSCR22 poolt, WBSCR22-TRMT112 kompleks paikneb raku tuumas ning WBSCR22-TRMT112 kompleksi teke on vajalik WBSCR22 stabiilsuse tagamiseks. WBSCR22 valgu hulk rakus on kontrollitud – tuumaplasmas on WBSCR22 seotud TRMT112-ga ning tuumakese piirkonnas ilmselt tuumakese valkude või rRNA-ga. Juhul kui rakkudes ei ole piisavalt TRMT112, näiteks WBSCR22 üleekspressiooni või TRMT112 vaigistamise korral, siis WBSCR22 valk märgistatakse ubikvitiini molekulidega ja lagundatakse 26S proteasoomi rajas.

Selle doktoritöö teises osas analüüsiti retroviiruste Gag valgu poolt indutseeritud viiruslaadsete partiklite (VLP-de) valgulist koostist ning avastati, et nendes VLP-des esineb suur hulk rakulisi valke, nende seas ka suur hulk ribosoomivalke. Paljude rakuliste valkude, sealhulgas ribosoomivalkude roll, viiruslaadsetes partiklites ei ole teada.

## REFERENCES

- Abbas, E., Cox, D.M., Smith, T., and Butler, M.G. (2016). The 7q11.23 Microduplication Syndrome: A Clinical Report with Review of Literature. *J. Pediatr. Genet.* *5*, 129–140.
- Abels, E.R., and Breakefield, X.O. (2016). Introduction to Extracellular Vesicles: Biogenesis, RNA Cargo Selection, Content, Release, and Uptake. *Cell. Mol. Neurobiol.* *36*, 301–312.
- Anger, A.M., Armache, J.-P., Berninghausen, O., Habeck, M., Subklewe, M., Wilson, D.N., and Beckmann, R. (2013). Structures of the human and *Drosophila* 80S ribosome. *Nature* *497*, 80–85.
- Ansel, K.M., Pastor, W.A., Rath, N., Lapan, A.D., Glasmacher, E., Wolf, C., Smith, L.C., Papadopoulou, N., Lamperti, E.D., Tahiliani, M., et al. (2008). Mouse Eri1 interacts with the ribosome and catalyzes 5.8S rRNA processing. *Nat. Struct. Mol. Biol.* *15*, 523–530.
- Armistead, J., Khatkar, S., Meyer, B., Mark, B.L., Patel, N., Coghlan, G., Lamont, R.E., Liu, S., Wiechert, J., Cattini, P.A., et al. (2009). Mutation of a gene essential for ribosome biogenesis, EMG1, causes Bowen-Conradi syndrome. *Am. J. Hum. Genet.* *84*, 728–739.
- Balakin, A.G., Smith, L., and Fournier, M.J. (1996). The RNA world of the nucleolus: two major families of small RNAs defined by different box elements with related functions. *Cell* *86*, 823–834.
- Baumer, A., Dutly, F., Balmer, D., Riegel, M., Tükel, T., Krajewska-Walasek, M., and Schinzel, A.A. (1998). High level of unequal meiotic crossovers at the origin of the 22q11.2 and 7q11.23 deletions. *Hum. Mol. Genet.* *7*, 887–894.
- Bayés, M., Magano, L.F., Rivera, N., Flores, R., and Pérez Jurado, L.A. (2003). Mutational mechanisms of Williams-Beuren syndrome deletions. *Am. J. Hum. Genet.* *73*, 131–151.
- Belin, S., Beghin, A., Solano-González, E., Bezin, L., Brunet-Manquat, S., Textoris, J., Prats, A.-C., Mertani, H.C., Dumontet, C., and Diaz, J.-J. (2009). Dysregulation of ribosome biogenesis and translational capacity is associated with tumor progression of human breast cancer cells. *PLoS One* *4*, e7147.
- Beuren, A.J., Apitz, J., and Harmjanz, D. (1962). Supravalvular aortic stenosis in association with mental retardation and a certain facial appearance. *Circulation* *26*, 1235–1240.
- de Beus, E., Brockenbrough, J.S., Hong, B., and Aris, J.P. (1994). Yeast NOP2 encodes an essential nucleolar protein with homology to a human proliferation marker. *J. Cell Biol.* *127*, 1799–1813.
- Blomen, V.A., Májek, P., Jae, L.T., Bigenzahn, J.W., Nieuwenhuis, J., Staring, J., Sacco, R., van Diemen, F.R., Olk, N., Stukalov, A., et al. (2015). Gene essentiality and synthetic lethality in haploid human cells. *Science* *350*, 1092–1096.
- Boisvert, F.-M., van Koningsbruggen, S., Navascués, J., and Lamond, A.I. (2007). The multifunctional nucleolus. *Nat. Rev. Mol. Cell Biol.* *8*, 574–585.
- Bourgeois, G., Ney, M., Gaspar, I., Aigueperse, C., Schaefer, M., Kellner, S., Helm, M., and Motorin, Y. (2015). Eukaryotic rRNA Modification by Yeast 5-Methylcytosine-Methyltransferases and Human Proliferation-Associated Antigen p120. *PLoS One* *10*, e0133321.

- Bourgeois, G., L toquart, J., van Tran, N., and Graille, M. (2017a). Trm112, a Protein Activator of Methyltransferases Modifying Actors of the Eukaryotic Translational Apparatus. *Biomolecules* 7.
- Bourgeois, G., Marcoux, J., Saliou, J.-M., Cianf rani, S., and Graille, M. (2017b). Activation mode of the eukaryotic m2G10 tRNA methyltransferase Trm11 by its partner protein Trm112. *Nucleic Acids Res.* 45, 1971–1982.
- Bousquet-Antonelli, C., Vanrobays, E., G lugne, J.P., Caizergues-Ferrer, M., and Henry, Y. (2000). Rrp8p is a yeast nucleolar protein functionally linked to Gar1p and involved in pre-rRNA cleavage at site A2. *RNA N. Y. N* 6, 826–843.
- Buchhaupt, M., Sharma, S., Kellner, S., Oswald, S., Paetzold, M., Peifer, C., Watzinger, P., Schrader, J., Helm, M., and Entian, K.-D. (2014). Partial methylation at Am100 in 18S rRNA of baker’s yeast reveals ribosome heterogeneity on the level of eukaryotic rRNA modification. *PLoS One* 9, e89640.
- Camus, G., Segura-Morales, C., Molle, D., Lopez-Verg es, S., Begon-Pescia, C., Cazevieuille, C., Schu, P., Bertrand, E., Berlioz-Torrent, C., and Basyuk, E. (2007). The Clathrin Adaptor Complex AP-1 Binds HIV-1 and MLV Gag and Facilitates Their Budding. *Mol. Biol. Cell* 18, 3193–3203.
- Carron, C., O’Donohue, M.-F., Choessel, V., Faubladi r, M., and Gleizes, P.-E. (2011). Analysis of two human pre-ribosomal factors, bystin and hTsr1, highlights differences in evolution of ribosome biogenesis between yeast and mammals. *Nucleic Acids Res.* 39, 280–291.
- Chattopadhyay, C., Hawke, D., Kobayashi, R., and Maity, S.N. (2004). Human p32, interacts with B subunit of the CCAAT-binding factor, CBF/NF-Y, and inhibits CBF-mediated transcription activation in vitro. *Nucleic Acids Res.* 32, 3632–3641.
- Cherniske, E.M., Carpenter, T.O., Klaiman, C., Young, E., Bregman, J., Insogna, K., Schultz, R.T., and Pober, B.R. (2004). Multisystem study of 20 older adults with Williams syndrome. *Am. J. Med. Genet. A.* 131, 255–264.
- Chu, S., Archer, R.H., Zengel, J.M., and Lindahl, L. (1994). The RNA of RNase MRP is required for normal processing of ribosomal RNA. *Proc. Natl. Acad. Sci. U. S. A.* 91, 659–663.
- Ciganda, M., and Williams, N. (2011). Eukaryotic 5S rRNA biogenesis. *Wiley Interdiscip. Rev. RNA* 2, 523–533.
- Colombo, M., Raposo, G., and Th ry, C. (2014). Biogenesis, secretion, and intercellular interactions of exosomes and other extracellular vesicles. *Annu. Rev. Cell Dev. Biol.* 30, 255–289.
- Cusc , I., Corominas, R., Bay s, M., Flores, R., Rivera-Brugu s, N., Campuzano, V., and P rez-Jurado, L.A. (2008). Copy number variation at the 7q11.23 segmental duplications is a susceptibility factor for the Williams-Beuren syndrome deletion. *Genome Res.* 18, 683–694.
- Decatur, W.A., and Fournier, M.J. (2002). rRNA modifications and ribosome function. *Trends Biochem. Sci.* 27, 344–351.
- Denisov, S., Lessard, F., Mayer, C., Stefanovsky, V., van Driel, M., Grummt, I., Moss, T., and Stunnenberg, H.G. (2011). A model for the topology of active ribosomal RNA genes. *EMBO Rep.* 12, 231–237.
- Derenzini, M., Pasquinelli, G., O’Donohue, M.-F., Ploton, D., and Thiry, M. (2006). Structural and functional organization of ribosomal genes within the mammalian cell nucleolus. *J. Histochem. Cytochem. Off. J. Histochem. Soc.* 54, 131–145.

- Doll, A., and Grzeschik, K.H. (2001). Characterization of two novel genes, WBSR20 and WBSR22, deleted in Williams-Beuren syndrome. *Cytogenet. Cell Genet.* *95*, 20–27.
- van Dommelen, S.M., Vader, P., Lakhal, S., Kooijmans, S. a. A., van Solinge, W.W., Wood, M.J.A., and Schiffelers, R.M. (2012). Microvesicles and exosomes: opportunities for cell-derived membrane vesicles in drug delivery. *J. Control. Release Off. J. Control. Release Soc.* *161*, 635–644.
- Dragon, F., Gallagher, J.E.G., Compagnone-Post, P.A., Mitchell, B.M., Porwancher, K.A., Wehner, K.A., Wormsley, S., Settlage, R.E., Shabanowitz, J., Osheim, Y., et al. (2002). A large nucleolar U3 ribonucleoprotein required for 18S ribosomal RNA biogenesis. *Nature* *417*, 967–970.
- Ebersberger, I., Simm, S., Leisegang, M.S., Schmitzberger, P., Mirus, O., von Haeseler, A., Bohnsack, M.T., and Schleiff, E. (2014). The evolution of the ribosome biogenesis pathway from a yeast perspective. *Nucleic Acids Res.* *42*, 1509–1523.
- Eichler, D.C., and Craig, N. (1994). Processing of eukaryotic ribosomal RNA. *Prog. Nucleic Acid Res. Mol. Biol.* *49*, 197–239.
- El Hage, A., Koper, M., Kufel, J., and Tollervey, D. (2008). Efficient termination of transcription by RNA polymerase I requires the 5' exonuclease Rat1 in yeast. *Genes Dev.* *22*, 1069–1081.
- Eronen, M., Peippo, M., Hiippala, A., Raatikka, M., Arvio, M., Johansson, R., and Kähkönen, M. (2002). Cardiovascular manifestations in 75 patients with Williams syndrome. *J. Med. Genet.* *39*, 554–558.
- Eschrich, D., Buchhaupt, M., Kötter, P., and Entian, K.-D. (2002). Nep1p (Emg1p), a novel protein conserved in eukaryotes and archaea, is involved in ribosome biogenesis. *Curr. Genet.* *40*, 326–338.
- Ewart, A.K., Morris, C.A., Atkinson, D., Jin, W., Sternes, K., Spallone, P., Stock, A.D., Leppert, M., and Keating, M.T. (1993). Hemizygoty at the elastin locus in a developmental disorder, Williams syndrome. *Nat. Genet.* *5*, 11–16.
- Faber, A.W., Vos, H.R., Vos, J.C., and Raué, H.A. (2006). 5'-end formation of yeast 5.8SL rRNA is an endonucleolytic event. *Biochem. Biophys. Res. Commun.* *345*, 796–802.
- Falnes, P.Ø., Jakobsson, M.E., Davydova, E., Ho, A., and Małeckı, J. (2016). Protein lysine methylation by seven- $\beta$ -strand methyltransferases. *Biochem. J.* *473*, 1995–2009.
- Farrar, J.E., Nater, M., Caywood, E., McDevitt, M.A., Kowalski, J., Takemoto, C.M., Talbot, C.C., Meltzer, P., Esposito, D., Beggs, A.H., et al. (2008). Abnormalities of the large ribosomal subunit protein, Rpl35a, in Diamond-Blackfan anemia. *Blood* *112*, 1582–1592.
- Ferrero, G.B., Howald, C., Micale, L., Biamino, E., Augello, B., Fusco, C., Turturo, M.G., Forzano, S., Reymond, A., and Merla, G. (2010). An atypical 7q11.23 deletion in a normal IQ Williams-Beuren syndrome patient. *Eur. J. Hum. Genet. EJHG* *18*, 33–38.
- Figaro, S., Scrima, N., Buckingham, R.H., and Heurgué-Hamard, V. (2008). HemK2 protein, encoded on human chromosome 21, methylates translation termination factor eRF1. *FEBS Lett.* *582*, 2352–2356.
- Figaro, S., Wacheul, L., Schillewaert, S., Graille, M., Huvelle, E., Mongeard, R., Zorbas, C., Lafontaine, D.L.J., and Heurgué-Hamard, V. (2012). Trm112 is required for Bud23-mediated methylation of the 18S rRNA at position G1575. *Mol. Cell. Biol.* *32*, 2254–2267.

- Fu, D., Brophy, J.A.N., Chan, C.T.Y., Atmore, K.A., Begley, U., Paules, R.S., Dedon, P.C., Begley, T.J., and Samson, L.D. (2010). Human AlkB homolog ABH8 is a tRNA methyltransferase required for wobble uridine modification and DNA damage survival. *Mol. Cell. Biol.* *30*, 2449–2459.
- Ganot, P., Bortolin, M.L., and Kiss, T. (1997). Site-specific pseudouridine formation in preribosomal RNA is guided by small nucleolar RNAs. *Cell* *89*, 799–809.
- Geerlings, T.H., Vos, J.C., and Raué, H.A. (2000). The final step in the formation of 25S rRNA in *Saccharomyces cerevisiae* is performed by 5'→3' exonucleases. *RNA* *6*, 1698–1703.
- Ghebrehiwet, B., Lim, B.L., Peerschke, E.I., Willis, A.C., and Reid, K.B. (1994). Isolation, cDNA cloning, and overexpression of a 33-kD cell surface glycoprotein that binds to the globular “heads” of C1q. *J. Exp. Med.* *179*, 1809–1821.
- Gigova, A., Duggimpudi, S., Pollex, T., Schaefer, M., and Koš, M. (2014). A cluster of methylations in the domain IV of 25S rRNA is required for ribosome stability. *RNA N. Y. N* *20*, 1632–1644.
- Gothelf, D., Farber, N., Raveh, E., Apter, A., and Attias, J. (2006). Hyperacusis in Williams syndrome: characteristics and associated neuroaudiologic abnormalities. *Neurology* *66*, 390–395.
- Grandi, P., Rybin, V., Bassler, J., Petfalski, E., Strauss, D., Marzioch, M., Schäfer, T., Kuster, B., Tschochner, H., Tollervey, D., et al. (2002). 90S pre-ribosomes include the 35S pre-rRNA, the U3 snoRNP, and 40S subunit processing factors but predominantly lack 60S synthesis factors. *Mol. Cell* *10*, 105–115.
- Granneman, S., and Baserga, S.J. (2004). Ribosome biogenesis: of knobs and RNA processing. *Exp. Cell Res.* *296*, 43–50.
- Granneman, S., Petfalski, E., and Tollervey, D. (2011). A cluster of ribosome synthesis factors regulate pre-rRNA folding and 5.8S rRNA maturation by the Rat1 exonuclease. *EMBO J.* *30*, 4006–4019.
- Haag, S., Kretschmer, J., and Bohnsack, M.T. (2015). WBSR22/Merm1 is required for late nuclear pre-ribosomal RNA processing and mediates N7-methylation of G1639 in human 18S rRNA. *RNA N. Y. N* *21*, 180–187.
- Hadjiolova, K.V., Nicoloso, M., Mazan, S., Hadjiolov, A.A., and Bachellerie, J.P. (1993). Alternative pre-rRNA processing pathways in human cells and their alteration by cycloheximide inhibition of protein synthesis. *Eur. J. Biochem. FEBS* *212*, 211–215.
- Hammarstedt, M., and Garoff, H. (2004). Passive and active inclusion of host proteins in human immunodeficiency virus type 1 gag particles during budding at the plasma membrane. *J. Virol.* *78*, 5686–5697.
- Heindl, K., and Martinez, J. (2010). Nol9 is a novel polynucleotide 5'-kinase involved in ribosomal RNA processing. *EMBO J.* *29*, 4161–4171.
- Heiss, N.S., Knight, S.W., Vulliamy, T.J., Klauck, S.M., Wiemann, S., Mason, P.J., Poustka, A., and Dokal, I. (1998). X-linked dyskeratosis congenita is caused by mutations in a highly conserved gene with putative nucleolar functions. *Nat. Genet.* *19*, 32–38.
- Henderson, A.S., Warburton, D., and Atwood, K.C. (1973). Letter: Ribosomal DNA connectives between human acrocentric chromosomes. *Nature* *245*, 95–97.
- Henras, A.K., Plisson-Chastang, C., O'Donohue, M.-F., Chakraborty, A., and Gleizes, P.-E. (2015). An overview of pre-ribosomal RNA processing in eukaryotes. *Wiley Interdiscip. Rev. RNA* *6*, 225–242.



- Henry, Y., Wood, H., Morrissey, J.P., Petfalski, E., Kearsey, S., and Tollervey, D. (1994). The 5' end of yeast 5.8S rRNA is generated by exonucleases from an upstream cleavage site. *EMBO J.* *13*, 2452–2463.
- Heurgué-Hamard, V., Graille, M., Scrima, N., Ulryck, N., Champ, S., van Tilbeurgh, H., and Buckingham, R.H. (2006). The zinc finger protein Ynr046w is plurifunctional and a component of the eRF1 methyltransferase in yeast. *J. Biol. Chem.* *281*, 36140–36148.
- Hong, B., Brockenbrough, J.S., Wu, P., and Aris, J.P. (1997). Nop2p is required for pre-rRNA processing and 60S ribosome subunit synthesis in yeast. *Mol. Cell. Biol.* *17*, 378–388.
- van Hoof, A., Lennertz, P., and Parker, R. (2000). Three conserved members of the RNase D family have unique and overlapping functions in the processing of 5S, 5.8S, U4, U5, RNase MRP and RNase P RNAs in yeast. *EMBO J.* *19*, 1357–1365.
- Hughes, J.M., and Ares, M. (1991). Depletion of U3 small nucleolar RNA inhibits cleavage in the 5' external transcribed spacer of yeast pre-ribosomal RNA and impairs formation of 18S ribosomal RNA. *EMBO J.* *10*, 4231–4239.
- Idol, R.A., Robledo, S., Du, H.-Y., Crimmins, D.L., Wilson, D.B., Ladenson, J.H., Bessler, M., and Mason, P.J. (2007). Cells depleted for RPS19, a protein associated with Diamond Blackfan Anemia, show defects in 18S ribosomal RNA synthesis and small ribosomal subunit production. *Blood Cells. Mol. Dis.* *39*, 35–43.
- Itahana, K., and Zhang, Y. (2008). Mitochondrial p32 is a critical mediator of ARF-induced apoptosis. *Cancer Cell* *13*, 542–553.
- Ito, S., Akamatsu, Y., Noma, A., Kimura, S., Miyauchi, K., Ikeuchi, Y., Suzuki, T., and Suzuki, T. (2014a). A single acetylation of 18 S rRNA is essential for biogenesis of the small ribosomal subunit in *Saccharomyces cerevisiae*. *J. Biol. Chem.* *289*, 26201–26212.
- Ito, S., Horikawa, S., Suzuki, T., Kawauchi, H., Tanaka, Y., Suzuki, T., and Suzuki, T. (2014b). Human NAT10 Is an ATP-dependent RNA Acetyltransferase Responsible for N4-Acetylcytidine Formation in 18 S Ribosomal RNA (rRNA). *J. Biol. Chem.* *289*, 35724–35730.
- Jangani, M., Poolman, T.M., Matthews, L., Yang, N., Farrow, S.N., Berry, A., Hanley, N., Williamson, A.J.K., Whetton, A.D., Donn, R., et al. (2014). The methyltransferase WBSR22/Merm1 enhances glucocorticoid receptor function and is regulated in lung inflammation and cancer. *J. Biol. Chem.* *289*, 8931–8946.
- Kalra, H., Simpson, R.J., Ji, H., Aikawa, E., Altevogt, P., Askenase, P., Bond, V.C., Borràs, F.E., Breakefield, X., Budnik, V., et al. (2012). Vesiclepedia: a compendium for extracellular vesicles with continuous community annotation. *PLoS Biol.* *10*, e1001450.
- Kass, S., Craig, N., and Sollner-Webb, B. (1987). Primary processing of mammalian rRNA involves two adjacent cleavages and is not species specific. *Mol. Cell. Biol.* *7*, 2891–2898.
- Kass, S., Tyc, K., Steitz, J.A., and Sollner-Webb, B. (1990). The U3 small nucleolar ribonucleoprotein functions in the first step of preribosomal RNA processing. *Cell* *60*, 897–908.
- Kiss, A.M., Jády, B.E., Bertrand, E., and Kiss, T. (2004). Human box H/ACA pseudouridylation guide RNA machinery. *Mol. Cell. Biol.* *24*, 5797–5807.
- Kiss-László, Z., Henry, Y., Bachellerie, J.P., Caizergues-Ferrer, M., and Kiss, T. (1996). Site-specific ribose methylation of preribosomal RNA: a novel function for small nucleolar RNAs. *Cell* *85*, 1077–1088.

- Konishi, M., and Sugiyama, M. (2003). Genetic analysis of adventitious root formation with a novel series of temperature-sensitive mutants of *Arabidopsis thaliana*. *Dev. Camb. Engl.* *130*, 5637–5647.
- Kos, M., and Tollervey, D. (2010). Yeast pre-rRNA processing and modification occur cotranscriptionally. *Mol. Cell* *37*, 809–820.
- Kowal, J., Arras, G., Colombo, M., Jouve, M., Morath, J.P., Primdal-Bengtson, B., Dingli, F., Loew, D., Tkach, M., and Théry, C. (2016). Proteomic comparison defines novel markers to characterize heterogeneous populations of extracellular vesicle subtypes. *Proc. Natl. Acad. Sci. U. S. A.* *113*, E968–977.
- Krogh, N., Jansson, M.D., Häfner, S.J., Tehler, D., Birkedal, U., Christensen-Dalsgaard, M., Lund, A.H., and Nielsen, H. (2016). Profiling of 2'-O-Me in human rRNA reveals a subset of fractionally modified positions and provides evidence for ribosome heterogeneity. *Nucleic Acids Res.* *44*, 7884–7895.
- Kufel, J., Dichtl, B., and Tollervey, D. (1999). Yeast Rnt1p is required for cleavage of the pre-ribosomal RNA in the 3' ETS but not the 5' ETS. *RNA N. Y. N* *5*, 909–917.
- Lafontaine, D., Delcour, J., Glasser, A.L., Desgrès, J., and Vandenhoute, J. (1994). The DIM1 gene responsible for the conserved m6(2)Am6(2)A dimethylation in the 3'-terminal loop of 18 S rRNA is essential in yeast. *J. Mol. Biol.* *241*, 492–497.
- Lafontaine, D., Vandenhoute, J., and Tollervey, D. (1995). The 18S rRNA dimethylase Dim1p is required for pre-ribosomal RNA processing in yeast. *Genes Dev.* *9*, 2470–2481.
- Lafontaine, D.L., Bousquet-Antonelli, C., Henry, Y., Caizergues-Ferrer, M., and Tollervey, D. (1998a). The box H + ACA snoRNAs carry Cbf5p, the putative rRNA pseudouridine synthase. *Genes Dev.* *12*, 527–537.
- Lafontaine, D.L., Preiss, T., and Tollervey, D. (1998b). Yeast 18S rRNA dimethylase Dim1p: a quality control mechanism in ribosome synthesis? *Mol. Cell. Biol.* *18*, 2360–2370.
- Langhendries, J.-L., Nicolas, E., Doumont, G., Goldman, S., and Lafontaine, D.L.J. (2016). The human box C/D snoRNAs U3 and U8 are required for pre-rRNA processing and tumorigenesis. *Oncotarget*.
- Larburu, N., Montellese, C., O'Donohue, M.-F., Kutay, U., Gleizes, P.-E., and Plisson-Chastang, C. (2016). Structure of a human pre-40S particle points to a role for RACK1 in the final steps of 18S rRNA processing. *Nucleic Acids Res.* *44*, 8465–8478.
- Lebaron, S., Schneider, C., van Nues, R.W., Swiatkowska, A., Walsh, D., Böttcher, B., Granneman, S., Watkins, N.J., and Tollervey, D. (2012). Proofreading of pre-40S ribosome maturation by a translation initiation factor and 60S subunits. *Nat. Struct. Mol. Biol.* *19*, 744–753.
- Lestrade, L., and Weber, M.J. (2006). snoRNA-LBME-db, a comprehensive database of human H/ACA and C/D box snoRNAs. *Nucleic Acids Res.* *34*, D158–D162.
- Létoquart, J., Huvelle, E., Wacheul, L., Bourgeois, G., Zorbas, C., Graille, M., Heurgué-Hamard, V., and Lafontaine, D.L.J. (2014). Structural and functional studies of Bud23-Trm112 reveal 18S rRNA N7-G1575 methylation occurs on late 40S precursor ribosomes. *Proc. Natl. Acad. Sci. U. S. A.* *111*, E5518–5526.
- Létoquart, J., van Tran, N., Caroline, V., Aleksandrov, A., Lazar, N., van Tilbeurgh, H., Liger, D., and Graille, M. (2015). Insights into molecular plasticity in protein complexes from Trm9-Trm112 tRNA modifying enzyme crystal structure. *Nucleic Acids Res.* *43*, 10989–11002.

- Leulliot, N., Bohnsack, M.T., Graille, M., Tollervey, D., and Van Tilbeurgh, H. (2008). The yeast ribosome synthesis factor Emg1 is a novel member of the superfamily of alpha/beta knot fold methyltransferases. *Nucleic Acids Res.* *36*, 629–639.
- Liger, D., Mora, L., Lazar, N., Figaro, S., Henri, J., Scrima, N., Buckingham, R.H., van Tilbeurgh, H., Heurgué-Hamard, V., and Graille, M. (2011). Mechanism of activation of methyltransferases involved in translation by the Trm112 “hub” protein. *Nucleic Acids Res.* *39*, 6249–6259.
- Lin, A.E., Basson, C.T., Goldmuntz, E., Magoulas, P.L., McDermott, D.A., McDonald-McGinn, D.M., McPherson, E., Morris, C.A., Noonan, J., Nowak, C., et al. (2008). Adults with genetic syndromes and cardiovascular abnormalities: Clinical history and management. *Genet. Med. Off. J. Am. Coll. Med. Genet.* *10*, 469–494.
- Lygerou, Z., Allmang, C., Tollervey, D., and Séraphin, B. (1996). Accurate processing of a eukaryotic precursor ribosomal RNA by ribonuclease MRP in vitro. *Science* *272*, 268–270.
- Machnicka, M.A., Milanowska, K., Osman Oglou, O., Purta, E., Kurkowska, M., Olchowik, A., Januszewski, W., Kalinowski, S., Dunin-Horkawicz, S., Rother, K.M., et al. (2013). MODOMICS: a database of RNA modification pathways – 2013 update. *Nucleic Acids Res.* *41*, D262–267.
- Maden, B.E., and Hughes, J.M. (1997). Eukaryotic ribosomal RNA: the recent excitement in the nucleotide modification problem. *Chromosoma* *105*, 391–400.
- Marcel, V., Ghayad, S.E., Belin, S., Therizols, G., Morel, A.-P., Solano-González, E., Vendrell, J.A., Hacot, S., Mertani, H.C., Albaret, M.A., et al. (2013). p53 acts as a safeguard of translational control by regulating fibrillar and rRNA methylation in cancer. *Cancer Cell* *24*, 318–330.
- Marfori, M., Mynott, A., Ellis, J.J., Mehdi, A.M., Saunders, N.F.W., Curmi, P.M., Forwood, J.K., Bodén, M., and Kobe, B. (2011). Molecular basis for specificity of nuclear import and prediction of nuclear localization. *Biochim. Biophys. Acta* *1813*, 1562–1577.
- Marimpietri, D., Petretto, A., Raffaghello, L., Pezzolo, A., Gagliani, C., Tacchetti, C., Mauri, P., Melioli, G., and Pistoia, V. (2013). Proteome profiling of neuroblastoma-derived exosomes reveal the expression of proteins potentially involved in tumor progression. *PloS One* *8*, e75054.
- Martens-Uzunova, E.S., Jalava, S.E., Dits, N.F., van Leenders, G.J.L.H., Møller, S., Trapman, J., Bangma, C.H., Litman, T., Visakorpi, T., and Jenster, G. (2012). Diagnostic and prognostic signatures from the small non-coding RNA transcriptome in prostate cancer. *Oncogene* *31*, 978–991.
- Martin, J.L., and McMillan, F.M. (2002). SAM (dependent) I AM: the S-adenosylmethionine-dependent methyltransferase fold. *Curr. Opin. Struct. Biol.* *12*, 783–793.
- Mazaauric, M.-H., Dirick, L., Purushothaman, S.K., Björk, G.R., and Lapeyre, B. (2010). Trm112p is a 15-kDa zinc finger protein essential for the activity of two tRNA and one protein methyltransferases in yeast. *J. Biol. Chem.* *285*, 18505–18515.
- Merla, G., Ucla, C., Guipponi, M., and Reymond, A. (2002). Identification of additional transcripts in the Williams-Beuren syndrome critical region. *Hum. Genet.* *110*, 429–438.
- Merla, G., Howald, C., Henrichsen, C.N., Lyle, R., Wyss, C., Zobot, M.-T., Antonarakis, S.E., and Reymond, A. (2006). Submicroscopic deletion in patients with Williams-Beuren syndrome influences expression levels of the nonhemizygous flanking genes. *Am. J. Hum. Genet.* *79*, 332–341.

- Merla, G., Brunetti-Pierri, N., Micale, L., and Fusco, C. (2010). Copy number variants at Williams-Beuren syndrome 7q11.23 region. *Hum. Genet.* *128*, 3–26.
- Meyer, B., Wurm, J.P., Kötter, P., Leisegang, M.S., Schilling, V., Buchhaupt, M., Held, M., Bahr, U., Karas, M., Heckel, A., et al. (2011). The Bowen-Conradi syndrome protein Nep1 (Emg1) has a dual role in eukaryotic ribosome biogenesis, as an essential assembly factor and in the methylation of  $\Psi$ 1191 in yeast 18S rRNA. *Nucleic Acids Res.* *39*, 1526–1537.
- Meyer, B., Wurm, J.P., Sharma, S., Immer, C., Pogoryelov, D., Kötter, P., Lafontaine, D.L.J., Wöhnert, J., and Entian, K.-D. (2016). Ribosome biogenesis factor Tsr3 is the aminocarboxypropyl transferase responsible for 18S rRNA hypermodification in yeast and humans. *Nucleic Acids Res.* *44*, 4304–4316.
- Miller, O.L., and Beatty, B.R. (1969). Visualization of nucleolar genes. *Science* *164*, 955–957.
- Morello, L.G., Hesling, C., Coltri, P.P., Castilho, B.A., Rimokh, R., and Zanchin, N.I.T. (2011). The NIP7 protein is required for accurate pre-rRNA processing in human cells. *Nucleic Acids Res.* *39*, 648–665.
- Motorin, Y., and Helm, M. (2011). RNA nucleotide methylation. *Wiley Interdiscip. Rev. RNA* *2*, 611–631.
- Mougey, E.B., O'Reilly, M., Osheim, Y., Miller, O.L., Beyer, A., and Sollner-Webb, B. (1993). The terminal balls characteristic of eukaryotic rRNA transcription units in chromatin spreads are rRNA processing complexes. *Genes Dev.* *7*, 1609–1619.
- Mulcahy, L.A., Pink, R.C., and Carter, D.R.F. (2014). Routes and mechanisms of extracellular vesicle uptake. *J. Extracell. Vesicles* *3*.
- Mullineux, S.-T., and Lafontaine, D.L.J. (2012). Mapping the cleavage sites on mammalian pre-rRNAs: where do we stand? *Biochimie* *94*, 1521–1532.
- Murayama, A., Ohmori, K., Fujimura, A., Minami, H., Yasuzawa-Tanaka, K., Kuroda, T., Oie, S., Daitoku, H., Okuwaki, M., Nagata, K., et al. (2008). Epigenetic control of rDNA loci in response to intracellular energy status. *Cell* *133*, 627–639.
- Muriaux, D., Mirro, J., Nagashima, K., Harvin, D., and Rein, A. (2002). Murine Leukemia Virus Nucleocapsid Mutant Particles Lacking Viral RNA Encapsidate Ribosomes. *J. Virol.* *76*, 11405–11413.
- Nakazawa, Y., Arai, H., and Fujita, N. (2011). The novel metastasis promoter Merm1/Wbscr22 enhances tumor cell survival in the vasculature by suppressing Zacl/p53-dependent apoptosis. *Cancer Res.* *71*, 1146–1155.
- Ni, L., and Snyder, M. (2001). A genomic study of the bipolar bud site selection pattern in *Saccharomyces cerevisiae*. *Mol. Biol. Cell* *12*, 2147–2170.
- Ni, J., Tien, A.L., and Fournier, M.J. (1997). Small nucleolar RNAs direct site-specific synthesis of pseudouridine in ribosomal RNA. *Cell* *89*, 565–573.
- Niewmierzycka, A., and Clarke, S. (1999). S-Adenosylmethionine-dependent methylation in *Saccharomyces cerevisiae*. Identification of a novel protein arginine methyltransferase. *J. Biol. Chem.* *274*, 814–824.
- Oeffinger, M., Zenklusen, D., Ferguson, A., Wei, K.E., Hage, A.E., Tollervey, D., Chait, B.T., Singer, R.H., and Rout, M.P. (2009). Rrp17p Is a Eukaryotic Exonuclease Required for 5' End Processing of Pre-60S Ribosomal RNA. *Mol. Cell* *36*, 768–781.
- O'Farrell, H.C., Xu, Z., Culver, G.M., and Rife, J.P. (2008). Sequence and structural evolution of the KsgA/Dim1 methyltransferase family. *BMC Res. Notes* *1*, 108.

- Ohbayashi, I., Konishi, M., Ebine, K., and Sugiyama, M. (2011). Genetic identification of Arabidopsis RID2 as an essential factor involved in pre-rRNA processing. *Plant J. Cell Mol. Biol.* *67*, 49–60.
- Osheim, Y.N., French, S.L., Keck, K.M., Champion, E.A., Spasov, K., Dragon, F., Baserga, S.J., and Beyer, A.L. (2004). Pre-18S Ribosomal RNA Is Structurally Compacted into the SSU Processome Prior to Being Cleaved from Nascent Transcripts in *Saccharomyces cerevisiae*. *Mol. Cell* *16*, 943–954.
- Peifer, C., Sharma, S., Watzinger, P., Lamberth, S., Kötter, P., and Entian, K.-D. (2013). Yeast Rrp8p, a novel methyltransferase responsible for m1A 645 base modification of 25S rRNA. *Nucleic Acids Res.* *41*, 1151–1163.
- Pelava, A., Schneider, C., and Watkins, N.J. (2016). The importance of ribosome production, and the 5S RNP-MDM2 pathway, in health and disease. *Biochem. Soc. Trans.* *44*, 1086–1090.
- Pertschy, B., Schneider, C., Gnädig, M., Schäfer, T., Tollervey, D., and Hurt, E. (2009). RNA helicase Prp43 and its co-factor Pfa1 promote 20 to 18 S rRNA processing catalyzed by the endonuclease Nob1. *J. Biol. Chem.* *284*, 35079–35091.
- Petersen-Mahrt, S.K., Estmer, C., Ohrmalm, C., Matthews, D.A., Russell, W.C., and Akusjärvi, G. (1999). The splicing factor-associated protein, p32, regulates RNA splicing by inhibiting ASF/SF2 RNA binding and phosphorylation. *EMBO J.* *18*, 1014–1024.
- Petrossian, T.C., and Clarke, S.G. (2011). Uncovering the human methyltransferasome. *Mol. Cell. Proteomics MCP* *10*, M110.000976.
- Piekna-Przybylska, D., Decatur, W.A., and Fournier, M.J. (2008). The 3D rRNA modification maps database: with interactive tools for ribosome analysis. *Nucleic Acids Res.* *36*, D178–183.
- Pober, B.R. (2010). Williams–Beuren Syndrome. *N. Engl. J. Med.* *362*, 239–252.
- Pober, B.R., Johnson, M., and Urban, Z. (2008). Mechanisms and treatment of cardiovascular disease in Williams–Beuren syndrome. *J. Clin. Invest.* *118*, 1606–1615.
- Popowski, T., Vialard, F., Leroy, B., Bault, J.P., and Molina-Gomes, D. (2011). Williams–Beuren syndrome: the prenatal phenotype. *Am. J. Obstet. Gynecol.* *205*, e6–e8.
- Preti, M., O’Donohue, M.-F., Montel-Lehry, N., Bortolin-Cavaillé, M.-L., Choismel, V., and Gleizes, P.-E. (2013). Gradual processing of the ITS1 from the nucleolus to the cytoplasm during synthesis of the human 18S rRNA. *Nucleic Acids Res.* *41*, 4709–4723.
- Purushothaman, S.K., Bujnicki, J.M., Grosjean, H., and Lapeyre, B. (2005). Trm11p and Trm112p are both required for the formation of 2-methylguanosine at position 10 in yeast tRNA. *Mol. Cell. Biol.* *25*, 4359–4370.
- Ramesh, M., and Woolford, J.L. (2016). Eukaryote-specific rRNA expansion segments function in ribosome biogenesis. *RNA N. Y. N* *22*, 1153–1162.
- Raska, I. (2003). Oldies but goldies: searching for Christmas trees within the nucleolar architecture. *Trends Cell Biol.* *13*, 517–525.
- Rouquette, J., Choismel, V., and Gleizes, P.-E. (2005). Nuclear export and cytoplasmic processing of precursors to the 40S ribosomal subunits in mammalian cells. *EMBO J.* *24*, 2862–2872.
- Samarsky, D.A., Balakin, A.G., and Fournier, M.J. (1995). Characterization of three new snRNAs from *Saccharomyces cerevisiae*: snR34, snR35 and snR36. *Nucleic Acids Res.* *23*, 2548–2554.
- Sardana, R., and Johnson, A.W. (2012). The methyltransferase adaptor protein Trm112 is involved in biogenesis of both ribosomal subunits. *Mol. Biol. Cell* *23*, 4313–4322.

- Sardana, R., White, J.P., and Johnson, A.W. (2013). The rRNA methyltransferase Bud23 shows functional interaction with components of the SSU processome and RNase MRP. *RNA N. Y. N* *19*, 828–840.
- Sardana, R., Zhu, J., Gill, M., and Johnson, A.W. (2014). Physical and functional interaction between the methyltransferase Bud23 and the essential DEAH-box RNA helicase Ecm16. *Mol. Cell. Biol.* *34*, 2208–2220.
- Schillewaert, S., Wacheul, L., Lhomme, F., and Lafontaine, D.L.J. (2012). The evolutionarily conserved protein Las1 is required for pre-rRNA processing at both ends of ITS2. *Mol. Cell. Biol.* *32*, 430–444.
- Schilling, V., Peifer, C., Buchhaupt, M., Lamberth, S., Lioutikov, A., Rietschel, B., Kötter, P., and Entian, K.-D. (2012). Genetic interactions of yeast NEP1 (EMG1), encoding an essential factor in ribosome biogenesis. *Yeast Chichester Engl.* *29*, 167–183.
- Schossere, M., Minois, N., Angerer, T.B., Amring, M., Dellago, H., Harreither, E., Calle-Perez, A., Pircher, A., Gerstl, M.P., Pfeifenberger, S., et al. (2015). Methylation of ribosomal RNA by NSUN5 is a conserved mechanism modulating organismal lifespan. *Nat. Commun.* *6*, 6158.
- Schubert, C. (2009). The genomic basis of the Williams-Beuren syndrome. *Cell. Mol. Life Sci. CMLS* *66*, 1178–1197.
- Segura-Morales, C., Pescia, C., Chatellard-Causse, C., Sadoul, R., Bertrand, E., and Basyuk, E. (2005). Tsg101 and Alix interact with murine leukemia virus Gag and cooperate with Nedd4 ubiquitin ligases during budding. *J. Biol. Chem.* *280*, 27004–27012.
- Sharma, S., and Lafontaine, D.L.J. (2015). “View From A Bridge”: A New Perspective on Eukaryotic rRNA Base Modification. *Trends Biochem. Sci.* *40*, 560–575.
- Sharma, S., Yang, J., Watzinger, P., Kötter, P., and Entian, K.-D. (2013a). Yeast Nop2 and Rcm1 methylate C2870 and C2278 of the 25S rRNA, respectively. *Nucleic Acids Res.* *41*, 9062–9076.
- Sharma, S., Watzinger, P., Kötter, P., and Entian, K.-D. (2013b). Identification of a novel methyltransferase, Bmt2, responsible for the N-1-methyl-adenosine base modification of 25S rRNA in *Saccharomyces cerevisiae*. *Nucleic Acids Res.* *41*, 5428–5443.
- Sharma, S., Yang, J., Düttmann, S., Watzinger, P., Kötter, P., and Entian, K.-D. (2014). Identification of novel methyltransferases, Bmt5 and Bmt6, responsible for the m3U methylations of 25S rRNA in *Saccharomyces cerevisiae*. *Nucleic Acids Res.* *42*, 3246–3260.
- Sharma, S., Langhendries, J.-L., Watzinger, P., Kötter, P., Entian, K.-D., and Lafontaine, D.L.J. (2015). Yeast Kre33 and human NAT10 are conserved 18S rRNA cytosine acetyltransferases that modify tRNAs assisted by the adaptor Tan1/THUMPDI. *Nucleic Acids Res.* *43*, 2242–2258.
- Shen, Q., Zheng, X., McNutt, M.A., Guang, L., Sun, Y., Wang, J., Gong, Y., Hou, L., and Zhang, B. (2009). NAT10, a nucleolar protein, localizes to the midbody and regulates cytokinesis and acetylation of microtubules. *Exp. Cell Res.* *315*, 1653–1667.
- Sloan, K.E., Mattijssen, S., Lebaron, S., Tollervey, D., Pruijn, G.J.M., and Watkins, N.J. (2013). Both endonucleolytic and exonucleolytic cleavage mediate ITS1 removal during human ribosomal RNA processing. *J. Cell Biol.* *200*, 577–588.
- Sloan, K.E., Warda, A.S., Sharma, S., Entian, K.-D., Lafontaine, D.L.J., and Bohnsack, M.T. (2016). Tuning the ribosome: the influence of rRNA modification on eukaryotic ribosome biogenesis and function. *RNA Biol.* doi: 10.1080/15476286.2016.1259781

- Somerville, M.J., Mervis, C.B., Young, E.J., Seo, E.-J., del Campo, M., Bamforth, S., Peregrine, E., Loo, W., Lilley, M., Pérez-Jurado, L.A., et al. (2005). Severe expressive-language delay related to duplication of the Williams-Beuren locus. *N. Engl. J. Med.* *353*, 1694–1701.
- Sørensen, P.D., and Frederiksen, S. (1991). Characterization of human 5S rRNA genes. *Nucleic Acids Res.* *19*, 4147–4151.
- Stanchi, F., Bertocco, E., Toppo, S., Dioguardi, R., Simionati, B., Cannata, N., Zimbello, R., Lanfranchi, G., and Valle, G. (2001). Characterization of 16 novel human genes showing high similarity to yeast sequences. *Yeast Chichester Engl.* *18*, 69–80.
- Stankiewicz, P., and Lupski, J.R. (2002). Genome architecture, rearrangements and genomic disorders. *Trends Genet. TIG* *18*, 74–82.
- Stefanska, B., Cheishvili, D., Suderman, M., Arakelian, A., Huang, J., Hallett, M., Han, Z.-G., Al-Mahtab, M., Akbar, S.M.F., Khan, W.A., et al. (2014). Genome-wide study of hypomethylated and induced genes in patients with liver cancer unravels novel anticancer targets. *Clin. Cancer Res. Off. J. Am. Assoc. Cancer Res.* *20*, 3118–3132.
- Strømme, P., Bjørnstad, P.G., and Ramstad, K. (2002). Prevalence estimation of Williams syndrome. *J. Child Neurol.* *17*, 269–271.
- Studte, P., Zink, S., Jablonowski, D., Bär, C., von der Haar, T., Tuite, M.F., and Schaffrath, R. (2008). tRNA and protein methylase complexes mediate zymocin toxicity in yeast. *Mol. Microbiol.* *69*, 1266–1277.
- Su, H., Xu, T., Ganapathy, S., Shadfan, M., Long, M., Huang, T.H.-M., Thompson, I., and Yuan, Z.-M. (2014). Elevated snoRNA biogenesis is essential in breast cancer. *Oncogene* *33*, 1348–1358.
- Sumita, M., Desaulniers, J.-P., Chang, Y.-C., Chui, H.M.-P., Clos, L., and Chow, C.S. (2005). Effects of nucleotide substitution and modification on the stability and structure of helix 69 from 28S rRNA. *RNA N. Y. N* *11*, 1420–1429.
- Tafforeau, L., Zorbas, C., Langhendries, J.-L., Mullineux, S.-T., Stamatopoulou, V., Mullier, R., Wacheul, L., and Lafontaine, D.L.J. (2013). The complexity of human ribosome biogenesis revealed by systematic nucleolar screening of Pre-rRNA processing factors. *Mol. Cell* *51*, 539–551.
- Tam, E., Young, E.J., Morris, C.A., Marshall, C.R., Loo, W., Scherer, S.W., Mervis, C.B., and Osborne, L.R. (2008). The common inversion of the Williams-Beuren syndrome region at 7q11.23 does not cause clinical symptoms. *Am. J. Med. Genet. A.* *146A*, 1797–1806.
- Taoka, M., Nobe, Y., Yamaki, Y., Yamauchi, Y., Ishikawa, H., Takahashi, N., Nakayama, H., and Isobe, T. (2016). The complete chemical structure of *Saccharomyces cerevisiae* rRNA: partial pseudouridylation of U2345 in 25S rRNA by snoRNA snR9. *Nucleic Acids Res.* *44*, 8951–8961.
- Thiry, M., and Lafontaine, D.L.J. (2005). Birth of a nucleolus: the evolution of nucleolar compartments. *Trends Cell Biol.* *15*, 194–199.
- Thiry, M., Lamaye, F., and Lafontaine, D.L.J. (2011). The nucleolus: when 2 became 3. *Nucl. Austin Tex* *2*, 289–293.
- Thomson, E., and Tollervey, D. (2010). The final step in 5.8S rRNA processing is cytoplasmic in *Saccharomyces cerevisiae*. *Mol. Cell. Biol.* *30*, 976–984.
- Tiedemann, R.E., Zhu, Y.X., Schmidt, J., Shi, C.X., Sereduk, C., Yin, H., Mousses, S., and Stewart, A.K. (2012). Identification of molecular vulnerabilities in human multiple myeloma cells by RNA interference lethality screening of the druggable genome. *Cancer Res.* *72*, 757–768.

- Tollervey, D., Lehtonen, H., Jansen, R., Kern, H., and Hurt, E.C. (1993). Temperature-sensitive mutations demonstrate roles for yeast fibrillar in pre-rRNA processing, pre-rRNA methylation, and ribosome assembly. *Cell* 72, 443–457.
- Towns, W.L., and Begley, T.J. (2012). Transfer RNA methyltransferases and their corresponding modifications in budding yeast and humans: activities, predications, and potential roles in human health. *DNA Cell Biol.* 31, 434–454.
- Tyc, K., and Steitz, J.A. (1989). U3, U8 and U13 comprise a new class of mammalian snRNPs localized in the cell nucleolus. *EMBO J.* 8, 3113–3119.
- Uhlén, M., Fagerberg, L., Hallström, B.M., Lindskog, C., Oksvold, P., Mardinoglu, A., Sivertsson, Å., Kampf, C., Sjöstedt, E., Asplund, A., et al. (2015). Proteomics. Tissue-based map of the human proteome. *Science* 347, 1260419.
- Urbán, Z., Helms, C., Fekete, G., Csiszár, K., Bonnet, D., Munnich, A., Donis-Keller, H., and Boyd, C.D. (1996). 7q11.23 deletions in Williams syndrome arise as a consequence of unequal meiotic crossover. *Am. J. Hum. Genet.* 59, 958–962.
- Van Den Heuvel, E., Manders, E., Swillen, A., and Zink, I. (2016). Developmental trajectories of structural and pragmatic language skills in school-aged children with Williams syndrome. *J. Intellect. Disabil. Res.* 60, 903–919.
- Velleman, S.L., and Mervis, C.B. (2011). Children with 7q11.23 Duplication Syndrome: Speech, Language, Cognitive, and Behavioral Characteristics and their Implications for Intervention. *Perspect. Lang. Learn. Educ.* 18, 108–116.
- Venema, J., and Tollervey, D. (1999). Ribosome synthesis in *Saccharomyces cerevisiae*. *Annu. Rev. Genet.* 33, 261–311.
- Waku, T., Nakajima, Y., Yokoyama, W., Nomura, N., Kako, K., Kobayashi, A., Shimizu, T., and Fukamizu, A. (2016). NML-mediated rRNA base methylation links ribosomal subunit formation to cell proliferation in a p53-dependent manner. *J. Cell Sci.* 129, 2382–2393.
- Wang, M., and Pestov, D.G. (2011). 5'-end surveillance by Xrn2 acts as a shared mechanism for mammalian pre-rRNA maturation and decay. *Nucleic Acids Res.* 39, 1811–1822.
- Wang, T., Birsoy, K., Hughes, N.W., Krupczak, K.M., Post, Y., Wei, J.J., Lander, E.S., and Sabatini, D.M. (2015a). Identification and characterization of essential genes in the human genome. *Science* 350, 1096–1101.
- Wang, Y., Yue, D., Xiao, M., Qi, C., Chen, Y., Sun, D., Zhang, N., and Chen, R. (2015b). CIQBP negatively regulates the activation of oncoprotein YBX1 in the renal cell carcinoma as revealed by interactomics analysis. *J. Proteome Res.* 14, 804–813.
- Warda, A.S., Freytag, B., Haag, S., Sloan, K.E., Görlich, D., and Bohnsack, M.T. (2016). Effects of the Bowen-Conradi syndrome mutation in EMG1 on its nuclear import, stability and nucleolar recruitment. *Hum. Mol. Genet.*
- Watkins, N.J., and Bohnsack, M.T. (2012). The box C/D and H/ACA snoRNPs: key players in the modification, processing and the dynamic folding of ribosomal RNA. *Wiley Interdiscip. Rev. RNA* 3, 397–414.
- White, J., Li, Z., Sardana, R., Bujnicki, J.M., Marcotte, E.M., and Johnson, A.W. (2008). Bud23 methylates G1575 of 18S rRNA and is required for efficient nuclear export of pre-40S subunits. *Mol. Cell. Biol.* 28, 3151–3161.
- Williams, J.C., Barratt-Boyes, B.G., and Lowe, J.B. (1961). Supravalvular aortic stenosis. *Circulation* 24, 1311–1318.
- Willms, E., Johansson, H.J., Mäger, I., Lee, Y., Blomberg, K.E.M., Sadik, M., Alaarg, A., Smith, C.I.E., Lehtio, J., Andaloussi, S.E., et al. (2016). Cells release subpopulations of exosomes with distinct molecular and biological properties. *Sci. Rep.* 6, 22519.



- Wilson, D.N., and Doudna Cate, J.H. (2012). The structure and function of the eukaryotic ribosome. *Cold Spring Harb. Perspect. Biol.* 4.
- Woolford, J.L., and Baserga, S.J. (2013). Ribosome Biogenesis in the Yeast *Saccharomyces cerevisiae*. *Genetics* 195, 643–681.
- Wurm, J.P., Meyer, B., Bahr, U., Held, M., Frolow, O., Kötter, P., Engels, J.W., Heckel, A., Karas, M., Entian, K.-D., et al. (2010). The ribosome assembly factor Nep1 responsible for Bowen-Conradi syndrome is a pseudouridine-N1-specific methyltransferase. *Nucleic Acids Res.* 38, 2387–2398.
- Yáñez-Mó, M., Siljander, P.R.-M., Andreu, Z., Zavec, A.B., Borràs, F.E., Buzas, E.I., Buzas, K., Casal, E., Cappello, F., Carvalho, J., et al. (2015). Biological properties of extracellular vesicles and their physiological functions. *J. Extracell. Vesicles* 4.
- Yang, L., Song, T., Chen, L., Kabra, N., Zheng, H., Koomen, J., Seto, E., and Chen, J. (2013). Regulation of SirT1-nucleomethylin binding by rRNA coordinates ribosome biogenesis with nutrient availability. *Mol. Cell. Biol.* 33, 3835–3848.
- Yoshikawa, H., Komatsu, W., Hayano, T., Miura, Y., Homma, K., Izumikawa, K., Ishikawa, H., Miyazawa, N., Tachikawa, H., Yamauchi, Y., et al. (2011). Splicing factor 2-associated protein p32 participates in ribosome biogenesis by regulating the binding of Nop52 and fibrillarin to preribosome particles. *Mol. Cell. Proteomics MCP* 10, M110.006148.
- Yoshikawa, H., Ishikawa, H., Izumikawa, K., Miura, Y., Hayano, T., Isobe, T., Simpson, R.J., and Takahashi, N. (2015). Human nucleolar protein Nop52 (RRP1/NNP-1) is involved in site 2 cleavage in internal transcribed spacer 1 of pre-rRNAs at early stages of ribosome biogenesis. *Nucleic Acids Res.* 43, 5524–5536.
- Yusupova, G., and Yusupov, M. (2014). High-resolution structure of the eukaryotic 80S ribosome. *Annu. Rev. Biochem.* 83, 467–486.
- Zhang, X., Zhang, F., Guo, L., Wang, Y., Zhang, P., Wang, R., Zhang, N., and Chen, R. (2013). Interactome analysis reveals that C1QBP (complement component 1, q subcomponent binding protein) is associated with cancer cell chemotaxis and metastasis. *Mol. Cell. Proteomics MCP* 12, 3199–3209.
- Zorbas, C., Nicolas, E., Wacheul, L., Huvelle, E., Heurgué-Hamard, V., and Lafontaine, D.L.J. (2015). The human 18S rRNA base methyltransferases DIMT1L and WBSR22-TRMT112 but not rRNA modification are required for ribosome biogenesis. *Mol. Biol. Cell* 26, 2080–2095.

## ACKNOWLEDGEMENTS

First, I would like to express my sincere gratitude to my supervisors Reet and Ants Kurg. Thank you for your guidance in the lab and for helpful scientific and non-scientific discussions, patience, and encouragement throughout my studies.

I am also thankful to Mart Ustav for enabling me to participate in the IMGEMEL project, and to Jaanus Remme for reading and commenting on this dissertation. In addition, I appreciate valuable discussions with Aare Abroi and Ivar Ilves. Furthermore, I would like to express my gratitude towards Merike and Inge for their administrative support.

I would like to thank all co-authors of the papers included in this thesis, and all the former and current members of our research group, especially Piia Kivipõld, Liisi Võsa, Lilian Leetsi, Ly Käsper, and Eve Toomsoo. Liisi and Piia, you have been really good friends of mine and I am grateful for the hours we spent together both in and out of the lab. In addition, I appreciate the friendly atmosphere in the laboratory of molecular biology and the help provided by Aivar Liiv and Margus Leppik.

I would also like to thank the helpful members of the Ustav lab and other colleagues from the Institute of Technology. I am grateful to Age, Airiin, Ervin, Eva-Maria, Liisi Henno, Margit, Marit, Matu, Rait, Sandra, Sergo, Sirle, Tom and Tormi for memorable times. In addition, adventurous moments with Annika and Margit often helped to recharge the batteries.

I would like to thank my friends Risse and Mari for always being there for me. I am also thankful to Marko and Jaan for occasionally querying about the progress of my doctoral studies. Now, finally, I can answer that the work is complete and the dissertation is finished.

I would like to thank my mother and grandmother for their love and belief in me throughout my studies. I am thankful to Kersti, for being the best sister and a very close friend. Last but not least, the encouragement and love from my dear Tiit have been crucial.

## **PUBLICATIONS**

## CURRICULUM VITAE

**Name:** Kadri Õunap  
**Date of birth:** 13<sup>th</sup> of August 1986  
**Citizenship:** Estonian  
**Address:** Institute of Technology, Nooruse 1, 50411 Tartu, Estonia  
**E-mail:** kadri.ounap@ut.ee

### Education:

2010–... University of Tartu, PhD studies in Molecular and Cell Biology  
2008–2010 University of Tartu, MSc  
2005–2008 University of Tartu, BSc  
2002–2005 Hugo Treffner Gymnasium

### Professional employment:

2012–... University of Tartu, Institute of Technology, specialist

### Supervised theses:

Heilike Toming bachelor's thesis, 2015 (co-supervisor: Reet Kurg, PhD)  
Lilian Leetsi master's thesis, 2014 (co-supervisor: Reet Kurg, PhD)  
Maarja Matsoo master's thesis, 2014 (co-supervisor: Reet Kurg, PhD)  
Lilian Leetsi bachelor's thesis, 2012 (co-supervisor: Reet Kurg, PhD)

### List of publications:

**Õunap K**, Käsper L, Kurg A, Kurg R (2013). The human WBSCR22 protein is involved in the biogenesis of the 40S ribosomal subunits in mammalian cells. *PLoS ONE* 8(9): e75686. doi:10.1371/journal.pone.0075686

**Õunap K**, Leetsi L, Matsoo M, Kurg R (2015). The stability of ribosome biogenesis factor WBSCR22 is regulated by interaction with TRMT112 via ubiquitin-proteasome pathway. *PLoS ONE* 10(7): e0133841. doi:10.1371/journal.pone.0133841

Kurg R, Reinsalu O, Jagur S, **Õunap K**, Vösa L, Kasvandik S, Padari K, Gilde-mann K, Ustav M (2016). Biochemical and proteomic characterization of retrovirus Gag based microparticles carrying melanoma antigens. *Sci. Rep.* 6, 29425; doi: 10.1038/srep29425

## ELULOOKIRJELDUS

**Nimi:** Kadri Õunap  
**Sünniaeg:** 13. august 1986  
**Rahvus:** eestlane  
**Aadress:** Tartu Ülikooli Tehnoloogiainstituut, Nooruse 1, Tartu, 50411, Eesti  
**E-post:** kadri.ounap@ut.ee  
**Telefon:** +372 5221074

### Haridustee:

2010–... Tartu Ülikool, Molekulaar- ja Rakubioloogia, doktorant  
2008–2010 Tartu Ülikool, geenitehnoloogia, MSc  
2005–2008 Tartu Ülikool, geenitehnoloogia, BSc  
2002–2005 Hugo Treffneri Gümnaasium

### Erialane töökogemus:

2012–... Tartu Ülikool, Tehnoloogiainstituut, spetsialist

### Juhendatud lõputööd:

Heilike Toming bakalaureusetöö, 2015 (kaasjuhendaja: Reet Kurg, PhD)  
Lilian Leetsi magistritöö, 2014 (kaasjuhendaja: Reet Kurg, PhD)  
Maarja Matsoo magistritöö, 2014 (kaasjuhendaja: Reet Kurg, PhD)  
Lilian Leetsi bakalaureusetöö, 2012 (kaasjuhendaja: Reet Kurg, PhD)

### Publikatsioonide loetelu:

**Õunap K**, Käsper L, Kurg A, Kurg R (2013). The human WBSCR22 protein is involved in the biogenesis of the 40S ribosomal subunits in mammalian cells. PLoS ONE 8(9): e75686. doi:10.1371/journal.pone.0075686  
**Õunap K**, Leetsi L, Matsoo M, Kurg R (2015). The stability of ribosome biogenesis factor WBSCR22 is regulated by interaction with TRMT112 via ubiquitin-proteasome pathway. PLoS ONE 10(7): e0133841. doi:10.1371/journal.pone.0133841  
Kurg R, Reinsalu O, Jagur S, **Õunap K**, Võsa L, Kasvandik S, Padari K, Gilde-  
mann K, Ustav M (2016). Biochemical and proteomic characterization of  
retrovirus Gag based microparticles carrying melanoma antigens. Sci. Rep.  
6, 29425; doi: 10.1038/srep29425

## DISSERTATIONES BIOLOGICAE UNIVERSITATIS TARTUENSIS

1. **Toivo Maimets.** Studies of human oncoprotein p53. Tartu, 1991, 96 p.
2. **Enn K. Seppet.** Thyroid state control over energy metabolism, ion transport and contractile functions in rat heart. Tartu, 1991, 135 p.
3. **Kristjan Zobel.** Epifüütsete makrosamblike väärtus õhu saastuse indikaatoritena Hamar-Dobani boreaalsetes mägimetsades. Tartu, 1992, 131 lk.
4. **Andres Mäe.** Conjugal mobilization of catabolic plasmids by transposable elements in helper plasmids. Tartu, 1992, 91 p.
5. **Maia Kivisaar.** Studies on phenol degradation genes of *Pseudomonas* sp. strain EST 1001. Tartu, 1992, 61 p.
6. **Allan Nurk.** Nucleotide sequences of phenol degradative genes from *Pseudomonas* sp. strain EST 1001 and their transcriptional activation in *Pseudomonas putida*. Tartu, 1992, 72 p.
7. **Ülo Tamm.** The genus *Populus* L. in Estonia: variation of the species biology and introduction. Tartu, 1993, 91 p.
8. **Jaanus Remme.** Studies on the peptidyltransferase centre of the *E.coli* ribosome. Tartu, 1993, 68 p.
9. **Ülo Langel.** Galanin and galanin antagonists. Tartu, 1993, 97 p.
10. **Arvo Käär.** The development of an automatic online dynamic fluorescence-based pH-dependent fiber optic penicillin flowthrough biosensor for the control of the benzylpenicillin hydrolysis. Tartu, 1993, 117 p.
11. **Lilian Järvekülg.** Antigenic analysis and development of sensitive immunoassay for potato viruses. Tartu, 1993, 147 p.
12. **Jaak Palumets.** Analysis of phytomass partition in Norway spruce. Tartu, 1993, 47 p.
13. **Arne Sellin.** Variation in hydraulic architecture of *Picea abies* (L.) Karst. trees grown under different environmental conditions. Tartu, 1994, 119 p.
13. **Mati Reeben.** Regulation of light neurofilament gene expression. Tartu, 1994, 108 p.
14. **Urmas Tartes.** Respiration rhythms in insects. Tartu, 1995, 109 p.
15. **Ülo Puurand.** The complete nucleotide sequence and infections *in vitro* transcripts from cloned cDNA of a potato A potyvirus. Tartu, 1995, 96 p.
16. **Peeter Hõrak.** Pathways of selection in avian reproduction: a functional framework and its application in the population study of the great tit (*Parus major*). Tartu, 1995, 118 p.
17. **Erkki Truve.** Studies on specific and broad spectrum virus resistance in transgenic plants. Tartu, 1996, 158 p.
18. **Illar Pata.** Cloning and characterization of human and mouse ribosomal protein S6-encoding genes. Tartu, 1996, 60 p.
19. **Ülo Niinemets.** Importance of structural features of leaves and canopy in determining species shade-tolerance in temperature deciduous woody taxa. Tartu, 1996, 150 p.

20. **Ants Kurg.** Bovine leukemia virus: molecular studies on the packaging region and DNA diagnostics in cattle. Tartu, 1996, 104 p.
21. **Ene Ustav.** E2 as the modulator of the BPV1 DNA replication. Tartu, 1996, 100 p.
22. **Aksel Soosaar.** Role of helix-loop-helix and nuclear hormone receptor transcription factors in neurogenesis. Tartu, 1996, 109 p.
23. **Maido Remm.** Human papillomavirus type 18: replication, transformation and gene expression. Tartu, 1997, 117 p.
24. **Tiiu Kull.** Population dynamics in *Cypripedium calceolus* L. Tartu, 1997, 124 p.
25. **Kalle Olli.** Evolutionary life-strategies of autotrophic planktonic microorganisms in the Baltic Sea. Tartu, 1997, 180 p.
26. **Meelis Pärtel.** Species diversity and community dynamics in calcareous grassland communities in Western Estonia. Tartu, 1997, 124 p.
27. **Malle Leht.** The Genus *Potentilla* L. in Estonia, Latvia and Lithuania: distribution, morphology and taxonomy. Tartu, 1997, 186 p.
28. **Tanel Tenson.** Ribosomes, peptides and antibiotic resistance. Tartu, 1997, 80 p.
29. **Arvo Tuvikene.** Assessment of inland water pollution using biomarker responses in fish *in vivo* and *in vitro*. Tartu, 1997, 160 p.
30. **Urmas Saarma.** Tuning ribosomal elongation cycle by mutagenesis of 23S rRNA. Tartu, 1997, 134 p.
31. **Henn Ojaveer.** Composition and dynamics of fish stocks in the gulf of Riga ecosystem. Tartu, 1997, 138 p.
32. **Lembi Lõugas.** Post-glacial development of vertebrate fauna in Estonian water bodies. Tartu, 1997, 138 p.
33. **Margus Pooga.** Cell penetrating peptide, transportan, and its predecessors, galanin-based chimeric peptides. Tartu, 1998, 110 p.
34. **Andres Saag.** Evolutionary relationships in some cetrarioid genera (Lichenized Ascomycota). Tartu, 1998, 196 p.
35. **Aivar Liiv.** Ribosomal large subunit assembly *in vivo*. Tartu, 1998, 158 p.
36. **Tatjana Oja.** Isoenzyme diversity and phylogenetic affinities among the eurasian annual bromes (*Bromus* L., Poaceae). Tartu, 1998, 92 p.
37. **Mari Moora.** The influence of arbuscular mycorrhizal (AM) symbiosis on the competition and coexistence of calcareous grassland plant species. Tartu, 1998, 78 p.
38. **Olavi Kurina.** Fungus gnats in Estonia (*Diptera: Bolitophilidae, Keroplattidae, Macroceridae, Ditomyiidae, Diadocidiidae, Mycetophilidae*). Tartu, 1998, 200 p.
39. **Andrus Tasa.** Biological leaching of shales: black shale and oil shale. Tartu, 1998, 98 p.
40. **Arnold Kristjuhan.** Studies on transcriptional activator properties of tumor suppressor protein p53. Tartu, 1998, 86 p.
41. **Sulev Ingerpuu.** Characterization of some human myeloid cell surface and nuclear differentiation antigens. Tartu, 1998, 163 p.

42. **Veljo Kisand**. Responses of planktonic bacteria to the abiotic and biotic factors in the shallow lake Võrtsjärv. Tartu, 1998, 118 p.
43. **Kadri Põldmaa**. Studies in the systematics of hypomyces and allied genera (Hypocreales, Ascomycota). Tartu, 1998, 178 p.
44. **Markus Vetemaa**. Reproduction parameters of fish as indicators in environmental monitoring. Tartu, 1998, 117 p.
45. **Heli Talvik**. Prepatent periods and species composition of different *Oesophagostomum* spp. populations in Estonia and Denmark. Tartu, 1998, 104 p.
46. **Katrin Heinsoo**. Cuticular and stomatal antechamber conductance to water vapour diffusion in *Picea abies* (L.) karst. Tartu, 1999, 133 p.
47. **Tarmo Annilo**. Studies on mammalian ribosomal protein S7. Tartu, 1998, 77 p.
48. **Indrek Ots**. Health state indices of reproducing great tits (*Parus major*): sources of variation and connections with life-history traits. Tartu, 1999, 117 p.
49. **Juan Jose Cantero**. Plant community diversity and habitat relationships in central Argentina grasslands. Tartu, 1999, 161 p.
50. **Rein Kalamees**. Seed bank, seed rain and community regeneration in Estonian calcareous grasslands. Tartu, 1999, 107 p.
51. **Sulev Kõks**. Cholecystokinin (CCK) – induced anxiety in rats: influence of environmental stimuli and involvement of endopioid mechanisms and serotonin. Tartu, 1999, 123 p.
52. **Ebe Sild**. Impact of increasing concentrations of O<sub>3</sub> and CO<sub>2</sub> on wheat, clover and pasture. Tartu, 1999, 123 p.
53. **Ljudmilla Timofejeva**. Electron microscopical analysis of the synaptosomal complex formation in cereals. Tartu, 1999, 99 p.
54. **Andres Valkna**. Interactions of galanin receptor with ligands and G-proteins: studies with synthetic peptides. Tartu, 1999, 103 p.
55. **Taavi Virro**. Life cycles of planktonic rotifers in lake Peipsi. Tartu, 1999, 101 p.
56. **Ana Rebane**. Mammalian ribosomal protein S3a genes and intron-encoded small nucleolar RNAs U73 and U82. Tartu, 1999, 85 p.
57. **Tiina Tamm**. Cocksfoot mottle virus: the genome organisation and translational strategies. Tartu, 2000, 101 p.
58. **Reet Kurg**. Structure-function relationship of the bovine papilloma virus E2 protein. Tartu, 2000, 89 p.
59. **Toomas Kivisild**. The origins of Southern and Western Eurasian populations: an mtDNA study. Tartu, 2000, 121 p.
60. **Niilo Kaldalu**. Studies of the TOL plasmid transcription factor XylS. Tartu, 2000, 88 p.
61. **Dina Lepik**. Modulation of viral DNA replication by tumor suppressor protein p53. Tartu, 2000, 106 p.



62. **Kai Vellak**. Influence of different factors on the diversity of the bryophyte vegetation in forest and wooded meadow communities. Tartu, 2000, 122 p.
63. **Jonne Kotta**. Impact of eutrophication and biological invasions on the structure and functions of benthic macrofauna. Tartu, 2000, 160 p.
64. **Georg Martin**. Phytobenthic communities of the Gulf of Riga and the inner sea the West-Estonian archipelago. Tartu, 2000, 139 p.
65. **Silvia Sepp**. Morphological and genetical variation of *Alchemilla L.* in Estonia. Tartu, 2000. 124 p.
66. **Jaan Liira**. On the determinants of structure and diversity in herbaceous plant communities. Tartu, 2000, 96 p.
67. **Priit Zingel**. The role of planktonic ciliates in lake ecosystems. Tartu, 2001, 111 p.
68. **Tiit Teder**. Direct and indirect effects in Host-parasitoid interactions: ecological and evolutionary consequences. Tartu, 2001, 122 p.
69. **Hannes Kollist**. Leaf apoplastic ascorbate as ozone scavenger and its transport across the plasma membrane. Tartu, 2001, 80 p.
70. **Reet Marits**. Role of two-component regulator system PehR-PehS and extracellular protease PrtW in virulence of *Erwinia Carotovora* subsp. *Carotovora*. Tartu, 2001, 112 p.
71. **Vallo Tilgar**. Effect of calcium supplementation on reproductive performance of the pied flycatcher *Ficedula hypoleuca* and the great tit *Parus major*, breeding in Northern temperate forests. Tartu, 2002, 126 p.
72. **Rita Hõrak**. Regulation of transposition of transposon Tn4652 in *Pseudomonas putida*. Tartu, 2002, 108 p.
73. **Liina Eek-Piirsoo**. The effect of fertilization, mowing and additional illumination on the structure of a species-rich grassland community. Tartu, 2002, 74 p.
74. **Krõõt Aasamaa**. Shoot hydraulic conductance and stomatal conductance of six temperate deciduous tree species. Tartu, 2002, 110 p.
75. **Nele Ingerpuu**. Bryophyte diversity and vascular plants. Tartu, 2002, 112 p.
76. **Neeme Tõnisson**. Mutation detection by primer extension on oligonucleotide microarrays. Tartu, 2002, 124 p.
77. **Margus Pensa**. Variation in needle retention of Scots pine in relation to leaf morphology, nitrogen conservation and tree age. Tartu, 2003, 110 p.
78. **Asko Lõhmus**. Habitat preferences and quality for birds of prey: from principles to applications. Tartu, 2003, 168 p.
79. **Viljar Jaks**. p53 – a switch in cellular circuit. Tartu, 2003, 160 p.
80. **Jaana Männik**. Characterization and genetic studies of four ATP-binding cassette (ABC) transporters. Tartu, 2003, 140 p.
81. **Marek Sammul**. Competition and coexistence of clonal plants in relation to productivity. Tartu, 2003, 159 p.
82. **Ivar Ilves**. Virus-cell interactions in the replication cycle of bovine papillomavirus type 1. Tartu, 2003, 89 p.

83. **Andres Männik**. Design and characterization of a novel vector system based on the stable replicator of bovine papillomavirus type 1. Tartu, 2003, 109 p.
84. **Ivika Ostonen**. Fine root structure, dynamics and proportion in net primary production of Norway spruce forest ecosystem in relation to site conditions. Tartu, 2003, 158 p.
85. **Gudrun Veldre**. Somatic status of 12–15-year-old Tartu schoolchildren. Tartu, 2003, 199 p.
86. **Ülo Väli**. The greater spotted eagle *Aquila clanga* and the lesser spotted eagle *A. pomarina*: taxonomy, phylogeography and ecology. Tartu, 2004, 159 p.
87. **Aare Abroi**. The determinants for the native activities of the bovine papillomavirus type 1 E2 protein are separable. Tartu, 2004, 135 p.
88. **Tiina Kahre**. Cystic fibrosis in Estonia. Tartu, 2004, 116 p.
89. **Helen Orav-Kotta**. Habitat choice and feeding activity of benthic suspension feeders and mesograzers in the northern Baltic Sea. Tartu, 2004, 117 p.
90. **Maarja Öpik**. Diversity of arbuscular mycorrhizal fungi in the roots of perennial plants and their effect on plant performance. Tartu, 2004, 175 p.
91. **Kadri Tali**. Species structure of *Neotinea ustulata*. Tartu, 2004, 109 p.
92. **Kristiina Tambets**. Towards the understanding of post-glacial spread of human mitochondrial DNA haplogroups in Europe and beyond: a phylogeographic approach. Tartu, 2004, 163 p.
93. **Arvi Jõers**. Regulation of p53-dependent transcription. Tartu, 2004, 103 p.
94. **Lilian Kadaja**. Studies on modulation of the activity of tumor suppressor protein p53. Tartu, 2004, 103 p.
95. **Jaak Truu**. Oil shale industry wastewater: impact on river microbial community and possibilities for bioremediation. Tartu, 2004, 128 p.
96. **Maire Peters**. Natural horizontal transfer of the *pheBA* operon. Tartu, 2004, 105 p.
97. **Ülo Maiväli**. Studies on the structure-function relationship of the bacterial ribosome. Tartu, 2004, 130 p.
98. **Merit Otsus**. Plant community regeneration and species diversity in dry calcareous grasslands. Tartu, 2004, 103 p.
99. **Mikk Heidemaa**. Systematic studies on sawflies of the genera *Dolerus*, *Empria*, and *Caliroa* (Hymenoptera: Tenthredinidae). Tartu, 2004, 167 p.
100. **Ilmar Tõnno**. The impact of nitrogen and phosphorus concentration and N/P ratio on cyanobacterial dominance and N<sub>2</sub> fixation in some Estonian lakes. Tartu, 2004, 111 p.
101. **Lauri Saks**. Immune function, parasites, and carotenoid-based ornaments in greenfinches. Tartu, 2004, 144 p.
102. **Siiri Rootsi**. Human Y-chromosomal variation in European populations. Tartu, 2004, 142 p.
103. **Eve Vedler**. Structure of the 2,4-dichloro-phenoxyacetic acid-degradative plasmid pEST4011. Tartu, 2005. 106 p.

104. **Andres Tover.** Regulation of transcription of the phenol degradation *pheBA* operon in *Pseudomonas putida*. Tartu, 2005, 126 p.
105. **Helen Udras.** Hexose kinases and glucose transport in the yeast *Hansenula polymorpha*. Tartu, 2005, 100 p.
106. **Ave Suija.** Lichens and lichenicolous fungi in Estonia: diversity, distribution patterns, taxonomy. Tartu, 2005, 162 p.
107. **Piret Lõhmus.** Forest lichens and their substrata in Estonia. Tartu, 2005, 162 p.
108. **Inga Lips.** Abiotic factors controlling the cyanobacterial bloom occurrence in the Gulf of Finland. Tartu, 2005, 156 p.
109. **Kaasik, Krista.** Circadian clock genes in mammalian clockwork, metabolism and behaviour. Tartu, 2005, 121 p.
110. **Juhan Javoš.** The effects of experience on host acceptance in ovipositing moths. Tartu, 2005, 112 p.
111. **Tiina Sedman.** Characterization of the yeast *Saccharomyces cerevisiae* mitochondrial DNA helicase Hmi1. Tartu, 2005, 103 p.
112. **Ruth Agurauja.** Hawaiian endemic fern lineage *Diellia* (Aspleniaceae): distribution, population structure and ecology. Tartu, 2005, 112 p.
113. **Riho Teras.** Regulation of transcription from the fusion promoters generated by transposition of Tn4652 into the upstream region of *pheBA* operon in *Pseudomonas putida*. Tartu, 2005, 106 p.
114. **Mait Metspalu.** Through the course of prehistory in india: tracing the mtDNA trail. Tartu, 2005, 138 p.
115. **Elin Lõhmussaar.** The comparative patterns of linkage disequilibrium in European populations and its implication for genetic association studies. Tartu, 2006, 124 p.
116. **Priit Kopper.** Hydraulic and environmental limitations to leaf water relations in trees with respect to canopy position. Tartu, 2006, 126 p.
117. **Heili Ilves.** Stress-induced transposition of Tn4652 in *Pseudomonas Putida*. Tartu, 2006, 120 p.
118. **Silja Kuusk.** Biochemical properties of Hmi1p, a DNA helicase from *Saccharomyces cerevisiae* mitochondria. Tartu, 2006, 126 p.
119. **Kersti Püssa.** Forest edges on medium resolution landsat thematic mapper satellite images. Tartu, 2006, 90 p.
120. **Lea Tummeleht.** Physiological condition and immune function in great tits (*Parus major* L.): Sources of variation and trade-offs in relation to growth. Tartu, 2006, 94 p.
121. **Toomas Esperk.** Larval instar as a key element of insect growth schedules. Tartu, 2006, 186 p.
122. **Harri Valdmann.** Lynx (*Lynx lynx*) and wolf (*Canis lupus*) in the Baltic region: Diets, helminth parasites and genetic variation. Tartu, 2006. 102 p.
123. **Priit Jõers.** Studies of the mitochondrial helicase Hmi1p in *Candida albicans* and *Saccharomyces cerevisia*. Tartu, 2006. 113 p.
124. **Kersti Lilleväli.** Gata3 and Gata2 in inner ear development. Tartu, 2007, 123 p.

125. **Kai Rünk.** Comparative ecology of three fern species: *Dryopteris carthusiana* (Vill.) H.P. Fuchs, *D. expansa* (C. Presl) Fraser-Jenkins & Jermy and *D. dilatata* (Hoffm.) A. Gray (Dryopteridaceae). Tartu, 2007, 143 p.
126. **Aveliina Helm.** Formation and persistence of dry grassland diversity: role of human history and landscape structure. Tartu, 2007, 89 p.
127. **Leho Tedersoo.** Ectomycorrhizal fungi: diversity and community structure in Estonia, Seychelles and Australia. Tartu, 2007, 233 p.
128. **Marko Mägi.** The habitat-related variation of reproductive performance of great tits in a deciduous-coniferous forest mosaic: looking for causes and consequences. Tartu, 2007, 135 p.
129. **Valeria Lulla.** Replication strategies and applications of Semliki Forest virus. Tartu, 2007, 109 p.
130. **Ülle Reier.** Estonian threatened vascular plant species: causes of rarity and conservation. Tartu, 2007, 79 p.
131. **Inga Jüriado.** Diversity of lichen species in Estonia: influence of regional and local factors. Tartu, 2007, 171 p.
132. **Tatjana Krama.** Mobbing behaviour in birds: costs and reciprocity based cooperation. Tartu, 2007, 112 p.
133. **Signe Saumaa.** The role of DNA mismatch repair and oxidative DNA damage defense systems in avoidance of stationary phase mutations in *Pseudomonas putida*. Tartu, 2007, 172 p.
134. **Reedik Mägi.** The linkage disequilibrium and the selection of genetic markers for association studies in european populations. Tartu, 2007, 96 p.
135. **Priit Kilgas.** Blood parameters as indicators of physiological condition and skeletal development in great tits (*Parus major*): natural variation and application in the reproductive ecology of birds. Tartu, 2007, 129 p.
136. **Anu Albert.** The role of water salinity in structuring eastern Baltic coastal fish communities. Tartu, 2007, 95 p.
137. **Kärt Padari.** Protein transduction mechanisms of transportans. Tartu, 2008, 128 p.
138. **Siiri-Lii Sandre.** Selective forces on larval colouration in a moth. Tartu, 2008, 125 p.
139. **Ülle Jõgar.** Conservation and restoration of semi-natural floodplain meadows and their rare plant species. Tartu, 2008, 99 p.
140. **Lauri Laanisto.** Macroecological approach in vegetation science: generality of ecological relationships at the global scale. Tartu, 2008, 133 p.
141. **Reidar Andreson.** Methods and software for predicting PCR failure rate in large genomes. Tartu, 2008, 105 p.
142. **Birgot Paavel.** Bio-optical properties of turbid lakes. Tartu, 2008, 175 p.
143. **Kaire Torn.** Distribution and ecology of charophytes in the Baltic Sea. Tartu, 2008, 98 p.
144. **Vladimir Vimberg.** Peptide mediated macrolide resistance. Tartu, 2008, 190 p.
145. **Daima Örd.** Studies on the stress-inducible pseudokinase TRB3, a novel inhibitor of transcription factor ATF4. Tartu, 2008, 108 p.

146. **Lauri Saag.** Taxonomic and ecologic problems in the genus *Lepraria* (*Stereocaulaceae*, lichenised *Ascomycota*). Tartu, 2008, 175 p.
147. **Ulvi Karu.** Antioxidant protection, carotenoids and coccidians in greenfinches – assessment of the costs of immune activation and mechanisms of parasite resistance in a passerine with carotenoid-based ornaments. Tartu, 2008, 124 p.
148. **Jaanus Remm.** Tree-cavities in forests: density, characteristics and occupancy by animals. Tartu, 2008, 128 p.
149. **Epp Moks.** Tapeworm parasites *Echinococcus multilocularis* and *E. granulosus* in Estonia: phylogenetic relationships and occurrence in wild carnivores and ungulates. Tartu, 2008, 82 p.
150. **Eve Eensalu.** Acclimation of stomatal structure and function in tree canopy: effect of light and CO<sub>2</sub> concentration. Tartu, 2008, 108 p.
151. **Janne Pullat.** Design, functionlization and application of an *in situ* synthesized oligonucleotide microarray. Tartu, 2008, 108 p.
152. **Marta Putrinš.** Responses of *Pseudomonas putida* to phenol-induced metabolic and stress signals. Tartu, 2008, 142 p.
153. **Marina Semtšenko.** Plant root behaviour: responses to neighbours and physical obstructions. Tartu, 2008, 106 p.
154. **Marge Starast.** Influence of cultivation techniques on productivity and fruit quality of some *Vaccinium* and *Rubus* taxa. Tartu, 2008, 154 p.
155. **Age Tats.** Sequence motifs influencing the efficiency of translation. Tartu, 2009, 104 p.
156. **Radi Tegova.** The role of specialized DNA polymerases in mutagenesis in *Pseudomonas putida*. Tartu, 2009, 124 p.
157. **Tsipe Aavik.** Plant species richness, composition and functional trait pattern in agricultural landscapes – the role of land use intensity and landscape structure. Tartu, 2009, 112 p.
158. **Kaja Kiiver.** Semliki forest virus based vectors and cell lines for studying the replication and interactions of alphaviruses and hepaciviruses. Tartu, 2009, 104 p.
159. **Meelis Kadaja.** Papillomavirus Replication Machinery Induces Genomic Instability in its Host Cell. Tartu, 2009, 126 p.
160. **Pille Hallast.** Human and chimpanzee Luteinizing hormone/Chorionic Gonadotropin beta (*LHB/CGB*) gene clusters: diversity and divergence of young duplicated genes. Tartu, 2009, 168 p.
161. **Ain Vellak.** Spatial and temporal aspects of plant species conservation. Tartu, 2009, 86 p.
162. **Triinu Remmel.** Body size evolution in insects with different colouration strategies: the role of predation risk. Tartu, 2009, 168 p.
163. **Jaana Salujõe.** Zooplankton as the indicator of ecological quality and fish predation in lake ecosystems. Tartu, 2009, 129 p.
164. **Ele Vahtmäe.** Mapping benthic habitat with remote sensing in optically complex coastal environments. Tartu, 2009, 109 p.

165. **Liisa Metsamaa**. Model-based assessment to improve the use of remote sensing in recognition and quantitative mapping of cyanobacteria. Tartu, 2009, 114 p.
166. **Pille Säälük**. The role of endocytosis in the protein transduction by cell-penetrating peptides. Tartu, 2009, 155 p.
167. **Lauri Peil**. Ribosome assembly factors in *Escherichia coli*. Tartu, 2009, 147 p.
168. **Lea Hallik**. Generality and specificity in light harvesting, carbon gain capacity and shade tolerance among plant functional groups. Tartu, 2009, 99 p.
169. **Mariliis Tark**. Mutagenic potential of DNA damage repair and tolerance mechanisms under starvation stress. Tartu, 2009, 191 p.
170. **Riinu Rannap**. Impacts of habitat loss and restoration on amphibian populations. Tartu, 2009, 117 p.
171. **Maarja Adojaan**. Molecular variation of HIV-1 and the use of this knowledge in vaccine development. Tartu, 2009, 95 p.
172. **Signe Altmäe**. Genomics and transcriptomics of human induced ovarian folliculogenesis. Tartu, 2010, 179 p.
173. **Triin Suvi**. Mycorrhizal fungi of native and introduced trees in the Seychelles Islands. Tartu, 2010, 107 p.
174. **Velda Lauringson**. Role of suspension feeding in a brackish-water coastal sea. Tartu, 2010, 123 p.
175. **Eero Talts**. Photosynthetic cyclic electron transport – measurement and variably proton-coupled mechanism. Tartu, 2010, 121 p.
176. **Mari Nelis**. Genetic structure of the Estonian population and genetic distance from other populations of European descent. Tartu, 2010, 97 p.
177. **Kaarel Krjutškov**. Arrayed Primer Extension-2 as a multiplex PCR-based method for nucleic acid variation analysis: method and applications. Tartu, 2010, 129 p.
178. **Egle Köster**. Morphological and genetical variation within species complexes: *Anthyllis vulneraria* s. l. and *Alchemilla vulgaris* (coll.). Tartu, 2010, 101 p.
179. **Erki Õunap**. Systematic studies on the subfamily Sterrhinae (Lepidoptera: Geometridae). Tartu, 2010, 111 p.
180. **Merike Jõesaar**. Diversity of key catabolic genes at degradation of phenol and *p*-cresol in pseudomonads. Tartu, 2010, 125 p.
181. **Kristjan Herkül**. Effects of physical disturbance and habitat-modifying species on sediment properties and benthic communities in the northern Baltic Sea. Tartu, 2010, 123 p.
182. **Arto Pulk**. Studies on bacterial ribosomes by chemical modification approaches. Tartu, 2010, 161 p.
183. **Maria Põllupüü**. Ecological relations of cladocerans in a brackish-water ecosystem. Tartu, 2010, 126 p.
184. **Toomas Silla**. Study of the segregation mechanism of the Bovine Papillomavirus Type 1. Tartu, 2010, 188 p.

185. **Gyaneshwer Chaubey**. The demographic history of India: A perspective based on genetic evidence. Tartu, 2010, 184 p.
186. **Katrin Kepp**. Genes involved in cardiovascular traits: detection of genetic variation in Estonian and Czech populations. Tartu, 2010, 164 p.
187. **Virve Sõber**. The role of biotic interactions in plant reproductive performance. Tartu, 2010, 92 p.
188. **Kersti Kangro**. The response of phytoplankton community to the changes in nutrient loading. Tartu, 2010, 144 p.
189. **Joachim M. Gerhold**. Replication and Recombination of mitochondrial DNA in Yeast. Tartu, 2010, 120 p.
190. **Helen Tammert**. Ecological role of physiological and phylogenetic diversity in aquatic bacterial communities. Tartu, 2010, 140 p.
191. **Elle Rajandu**. Factors determining plant and lichen species diversity and composition in Estonian *Calamagrostis* and *Hepatica* site type forests. Tartu, 2010, 123 p.
192. **Paula Ann Kivistik**. ColR-ColS signalling system and transposition of Tn4652 in the adaptation of *Pseudomonas putida*. Tartu, 2010, 118 p.
193. **Siim Sõber**. Blood pressure genetics: from candidate genes to genome-wide association studies. Tartu, 2011, 120 p.
194. **Kalle Kipper**. Studies on the role of helix 69 of 23S rRNA in the factor-dependent stages of translation initiation, elongation, and termination. Tartu, 2011, 178 p.
195. **Triinu Siibak**. Effect of antibiotics on ribosome assembly is indirect. Tartu, 2011, 134 p.
196. **Tambet Tõnissoo**. Identification and molecular analysis of the role of guanine nucleotide exchange factor RIC-8 in mouse development and neural function. Tartu, 2011, 110 p.
197. **Helin Räägel**. Multiple faces of cell-penetrating peptides – their intracellular trafficking, stability and endosomal escape during protein transduction. Tartu, 2011, 161 p.
198. **Andres Jaanus**. Phytoplankton in Estonian coastal waters – variability, trends and response to environmental pressures. Tartu, 2011, 157 p.
199. **Tiit Nikopensius**. Genetic predisposition to nonsyndromic orofacial clefts. Tartu, 2011, 152 p.
200. **Signe Värv**. Studies on the mechanisms of RNA polymerase II-dependent transcription elongation. Tartu, 2011, 108 p.
201. **Kristjan Välk**. Gene expression profiling and genome-wide association studies of non-small cell lung cancer. Tartu, 2011, 98 p.
202. **Arno Põllumäe**. Spatio-temporal patterns of native and invasive zooplankton species under changing climate and eutrophication conditions. Tartu, 2011, 153 p.
203. **Egle Tammeleht**. Brown bear (*Ursus arctos*) population structure, demographic processes and variations in diet in northern Eurasia. Tartu, 2011, 143 p.

205. **Teele Jairus**. Species composition and host preference among ectomycorrhizal fungi in Australian and African ecosystems. Tartu, 2011, 106 p.
206. **Kessy Abarenkov**. PlutoF – cloud database and computing services supporting biological research. Tartu, 2011, 125 p.
207. **Marina Grigorova**. Fine-scale genetic variation of follicle-stimulating hormone beta-subunit coding gene (*FSHB*) and its association with reproductive health. Tartu, 2011, 184 p.
208. **Anu Tiitsaar**. The effects of predation risk and habitat history on butterfly communities. Tartu, 2011, 97 p.
209. **Elin Sild**. Oxidative defences in immunoeological context: validation and application of assays for nitric oxide production and oxidative burst in a wild passerine. Tartu, 2011, 105 p.
210. **Irja Saar**. The taxonomy and phylogeny of the genera *Cystoderma* and *Cystodermella* (Agaricales, Fungi). Tartu, 2012, 167 p.
211. **Pauli Saag**. Natural variation in plumage bacterial assemblages in two wild breeding passerines. Tartu, 2012, 113 p.
212. **Aleksei Lulla**. Alphaviral nonstructural protease and its polyprotein substrate: arrangements for the perfect marriage. Tartu, 2012, 143 p.
213. **Mari Järve**. Different genetic perspectives on human history in Europe and the Caucasus: the stories told by uniparental and autosomal markers. Tartu, 2012, 119 p.
214. **Ott Scheler**. The application of tmRNA as a marker molecule in bacterial diagnostics using microarray and biosensor technology. Tartu, 2012, 93 p.
215. **Anna Balikova**. Studies on the functions of tumor-associated mucin-like leukosialin (CD43) in human cancer cells. Tartu, 2012, 129 p.
216. **Triinu Kõressaar**. Improvement of PCR primer design for detection of prokaryotic species. Tartu, 2012, 83 p.
217. **Tuul Sepp**. Hematological health state indices of greenfinches: sources of individual variation and responses to immune system manipulation. Tartu, 2012, 117 p.
218. **Rya Ero**. Modifier view of the bacterial ribosome. Tartu, 2012, 146 p.
219. **Mohammad Bahram**. Biogeography of ectomycorrhizal fungi across different spatial scales. Tartu, 2012, 165 p.
220. **Annely Lorents**. Overcoming the plasma membrane barrier: uptake of amphipathic cell-penetrating peptides induces influx of calcium ions and downstream responses. Tartu, 2012, 113 p.
221. **Katrin Männik**. Exploring the genomics of cognitive impairment: whole-genome SNP genotyping experience in Estonian patients and general population. Tartu, 2012, 171 p.
222. **Marko Prouš**. Taxonomy and phylogeny of the sawfly genus *Empria* (Hymenoptera, Tenthredinidae). Tartu, 2012, 192 p.
223. **Triinu Visnapuu**. Levansucrases encoded in the genome of *Pseudomonas syringae* pv. tomato DC3000: heterologous expression, biochemical characterization, mutational analysis and spectrum of polymerization products. Tartu, 2012, 160 p.



224. **Nele Tamberg.** Studies on Semliki Forest virus replication and pathogenesis. Tartu, 2012, 109 p.
225. **Tõnu Esko.** Novel applications of SNP array data in the analysis of the genetic structure of Europeans and in genetic association studies. Tartu, 2012, 149 p.
226. **Timo Arula.** Ecology of early life-history stages of herring *Clupea harengus membras* in the northeastern Baltic Sea. Tartu, 2012, 143 p.
227. **Inga Hiiesalu.** Belowground plant diversity and coexistence patterns in grassland ecosystems. Tartu, 2012, 130 p.
228. **Kadri Koorem.** The influence of abiotic and biotic factors on small-scale plant community patterns and regeneration in boreonemoral forest. Tartu, 2012, 114 p.
229. **Liis Andresen.** Regulation of virulence in plant-pathogenic pectobacteria. Tartu, 2012, 122 p.
230. **Kaupo Kohv.** The direct and indirect effects of management on boreal forest structure and field layer vegetation. Tartu, 2012, 124 p.
231. **Mart Jüssi.** Living on an edge: landlocked seals in changing climate. Tartu, 2012, 114 p.
232. **Riina Klais.** Phytoplankton trends in the Baltic Sea. Tartu, 2012, 136 p.
233. **Rauno Veeroja.** Effects of winter weather, population density and timing of reproduction on life-history traits and population dynamics of moose (*Alces alces*) in Estonia. Tartu, 2012, 92 p.
234. **Marju Keis.** Brown bear (*Ursus arctos*) phylogeography in northern Eurasia. Tartu, 2013, 142 p.
235. **Sergei Põlme.** Biogeography and ecology of *alnus*- associated ectomycorrhizal fungi – from regional to global scale. Tartu, 2013, 90 p.
236. **Liis Uusküla.** Placental gene expression in normal and complicated pregnancy. Tartu, 2013, 173 p.
237. **Marko Lõoke.** Studies on DNA replication initiation in *Saccharomyces cerevisiae*. Tartu, 2013, 112 p.
238. **Anne Aan.** Light- and nitrogen-use and biomass allocation along productivity gradients in multilayer plant communities. Tartu, 2013, 127 p.
239. **Heidi Tamm.** Comprehending phylogenetic diversity – case studies in three groups of ascomycetes. Tartu, 2013, 136 p.
240. **Liina Kangur.** High-Pressure Spectroscopy Study of Chromophore-Binding Hydrogen Bonds in Light-Harvesting Complexes of Photosynthetic Bacteria. Tartu, 2013, 150 p.
241. **Margus Leppik.** Substrate specificity of the multisite specific pseudouridine synthase RluD. Tartu, 2013, 111 p.
242. **Lauris Kaplinski.** The application of oligonucleotide hybridization model for PCR and microarray optimization. Tartu, 2013, 103 p.
243. **Merli Pärnoja.** Patterns of macrophyte distribution and productivity in coastal ecosystems: effect of abiotic and biotic forcing. Tartu, 2013, 155 p.
244. **Tõnu Margus.** Distribution and phylogeny of the bacterial translational GTPases and the Mqsr/YgiT regulatory system. Tartu, 2013, 126 p.

245. **Pille Mänd.** Light use capacity and carbon and nitrogen budget of plants: remote assessment and physiological determinants. Tartu, 2013, 128 p.
246. **Mario Plaas.** Animal model of Wolfram Syndrome in mice: behavioural, biochemical and psychopharmacological characterization. Tartu, 2013, 144 p.
247. **Georgi Hudjašov.** Maps of mitochondrial DNA, Y-chromosome and tyrosinase variation in Eurasian and Oceanian populations. Tartu, 2013, 115 p.
248. **Mari Lepik.** Plasticity to light in herbaceous plants and its importance for community structure and diversity. Tartu, 2013, 102 p.
249. **Ede Leppik.** Diversity of lichens in semi-natural habitats of Estonia. Tartu, 2013, 151 p.
250. **Ülle Saks.** Arbuscular mycorrhizal fungal diversity patterns in boreo-nemoral forest ecosystems. Tartu, 2013, 151 p.
251. **Eneli Oitmaa.** Development of arrayed primer extension microarray assays for molecular diagnostic applications. Tartu, 2013, 147 p.
252. **Jekaterina Jutkina.** The horizontal gene pool for aromatics degradation: bacterial catabolic plasmids of the Baltic Sea aquatic system. Tartu, 2013, 121 p.
253. **Helen Vellau.** Reaction norms for size and age at maturity in insects: rules and exceptions. Tartu, 2014, 132 p.
254. **Randel Kreitsberg.** Using biomarkers in assessment of environmental contamination in fish – new perspectives. Tartu, 2014, 107 p.
255. **Krista Takkis.** Changes in plant species richness and population performance in response to habitat loss and fragmentation. Tartu, 2014, 141 p.
256. **Liina Nagirnaja.** Global and fine-scale genetic determinants of recurrent pregnancy loss. Tartu, 2014, 211 p.
257. **Triin Triisberg.** Factors influencing the re-vegetation of abandoned extracted peatlands in Estonia. Tartu, 2014, 133 p.
258. **Villu Soon.** A phylogenetic revision of the *Chrysis ignita* species group (Hymenoptera: Chrysididae) with emphasis on the northern European fauna. Tartu, 2014, 211 p.
259. **Andrei Nikonov.** RNA-Dependent RNA Polymerase Activity as a Basis for the Detection of Positive-Strand RNA Viruses by Vertebrate Host Cells. Tartu, 2014, 207 p.
260. **Eele Õunapuu-Pikas.** Spatio-temporal variability of leaf hydraulic conductance in woody plants: ecophysiological consequences. Tartu, 2014, 135 p.
261. **Marju Männiste.** Physiological ecology of greenfinches: information content of feathers in relation to immune function and behavior. Tartu, 2014, 121 p.
262. **Katre Kets.** Effects of elevated concentrations of CO<sub>2</sub> and O<sub>3</sub> on leaf photosynthetic parameters in *Populus tremuloides*: diurnal, seasonal and interannual patterns. Tartu, 2014, 115 p.

263. **Küllil Lokko**. Seasonal and spatial variability of zoopsammon communities in relation to environmental parameters. Tartu, 2014, 129 p.
264. **Olga Žilina**. Chromosomal microarray analysis as diagnostic tool: Estonian experience. Tartu, 2014, 152 p.
265. **Kertu Lõhmus**. Colonisation ecology of forest-dwelling vascular plants and the conservation value of rural manor parks. Tartu, 2014, 111 p.
266. **Anu Aun**. Mitochondria as integral modulators of cellular signaling. Tartu, 2014, 167 p.
267. **Chandana Basu Mallick**. Genetics of adaptive traits and gender-specific demographic processes in South Asian populations. Tartu, 2014, 160 p.
268. **Riin Tamme**. The relationship between small-scale environmental heterogeneity and plant species diversity. Tartu, 2014, 130 p.
269. **Liina Remm**. Impacts of forest drainage on biodiversity and habitat quality: implications for sustainable management and conservation. Tartu, 2015, 126 p.
270. **Tiina Talve**. Genetic diversity and taxonomy within the genus *Rhinanthus*. Tartu, 2015, 106 p.
271. **Mehis Rohtla**. Otolith sclerochronological studies on migrations, spawning habitat preferences and age of freshwater fishes inhabiting the Baltic Sea. Tartu, 2015, 137 p.
272. **Alexey Reshchikov**. The world fauna of the genus *Lathrolestes* (Hymenoptera, Ichneumonidae). Tartu, 2015, 247 p.
273. **Martin Pook**. Studies on artificial and extracellular matrix protein-rich surfaces as regulators of cell growth and differentiation. Tartu, 2015, 142 p.
274. **Mai Kukumägi**. Factors affecting soil respiration and its components in silver birch and Norway spruce stands. Tartu, 2015, 155 p.
275. **Helen Karu**. Development of ecosystems under human activity in the North-East Estonian industrial region: forests on post-mining sites and bogs. Tartu, 2015, 152 p.
276. **Hedi Peterson**. Exploiting high-throughput data for establishing relationships between genes. Tartu, 2015, 186 p.
277. **Priit Adler**. Analysis and visualisation of large scale microarray data, Tartu, 2015, 126 p.
278. **Aigar Niglas**. Effects of environmental factors on gas exchange in deciduous trees: focus on photosynthetic water-use efficiency. Tartu, 2015, 152 p.
279. **Silja Laht**. Classification and identification of conopeptides using profile hidden Markov models and position-specific scoring matrices. Tartu, 2015, 100 p.
280. **Martin Kesler**. Biological characteristics and restoration of Atlantic salmon *Salmo salar* populations in the Rivers of Northern Estonia. Tartu, 2015, 97 p.
281. **Pratyush Kumar Das**. Biochemical perspective on alphaviral nonstructural protein 2: a tale from multiple domains to enzymatic profiling. Tartu, 2015, 205 p.

282. **Priit Palta**. Computational methods for DNA copy number detection. Tartu, 2015, 130 p.
283. **Julia Sidorenko**. Combating DNA damage and maintenance of genome integrity in pseudomonads. Tartu, 2015, 174 p.
284. **Anastasiia Kovtun-Kante**. Charophytes of Estonian inland and coastal waters: distribution and environmental preferences. Tartu, 2015, 97 p.
285. **Ly Lindman**. The ecology of protected butterfly species in Estonia. Tartu, 2015, 171 p.
286. **Jaanis Lodjak**. Association of Insulin-like Growth Factor I and Corticosterone with Nestling Growth and Fledging Success in Wild Passerines. Tartu, 2016, 113 p.
287. **Ann Kraut**. Conservation of Wood-Inhabiting Biodiversity – Semi-Natural Forests as an Opportunity. Tartu, 2016, 141 p.
288. **Tiit Örd**. Functions and regulation of the mammalian pseudokinase TRIB3. Tartu, 2016, 182. p.
289. **Kairi Käiro**. Biological Quality According to Macroinvertebrates in Streams of Estonia (Baltic Ecoregion of Europe): Effects of Human-induced Hydromorphological Changes. Tartu, 2016, 126 p.
290. **Leidi Laurimaa**. *Echinococcus multilocularis* and other zoonotic parasites in Estonian canids. Tartu, 2016, 144 p.
291. **Helerin Margus**. Characterization of cell-penetrating peptide/nucleic acid nanocomplexes and their cell-entry mechanisms. Tartu, 2016, 173 p.
292. **Kadri Runnel**. Fungal targets and tools for forest conservation. Tartu, 2016, 157 p.
293. **Urmo Võsa**. MicroRNAs in disease and health: aberrant regulation in lung cancer and association with genomic variation. Tartu, 2016, 163 p.
294. **Kristina Mäemets-Allas**. Studies on cell growth promoting AKT signaling pathway – a promising anti-cancer drug target. Tartu, 2016, 146 p.
295. **Janeli Viil**. Studies on cellular and molecular mechanisms that drive normal and regenerative processes in the liver and pathological processes in Dupuytren's contracture. Tartu, 2016, 175 p.
296. **Ene Kook**. Genetic diversity and evolution of *Pulmonaria angustifolia* L. and *Myosotis laxa sensu lato* (Boraginaceae). Tartu, 2016, 106 p.
297. **Kadri Peil**. RNA polymerase II-dependent transcription elongation in *Saccharomyces cerevisiae*. Tartu, 2016, 113 p.
298. **Katrin Ruisu**. The role of RIC8A in mouse development and its function in cell-matrix adhesion and actin cytoskeletal organisation. Tartu, 2016, 129 p.
299. **Janely Pae**. Translocation of cell-penetrating peptides across biological membranes and interactions with plasma membrane constituents. Tartu, 2016, 126 p.
300. **Argo Ronk**. Plant diversity patterns across Europe: observed and dark diversity. Tartu, 2016, 153 p.

301. **Kristiina Mark.** Diversification and species delimitation of lichenized fungi in selected groups of the family Parmeliaceae (Ascomycota). Tartu, 2016, 181 p.
302. **Jaak-Albert Metsoja.** Vegetation dynamics in floodplain meadows: influence of mowing and sediment application. Tartu, 2016, 140 p.
303. **Hedvig Tamman.** The GraTA toxin-antitoxin system of *Pseudomonas putida*: regulation and role in stress tolerance. Tartu, 2016, 154 p.
304. **Kadri Pärtel.** Application of ultrastructural and molecular data in the taxonomy of helotialean fungi. Tartu, 2016, 183 p.
305. **Maris Hindrikson.** Grey wolf (*Canis lupus*) populations in Estonia and Europe: genetic diversity, population structure and -processes, and hybridization between wolves and dogs. Tartu, 2016, 121 p.
306. **Polina Degtjarenko.** Impacts of alkaline dust pollution on biodiversity of plants and lichens: from communities to genetic diversity. Tartu, 2016, 126 p.
307. **Liina Pajusalu.** The effect of CO<sub>2</sub> enrichment on net photosynthesis of macrophytes in a brackish water environment. Tartu, 2016, 126 p.
308. **Stoyan Tankov.** Random walks in the stringent response. Tartu, 2016, 94 p.
309. **Liis Leitsalu.** Communicating genomic research results to population-based biobank participants. Tartu, 2016, 158 p.
310. **Richard Meitern.** Redox physiology of wild birds: validation and application of techniques for detecting oxidative stress. Tartu, 2016, 134 p.
311. **Kaie Lokk.** Comparative genome-wide DNA methylation studies of healthy human tissues and non-small cell lung cancer tissue. Tartu, 2016, 127 p.
312. **Mihhail Kurašin.** Processivity of cellulases and chitinases. Tartu, 2017, 132 p.
313. **Carmen Tali.** Scavenger receptors as a target for nucleic acid delivery with peptide vectors. Tartu, 2017, 155 p.
314. **Katarina Oganjan.** Distribution, feeding and habitat of benthic suspension feeders in a shallow coastal sea. Tartu, 2017, 132 p.
315. **Taavi Paal.** Immigration limitation of forest plants into wooded landscape corridors. Tartu, 2017, 145 p.